Fig. 1. The mean tongue flick-attack score for each stimulus item.

significant ($p < 0.01$, two-tailed). The response rate was higher for the first five trials.

The total number of responses for the 10 trials given each S also varied considerably. The range was from 63 to 1063. Because of this variability, the data were normalized. A single S's score for each stimulus item was transformed to a per cent of that S's total score. The data were then analyzed by Friedman's two-way analysis of variance (Siegel, 1956), a nonparametric test for related samples. A first test was performed across all stimulus items. The difference was significant ($p < 0.05$). A second test was performed across all stimulus items with the exception of the control stimulus (distilled water). The difference was not significant ($0.50 < p < 0.70$). It was concluded that the response to the prey-object extracts did not differ among themselves, and that the

response to the prey-objects was greater than the response to the control stimulus.

Eight days after initial testing, one live baby deer mouse was introduced into each jar containing two Ss. The mouse was attacked by one of the Ss within 30 sec, in all cases. In three of the six jars, both of the Ss attempted to eat the young mouse.

DISCUSSION

The stimulus items can be separated into several categories: (1) prey frequently eaten by adult snakes but too large for juveniles, i.e., cotton rat, lab mouse, deer mouse, and chicken; (2) prey frequently eaten by adults and small enough for juveniles, i.e., baby deer mouse; and (3) prey not usually eaten by *Elaphe* yet small enough to be eaten by juveniles, i.e., grasshopper, fish, frog, and worm.

Baby deer mice, which were accepted quite readily as a first meal, had a rather low

mean attack score in comparison to the other prey-object extracts. This suggests that if a "low" attack score is correlated with a food animal that was accepted, any stimulus receiving a higher attack score would also be accepted. Considering Burghardt's evaluation of the significance of an attack score, these results imply that a newborn *Elaphe* confronted with an adult cotton rat will attempt to eat it. It follows that all prey of the first category would be acceptable targets for the young rat snake. This conclusion seems unlikely. It seems equally improbable that a 6-ft adult *Elaphe* would waste its time devouring the excessive numbers of worms, grasshoppers, or fish required to satiate it.

The data indicate that young rat snakes enter the world accepting a wide variety of prey objects—those normally eaten by the adult and prey of a convenient size. Factors that may inhibit young snakes from attacking animals too large to be eaten remain undetermined. It is the authors' opinion, however, that size and/or the behavior of prey-objects would be a profitable variable to investigate.

REFERENCES

- BURGHARDT, G. M. Stimulus control of the prey attack response in naive garter snakes. *Psychonomic Science*, 1966, 4, 37-38.
- BURGHARDT, G. M. Chemical-cue preferences of inexperienced snakes: Comparative aspects. *Science*, 1967a, 157, 718-721.
- BURGHARDT, G. M. Chemical perception in newborn snakes. *Psychology Today*, 1967b, August, 51-59.
- SIEGEL, S. *Nonparametric statistics for the behavioral sciences*. New York: McGraw-Hill, 1956. Pp. 83-87, 166-172.

NOTES

1. Work supported by the U.S. Atomic Energy Commission.
2. Summer student aide, now a graduate student in the Department of Psychology at Florida State University, Tallahassee, Florida.

ERRATUM

DAVIS, JOEL L., & THOMPSON, RICHARD F. Sensory preconditioning of cats in a shuttle box avoidance situation. *Psychonomic Science*, 1968, 13 (1), 37-38.—Due to an oversight by the authors, an error was made in the above paper. The second sentence of the second paragraph in the methods section should read "During this time the auditory stimulus was presented for 6 sec with the light presented contiguously for the last 2 sec . . ." The italicized terms were incorrectly interchanged in the original paper. It, of course, would have been a "backward" sensory preconditioning situation, which was not the case. The authors are indebted to Dr. M. E. Bitterman for calling the error to their attention.