volume which had been consumed was recorded. Plain tap water and quinine water test days were alternated; solutions of 0.15 mg and 0.25 mg of quinine per cc of water gave similar results which were pooled.

Results and Discussion

The results are given as Fig. 1. Thirsty hyperphagics and normals drink the same amount of tap water, but hyperphagics drink very little quinine water compared to normal controls.²

The percent decrement in consumption due to adding quinine to the drinking water was computed for each rat and a t test done to compare the groups; $t = 6.26$ with 20 degrees of freedom ($p < .001$).

Thus, hyperphagics will tolerate quinine in neither food nor water.

It is often observed that ventromedial rats are sluggish (Hetherington & Ranson, 1942) and "hyperemotional" (Anand & Brobeck, 1951; Hetherington & Ranson, 1942); these symptoms, as well as lack of tolerance for bad taste, may be non drive-specific dysfunctions which are produced in addition to (but independently of) increased hunger. In any case the present results further emphasize that ventromedially lesioned rats are abnormal in many ways and not only in their regulation of food intake.

Comment on Jacobson et al by David J. Barker

Jacobson et al (1966) report data from an experiment in which RNA from rats rewarded for choosing one or the other side of a two-choice apparatus was injected into naive rats, whose side "preferences" in the apparatus were then measured. Table 1 of their data shows "preferences" of test animals as indexed by their "predominant" choices on 25 test trials, regardless of the strength of "preference." The authors, however, do not make explicit what they mean by a predominant choice. Examination of Table 1 shows that of the 42 test rats, 29 chose the alternative to which their donor rats had been trained (a+d diagonal). Referring now to Table 2, the distribution of scores of the test rats, it is clear that these 29 rats must have come from the upper half of the distribution which includes the scores from 13 to 21. Thus, one would assume that the definition of a "predominant" choice was selecting the alternative to which the donor had been trained 13 or more times out of 25. Under the assumption that the injection of RNA from a donor rat into a recipient rat did not influence the recipient's choices, that its choices were random, the probability of 13 or more choices of the donor-trained alternative out of 25 is quite high ($p = .50$). Thus, while the fact that 29 out of 42 rats showed a "preference" for the side to which their donor had been trained is statistically significant ($p = .01$), it is highly likely that many of these "preferences" were random choices, and not due to the injection of RNA. If the criterion for "predominant" choice had been 14 out of 25 instead of 13 out of 25, the number of rats choosing the alternative to which their

References

- Anand, B. K., & Brobeck, J. R. Hypothalamic control of food intake in rats and cats. *Yale J. Biol. Med.*, 1951, 24, 123-140.
- Brobeck, J. R. Regulation of feeding and drinking. In J. Field (Ed.), *Handbook of physiology*, Washington, D. C.: American Physiological Soc., 1960. Section 1, Chap. 48.
- deGroot, J. The rat hypothalamus in stereotaxic coordinates. *J. comp. Neurol.*, 1959, 113, 389-400.
- Graff, H., & Stellar, E. Hyperphagia, obesity, and finickiness. *J. comp. physiol. Psychol.*, 1962, 55, 418-24.
- Hetherington, A. W., & Ranson, S. W. The spontaneous activity and food intake of rats with hypothalamic lesions. *Amer. J. Physiol.*, 1942, 136, 609.
- Miller, N. E. Motivational effects of brain stimulation and drugs. *Fed. Proc.*, 1960, 19, 846-854.
- Miller, N. E., Bailey, C. J., & Stevenson, J. A. Decreased "hunger" but increased food intake resulting from hypothalamic lesions. *Science*, 1950, 112, 256-259.
- Stevenson, J. A. F., Walt, L. G., & Orloff, J. Abnormalities of water and electrolyte metabolism in rats with hypothalamic lesions. *Amer. J. Physiol.*, 1950, 161, 35-39.

Notes

1. The work reported here was done at Yale University during the tenure of a National Science Foundation Predoctoral Fellowship. It was supported by Grant M647 from the National Institute of Mental Health, United States Public Health Service. The writer wishes to thank Professor Neal E. Miller for helpful suggestions.
2. The present lack of a between-group difference in tap water consumption should be compared with results reported in Stevenson et al (1950).

donor had been trained would have been 21 out of 42, a result which is not significant ($p = .50$). It would seem that in such an important and controversial research area, a more conservative definition of "predominant" choice should have been adopted, with strength of preference taken into account. This seems especially important in view of the fact that Gross & Carey (1965) twice failed to replicate the Babich et al (1965) study of the transfer of a response by injection of RNA.

The authors conclude from their results that "... specific training given to donor rats was a determinant of the subsequent choices of recipient rats." This strong conclusion does not follow from the results presented. All that a significantly large Chi square implies, in this case, is a lack of independence between "preference of injected rats" and "side to which donor was trained." It does not indicate that one determined the other. A quote from Hays (1963) may be appropriate here: "Given a large enough sample size, the chances are very good that the psychologist can demonstrate the association of any two qualitative attributes via a Chi square test."

References

- Babich, F. R., Jacobson, A. L., Bubash, S., & Jacobson, Ann. Transfer of a response to naive rats by injection of ribonucleic acid extracted from trained rats. *Science*, 1965, 149, 656-657.
- Gross, C., & Carey, F. Transfer of learned response by RNA injections: Failure of attempts to replicate. *Science*, 1965, 150, 1749.
- Hays, W. *Statistics for psychologists*. New York: Holt, Rinehart, & Winston, 1963. P. 613.
- Jacobson, A. L., Babich, F. R., Bubash, S., & Goren, C. Maze preferences in naive rats produced by injection of ribonucleic acid from trained rats. *Psychon. Sci.*, 1966, 4, 3-4.

For Reply to David J. Barker see page 334.

of pigeon IV may have been caused by erratic behavior which was exhibited throughout the experiment.

Discussion

Three things may be concluded from the data: first, that pigeons are capable of forming associations between colors; second, that these pigeons formed a backward association; and third, that the forward association was not significantly stronger than the backward association.

It is likely that these results are related to the manner of training; specifically, requiring the pigeon to peck at the stimulus color before pecking at the response color. In terms of the Asch-Ebenholtz studies, this procedure may help to equalize the "availability" of the S- and R-colors as contrasted to a procedure which requires pecking at R only. The present procedure was adopted in the belief that it would enhance the likelihood that the pigeons would pay attention to the S-color. Using a procedure which requires pecking the R-color only might result in the backward association being significantly weaker than the forward association.

It could be argued that the blackout after an incorrect response in backward testing strengthened the backward association. If the blackout explanation is correct, then the number of correct responses in the last half of the tests should be greater than those in the first half. This explanation is not supported by the data; the total number of correct pecks in the first 14 testing trials

over all pigeons equaled the total correct in the last 14 trials. The same result was also true for the first versus the last 10 trials, as well as for the first versus the last 5 trials. Therefore, it would appear that the blackout during testing did not strengthen the backward association.

It should be noted that the results of this study are based upon intra-individual performance and not upon comparisons between pigeons. Therefore, the loss of pigeon II and the erratic behavior of pigeon IV do not destroy the conclusion drawn from the experiment.

The capacity to form backward associations may be used to evaluate differences in intelligence or learning ability, in much the same way that Bitterman (1965) has used the tasks of habit reversal and probability learning for evolutionary comparisons of learning ability.

References

- Asch, S., & Ebenholtz, S. The principle of associative symmetry. *Proc. Amer. Phil. Soc.*, 1962, 106, 135-163.
Bitterman, M. E. Phyletic differences in learning. *Amer. Psychol.*, 1965, 20, 396-410.

Note

1. This study started as a Science Fair project at Martin Junior High School, Raleigh, North Carolina, in 1962. I am indebted to Norman Guttman and Slater E. Newman for advice and to Gilbert Gottlieb and my father, Clifton Gray, for guidance. I am also indebted to Ray Moore for instruction and help in circuitry design, to Connecticut College for shop facilities and laboratory space, and to my father for statistical analysis. I also appreciate the secretarial help of the North Carolina Department of Mental Health.

Reply to David J. Barker by Allan L. Jacobson

Barker is correct in concluding that by "predominant choice," we meant the alternative chosen 13 or more times on the 25 test trials. We did not intend to suggest anything else, and in fact, we pointed out in the text that "absolute preferences were on the whole small."

As for the statistical test; the approximate number of Ss and the way we would treat the data were decided in advance. We did not expect to obtain strong preferences,

even if our results were positive, since (among other reasons) recipient rats were never rewarded during testing; thus we chose a "sensitive" criterion of preference. Perhaps in so doing, we inadvertently capitalized on the vagaries of χ^2 . If so, Dr. Barker is indeed justified in considering our conclusion too strongly stated. We hope at some future time to resolve this question more satisfactorily in the laboratory.

For Comment on Jacobson et al see page 314.