Fig. 2. Successive recovery functions with rest number as the independent variable and duration of rest as the parameter.

parameter and ordinal rest number is the independent variable.

The curves of Fig. 2 converge on a common value at Rest 7, and an F less than unity at Rest 7 bears this out. After Rest 7, the recoveries of the most massed ($1/4'$) and the most spaced ($8'$) groups decline, and appear no different in magnitude. The intermediate rest group is considerably and unaccountably variable and shows no marked trend. An F of 10.29 at Rest 14 is significant beyond the .01 level with 2/189 df and is attributed to Group $1'$.

Abstract

BRAINE, M. D. S., & SHANKS, BETTY L. (Walter Reed Army Institute of Research). The development of conservation of size. *J. verb. Learn. verb. Behav.*, in press.—Using several illusions, a stimulus was first made to look bigger and then to look smaller than another stimulus, on successive trials. 4 experiments required correct differential responses to the questions “Which looks bigger?” and “Which is really bigger?”, and one experiment required that the question “Which is bigger?” be interpreted as a question about real size.

Discussion

Except for the failure of the recovery scores of Group $1'$ to turn downward after several rest periods, the magnitudes and trends in Fig. 2 are in substantial agreement with the four phenomena previously obtained on the crank or work task. The interpretation of the work phenomena borrowed freely from I_R theory developed in learning contexts (Bilodeau, 1954). Hence, it should suffice for the present learning task as well.

Ten years have passed since the writer first noted that the classical form of recovery with rest duration was limited to Rest 1 and ought not to be generalized to Rest n . The present findings of decreasing and equal recovery between Groups $1/4'$ and $8'$ again pose serious problems for students of Rest 1. Research on distribution of practice in this decade has grown out of popularity; hopefully, not because it is thought to be well understood. The terminal results of Groups $1/4'$ and $8'$ suggest the future usefulness of such ideas as little I_R (or fatigue) and possible equality or near equality in I_R .

References

- BILODEAU, E. A. Rate recovery in a repetitive motor task as a function of successive rest periods. *J. exp. Psychol.*, 1954, 48, 197-203.
- BILODEAU, E. A., & BILODEAU, I. McD. Motor-skills learning. *Annu. Rev. Psychol.*, 1961, 12, 243-280.
- KIMBLE, G. A. & HORENSTEIN, B. R. Reminiscence as a function of the length of interpolated rest. *J. exp. Psychol.*, 1948, 38, 239-244.
- MELTON, A. W. (Ed) Apparatus tests. AAF Aviation Psychology Program Research Report No. 4, 1947.

By age 5 yr. a majority of children are capable of a quite general distinction between real and phenomenal size. The earliest stages of size conservation are not amenable to study with traditional Piagetian procedures because children under 7 do not interpret “bigger” as “really bigger,” unless feedback information forces this interpretation. It is proposed that what has to be explained in accounting for the conservations is the emergence of a general distinction between real and phenomenal properties of objects. (Pre-publication copies available from the authors.)