Results

The method and apparatus employed resulted in rapid acquisition of the bar press response. Shaping of the response occurred within the first few minutes of the initial test session. A single failure in obtaining operant conditioning was with a turtle, not included in the present experiment, whose size hindered feeding in the limited depth of water and the restricted area of the test chamber.

The relationship between operant rate and water temperature is shown in Fig. 1. Each point represents a range of 4°F. The number of daily observations occurring for each temperature range is indicated by the numbers appearing in parentheses. The heavier line in the figure represents the combined performance of the four Ss. The operant rates for the individual Ss are not identical but they are relatively uniform and consistent, and the individual curves tend to parallel each other. Although each of the lines continues to rise at the higher temperatures, with the exception of the final point for S No. 3, it is probably better to assume that there would be observed an eventual decline in rate with even higher temperatures, since the 85°F water temperature corresponds to an air temperature of 95°F under the conditions tested. Higher water and air temperatures would eventually approach lethal limits.

Abstract

ERIKSEN, C. W., & STEFFY, R. A. (U. Illinois & Veterans Administration Hospital, Danville, Illinois). Short term memory and retroactive interference in visual perception. *J. exp. Psychol.*, in press.—3 experiments were performed on the relation of backward masking effects in visual perception and short term perceptual memory. In all experiments the 1st stimulus was a brief presentation of a display consisting of a random pattern of Xs and Os arranged in a circular pattern. At predetermined intervals after termination of the display (10–700 ms) an arrow appeared indicating a position on the preceding display and S reported whether the position had contained an X or O. Experiment 1 in addition to an age variable, studied the effect of an illuminated indicator field. Experiment 2 was similar to Experiment 1 except that possible brightness summation between display and indicator stimuli was eliminated.

The body temperature of turtles, being exothermal, tends to conform to that of the environment and is related to metabolic rate. This relationship is commonly expressed as a factor by which metabolic rate increases with a temperature rise of 10°C and is referred to as the Q_{10} approximation. According to Prosser & Brown (1961), this factor lies between 2.0 and 2.5. Van Sommers (1963) reported an increase in response rate close to a factor of 2 for the temperature range of 22–32°C (70.6–89.6°F) employed in his study. In the present study, the factor was approximately 1.6 (1.42–1.79) for a similar temperature range of 66–85°F. The fact that essentially linear rates were produced with a temperature range about one-third larger than that employed by Van Sommers indicates that an operant response for food reinforcement is highly reliable method for the behavioral study of the turtle.

References

- PROSSER, C. L., & BROWN, F. H. Comparative animal physiology. (2nd ed.) Philadelphia: Saunders, 1961.
VAN SOMMERS, P. Air-motivated behavior in the turtle. *J. comp. physiol. Psychol.*, 1963, 56, 590–596.

Notes

1. Supported in part by a grant-in-aid from the Florida State University Research Council.
2. Now at North Carolina State College.

Experiment 3 was essentially a replication of Experiment 2 using 4 practiced Ss on whom sufficient observations were obtained to analyze the Ss' functions individually.

In none of the experiments was any evidence found of a brief perceptual memory. Experiment 1 showed impaired performance when indicator followed display within 100 ms. These results were consistent with an interpretation in terms of brightness summation between display and indicator fields with resulting loss of figure-ground contrast for display symbols. Experiment 2 confirmed this interpretation since elimination of possibility of brightness summation resulted in no significant effect attributable to delay of the indicator. Experiment 3 indicated marked individual differences in retroactive interference functions occasioned by the delay of the indicator. (Pre-publication copies available from the authors.)