References

- HYMAN, R. Stimulus information as a determinant of reaction time. *J. exp. Psychol.*, 1953, 45, 188-196.
- MOWBRAY, G. H. Choice reaction times for skilled responses. *Quart. J. exp. Psychol.*, 1960, 12, 193-202.

Note

1. The research reported here was supported by Grant MH 06856-01 from NIH.

Comment

The report of Pick and Hay (*Psychon. Sci.*, 1964, 1, 199-200), on the prolonged wearing of 20-diopter prisms, suggests that the varying rates and amounts of adaptation to the different dimensions of distortion are "best understood in terms of the changes in stimulus dependencies involved." On the surface this is true, but the most important variable is the amount of behavioral adaptation to the distorted retinal input. In my own experiment (J. G. Taylor, *The Behavioral Basis of Perception*. Yale University Press, 1962) there was virtually complete adaptation to the lateral displacement of objects on my work table, but much less to the displacement of points on the ground on which I stood or walked. The essential difference between these situations is that in the former, movements designed to make contact with objects had to be accurately directed, but not in the latter, since one part of the floor will support the feet as well as another.

Pick and Hay report that the smallest amount of adaptation was found with respect to the instability of perceived shapes under rotation of the head about its vertical axis. In my experiment the work table and the objects on it remained perceptually stable under rotation of the head within the range that was customary while I was at work, but there was no reduction in the instability of objects beyond my reach, at the end of the experiment as compared with the beginning. Evidently such results cannot easily be related to "stimulus dependencies."

If the stimulus dependency hypothesis were valid there could be no objection to the tests used by Pick and

Hay to measure adaptation, but if we adopt a behavior dependency hypothesis their validity is questionable. For example, adaptation to the convergence of horizontal lines above and below eye level was measured by having Ss arrange four dots to make a square, presumably with its center at eye level. Now we are constantly taking action relative to horizontal surfaces below eye level, such as tables and floors; and if these surfaces, seen through prisms, appear to be oblique, the resulting action is erroneous and has to be corrected. But we do not live on our ceilings, and any behavioral errors, relative to them, that the prisms might induce remain uncorrected. Surfaces such as high shelves require occasional action, but not nearly as much as tables and floors. It is evident then that the "square" test measures two dimensions of adaptation and not one. Complete adaptation to obliquity below eye level together with zero adaptation to obliquity above eye level would yield a measure of 50% in the test. The average obtained by Pick and Hay was 51.5%.

The behavioral theory of perception leads to the conclusion that tests applied in laboratory conditions remote from those in which behavioral errors are corrected are likely to yield erroneous measures of perceptual adaptation. My own test depends on the fact that if the spectacles are worn for only part of each day the customary after-effects of continuous wearing are eliminated. This makes possible a direct comparison between (a) present perception through the prisms, (b) normal perception without prisms, and (c) perception through the prisms as it was at the beginning of the experiment. To obtain (c) one has only to turn the spectacles so that the bases are on the other side. The resulting distortion is the same as at the beginning but in the opposite direction. This test is applicable in all situations and therefore yields more valid results than any test applied in an artificial laboratory situation.

James G. Taylor
Defence Research Medical Laboratories

(See page 414 for a reply by H. L. Pick, Jr. and J. C. Hay)