Fig. 1.

metric concepts of the coulomb type. The B dimension provides a demonstration of the results of varying frequency; a curvilinear relationship seems to exist with rate of responding. It will be noted that the range in which high rates of responding occur, is quite small. The main reason for varying frequency in this experiment, was not however to examine a functional relationship, but to seek an interaction. Clearly, if there is nothing varying in A there can be no interaction

Comment

Surridge & Amsel (1965), consistent with Amsel, MacKinnon, Rashotte & Surridge (1964), conclude that massed and spaced trial patterning is contaminated or regulated by visual, olfactory, or auditory cues. Perhaps so. But available evidence fails to support the contention and much opposes it. Fairness dictates that the erroneous impression fostered by Amsel and his associates be corrected.

If small or vision regulated patterning it would be expected that running would be slower on nonrewarded (N) than on rewarded (R) trials under any reward schedule. Early investigations by the writer and others routinely included a "randomly" rewarded group which, unlike single alternation (SA), showed no tendency to pattern. Figure 1 of a later study by Bloom & Capaldi (1961) clearly shows that while SA patterned, DA (double alternation) ran as rapidly on N as on R trials and as rapidly as SA on R. Later when SA was shifted to DA, as predicted, running continued to be slow following R, fast

with B. The presence of a significant A x B interaction, is additional evidence in favor of the suggestion that there is variance in A, not accounted for by changes in charge.

In summary, it is suggested that if significant differences in rate of responding may be obtained under conditions of constant frequency and charge, using a design of limited statistical power, it would seem reasonable to conclude that stimuli should be specified in terms of amplitude and duration, and that the microcoulomb is no substitute for such specification, and no accurate indicator of magnitude of reinforcement.

References

- Keesey, Richard E. The relation between pulse frequency, intensity, and the rate of responding for intracranial stimulation. *J. comp. physiol. Psychol.*, 1962, 55, 671-678.
- McIntire, R. W., & Wright, J. E. Parameters related to response-rate for septal and medial forebrain bundle stimulation. *J. comp. physiol. Psychol.*, 1965, 59, 131-134.
- Pliskoff, S. S., Wright, J. E., & Hawkins, T. D. Brain stimulation as a reinforcer: Intermittent schedules. *J. exp. Anal. Behav.*, 1965, 8, 75-88.
- Uyeda, A. H., & Gegerelli, J. A. Influences of rectangular pulses and sine waves of varying frequencies on brain self-stimulation in the laboratory rat. *Psychol. Rep.*, 1959, 5, 641-647.
- Ward, Henry P. Stimulus factors in septal self-stimulation. *Amer. J. Physiol.*, 1959, 196, 779-782.

Note

1. This research constitutes part of a dissertation by the senior author. The work was supported by contract No. DA-49-193-MD-2288, between the Office of the Surgeon General, U. S. Army, and the University of Maryland.

following N. Note, running was now *slow* on half the R trials and *fast* on half the N trials, rather foolish behavior if the presence or absence of food could be seen, smelled, or heard. Capaldi & Senko (1962) showed that when a R trial follows two successive N trials (N_1N_2R), as predicted, running was slow on N_1 , rapid and non-differential on N_2 and R. Presumably, N_1 and N_2 smell and look the same, and since the position of the food is changed on N_1 and R these sound the same. These data invite Amsel et al to be specific rather than general in their conclusions.

Aside from food under the goal box, Surridge & Amsel's procedure differed in many particulars from Capaldi & Spivey (1964), but unaccountably none of these differences is mentioned by them as a possible basis for the discrepant findings. Concerning one difference between investigations, published information is actually available. Burt & Wike (1963), employing three SA groups, showed that N-trial running is slower and develops sooner, the longer the confinement on N

(Continued on page 132)

performance on R and NR trials throughout the experiment. The starting and running curves are virtually inseparable, while the curves for goal time show somewhat greater variability from one day to the next but are completely interlaced. Statistical analysis of these data seems superfluous.

It seems clear that rats do not exhibit pattern running at 24-hr. ITIs after 192 trials, at least in the present experiment which used procedures designed to reduce discrimination on any basis other than that provided by the reinforcement pattern. In our experiment, the SA rats were run in squads of 11, along with two other groups of 11 Ss which were under different conditions. Within squads, the assignment of Ss to carrying cages and the order of running Ss from day to day was made unsystematic. Moreover, every attempt was made not to provide S with systematic cues for R and NR trials. For example, E "false-baited" the food cup prior to NR trials and brushed out the goal box after every trial, rewarded or not. S was unable to see into the food cup until the final photobeam had been interrupted; moreover, the possibility of olfactory discrimination was reduced by having the floor under the goal box (not visible to S) covered with food dust. The use of dry food as reward, instead of wet mash, would seem further to reduce the possibility of olfactory discrimination.

Our results suggest that the pattern running reported by Capaldi & Spivey occurred on some basis other than "memories." Olfactory discrimination seems a plausible explanation of the early discrimination of R and NR trials shown in the goal measure of their study; the smaller discrimination evident in the starting and running measures in the last 30 trials might be due to the same factor or to differential cues provided inadvertently

trials (20, 80, 120 sec.). Anticipating difficulty in obtaining patterning at 24 hr., Capaldi & Spivey employed 120-sec. N-trial confinement; Surridge & Amsel, only 30-35 sec. We are reasonably confident that had Capaldi & Spivey reduced N-confinement much below 120 sec., the small differences obtained would have been reduced and very possibly eliminated.

Burt & Wike's finding suggests that N-confinement differences probably account to some extent for differences in rate of development of patterning across investigations. Surridge & Amsel characterize such differences as "strange," which suggests either unfamiliarity with Burt & Wike or a failure to understand the result. Moreover, Surridge & Amsel augment "strangeness" by comparing goal times from one investigation (Capaldi & Spivey) with total times from others (Bloom & Capaldi; Capaldi & Stanley), ignoring the fact that patterning develops differentially in different sections of the runway. Actually, their method of comparison would produce widely divergent trial figures

(See article by Surridge and Amsel, page 131)

ly by the experimenter just before a trial, e. g., baiting vs not baiting the food cup, etc. The sharp decline in running speed evident in all three measures in the Capaldi & Spivey data was not duplicated in our experiment and remains unexplained.

The discrepancy between our results and those of Capaldi & Spivey (1964) indicates that a great deal of control must be exercised over stimuli in the experimental situation if performance on a runway trial is to be attributed to "after-effects" from a previous trial in partial reinforcement experiments. This is particularly true of experiments in which a number of trials are run each day. While it is not unreasonable to expect "after-effects" to operate when ITIs are very short, some of previous studies by Capaldi and his co-workers which have employed massed trials (e. g., Capaldi & Senko, 1962; Capaldi & Stanley, 1963; Capaldi, Turner & Wynn, 1962) have produced results which may be contaminated by the inclusion of possible bases for visual, auditory and olfactory discrimination, and olfactory discrimination.

References

- Bloom, J. M., & Capaldi, E. J. The behavior of rats in relation to complex patterns of partial reinforcement. *J. comp. physiol. Psychol.*, 1961, 54, 261-265.
- Capaldi, E. J., & Senko, M. G. Acquisition and transfer in partial reinforcement. *J. exp. Psychol.*, 1962, 63, 155-159.
- Capaldi, E. J., & Spivey, J. E. Stimulus consequences of reinforcement and nonreinforcement: Stimulus traces or memory. *Psychon. Sci.*, 1964, 1, 403-404.
- Capaldi, E. J., & Stanley, L. R. Temporal properties of reinforcement aftereffects. *J. exp. Psychol.*, 1963, 65, 169-175.
- Capaldi, E. J., Turner, L., & Wynn, W. H. Decremental and facilitative effects in the straight alley runway under partial reinforcement. *J. comp. physiol. Psychol.*, 1962, 55, 545-549.

Note

1. Supported by a grant (GB-143) from the National Science Foundation.

within the Capaldi & Spivey investigation itself (see Fig. 1)

References

- Amsel, A., MacKinnon, J. R., Rashotte, M. E., & Surridge, C. T. Partial reinforcement (acquisition) effects within subjects. *J. exp. Anal. Behav.*, 1964, 7, 135-138.
- Bloom, J. M., & Capaldi, E. J. The behavior of rats in relation to complex patterns of partial reinforcement. *J. comp. physiol. Psychol.*, 1961, 54, 261-265.
- Burt, D. H., & Wike, E. L. Effects of alternating partial reinforcement and alternating delay of reinforcement on a runway response. *Psychol. Rep.*, 1963, 13, 439-442.
- Capaldi, E. J., & Senko, M. G. Acquisition and transfer in partial reinforcement. *J. exp. Psychol.*, 1962, 63, 155-159.
- Capaldi, E. J., & Spivey, J. E. Stimulus consequences of reinforcement and nonreinforcement: Stimulus traces or memory. *Psychon. Sci.*, 1964, 1, 403-404.
- Capaldi, E. J., & Stanley, L. R. Temporal properties of reinforcement aftereffects. *J. exp. Psychol.*, 1963, 65, 169-175.
- Surridge, C. T., & Amsel, A. Performance under a single alternation schedule of reinforcement at 24-hour intertrial intervals. *Psychon. Sci.*, 1965, 2, 131-132.

E. J. Capaldi and James E. Spivey
University of Texas