discriminative stimuli produced by the S by the number of food reinforcements. A ratio of 1.00 indicates one observing response per reinforcement. All data presented represent means of the last four days of data collection under a given condition.

The functions are different for the two Ss. This is mainly because of the high observing response output for P-4B at FR20 FR50. This high output is due to the regular emission of two observing responses during FR20. The event records showed that both birds made observing responses after reinforcement, before responding on the food key again. P-4A was able to run through FR20 within 5 sec. after he had made an observing response. P-4B did not do this and interrupted the FR20 in order to make another observing response. Inspection of the event records indicated that both birds infrequently made two observing response during FR50, although FR50 could not be completed within 5 sec.

Both birds give approximately the same observing ratios at FR10 FR50 and FR30 FR50. Observing behavior is strong at FR10 FR50 and very weak at FR30 FR50.

Discussion

The original plan of the experiment had been to increase the value of the lower FR to 30 in steps of 10 and then raise it in smaller steps. It was expected that as the lower FR was increased, the observing ratio would show a systematic decrease, eventually falling off to 0. The original plan of the experiment was impossible to carry out since the observing ratio fell nearly to zero when the lower FR reached FR30.

Another study of ratio discriminability (Rilling & McDiarmid, 1965) found that pigeons can easily discriminate between FR35 and FR50. In fact, fair discrimination was shown between FR45 and FR50. It is evident that in the present experiment the discriminative stimuli lost their reinforcing power before discrimination between the two schedules broke down.

The fact that pigeons are able to discriminate between two FRs does not mean that they will under all circumstances. In an observing response experiment discrimination does not increase the probability of reinforcement (Wyckoff, 1952, 1959). In retrospect, it is not surprising that the Rilling and McDiarmid experiment yields quite different results from the present experiment. In their experiment reinforcement was contingent on discriminative behavior whereas in the present experiment the observing response only provides information as to which schedule is currently present.

References

- Kelleher, R. T. Stimulus-producing responses and attention in the chimpanzee. *J. exp. Anal. Behav.*, 1958, 1, 87-103.
- Kendall, S. B. Fixed interval behavior in mixed and multiple schedules with alternating fixed interval-fixed ratio components. *Psychon. Sci.*, 1964, 1, 165-166.
- Kendall, S. B. The distribution of observing responses in a mixed FI-FR schedule. *J. exp. Anal. Behav.*, in press.
- Kendall, S. B. Competing behavior and observing responses. *Psychon. Sci.*, 1965, 3, 279-280.
- Rilling, M., & McDiarmid, C. Signal detection in fixed-ratio schedules. *Science*, 1965, 148, 526-527.
- Wyckoff, L. B., Jr. The role of observing responses in discrimination learning: Part I. *Psychol. Rev.*, 1952, 59, 431-442.
- Wyckoff, L. B., Jr. Toward a quantitative theory of secondary reinforcement. *Psychol. Rev.*, 1959, 66, 68-78.

Errata

KASPER, PHYLLIS. Disruption of position habit reversal by septal stimulation. *Psychon. Sci.*, 1965, 3, 111-112—In the 2nd paragraph under Procedure, line 6, it should read "at 20 a"; in the 1st column of page 112, line 12, "Zucker & McCleary's (1964)"; and the label on the abscissa of Fig. 2 should be "Days."

ROUTTENBERG, A. The effects of chemical stimulation in dorsal midbrain tegmentum on self-stimulation in hypothalamus and septal area. *Psychon. Sci.*, 1965, 3, 41-42—The third to the last line of the Method section should read "... study, carbachol, choline chloride, and carbachol were ..."