therefore difficult with some Ss to shape up the bar press response initially, and in a few Ss freezing on the floor came to replace bar pressing even after the latter had been established.

In addition to this effect, there appears to be an intrinsic relationship between shock intensity and gross body activity (Kimble, 1955). In the shuttle box a high activity level, whether learned or unlearned, is an advantage to S since, typically, locomotion is required both for shock termination and for avoidance. However, in the bar press situation, a high activity level is a handicap since S can perform best by remaining at or near the bar. A further difficulty arises in the bar press situation in that if non-locomotion is conditionable, S may learn to be too immobile, i.e., freeze on the bar, then S may not be able to make appropriate avoidance responses even though it is still able to make very quick and effective escape responses (Chapman & Bolles, 1964). In view of the different implications of heightened activity in the two situations it is perhaps not surprising that greater shock intensity, at least up to a point, improves the acquisition of avoidance in the shuttle box (Brush, 1957), but, as we have found, impairs acquisition in the bar press situation.

References

- BRUSH, F. R. The effects of shock intensity on the acquisition and extinction of an avoidance response in dogs. *J. comp. physiol. Psychol.*, 1957, 50, 547-552.
- CHAPMAN, J. A., & BOLLES, R. C. Effect of UCS duration on classical avoidance learning of the bar press response. *Psychol. Rep.*, 1964, 14, 559-563.
- KIMBLE, G. A. Shock intensity and avoidance learning. *J. comp. physiol. Psychol.*, 1955, 48, 281-284.
- MEYER, D. R., CHO, C., & WESEMAN, A. S. On problems of conditioning discriminated lever-press avoidance responses. *Psychol. Rev.*, 1960, 67, 224-228.

Notes

1. This study was supported by Research Grant GB-1650 from the National Science Foundation.
2. Now at McMaster University.

Comment on Kenyon & Kriekhaus by Irving Zucker

Kenyon & Kriekhaus (1965) reported that rats with septal lesions were impaired on the acquisition of a one-way Active-Avoidance Response (AAR). On the other hand, Zucker & McCleary (1964) found that cats with lesions of the septum were not impaired in the acquisition of a one-way AAR. In fact, Zucker (1965) showed that cats with lesions of the septal-limbic area that were deficient in the performance of a passive-avoidance response, acquired a one-way AAR in significantly *fewer* trials than controls.

The possibility exists that the impaired performance of rats with septal lesions on the one-way AAR is due to the fact that Ss were handled following each trial (Kenyon & Kriekhaus, 1965; Vanderwolf, 1964). The comparable experiments on the cat, in which no impairments were found, employed an avoidance apparatus which eliminated the handling of Ss between trials (Zucker, 1965). It should also be noted that in the two-way active-avoidance test, during which Ss presumably were not handled, the performance of rats with septal lesions was superior to that of controls (Kenyon & Kriekhaus, 1965).

Before assuming that septal lesions differentially affect performance on one and two-way active-avoidance responses, it would appear to be necessary to establish that avoidance learning of rats with septal lesions is not influenced by the handling of Ss during training.

References

- Kenyon, J. & Kriekhaus, E. E. Decrements in one-way avoidance learning following septal lesions in rats. *Psychon. Sci.*, 1965, 3, 113-114.
- Vanderwolf, C. H. Effect of combined medial thalamic and septal lesions on active-avoidance behavior. *J. comp. physiol. Psychol.*, 1964, 58, 31-37.
- Zucker, I. & McCleary, R. A. Perseveration in septal cats. *Psychon. Sci.*, 1964, 1, 387-388.
- Zucker, I. Effect of lesions of the septal-limbic area on the behavior of cats. *J. comp. physiol. Psychol.* 1965, in press.