was segregating. However, additional work with the RIII/ Crg1, DBA/2J and C57BL/10J strains indicates that the discontinuity of behavioral development is a general phenomenon. Figure 3 gives data for these strains over a period of 10 days. Arrows indicate the day of eye opening. Apparently the onset of visual stimulation need not correspond closely with activity depression, since eye opening is seen to precede, correspond with and follow declines in activity.

Discussion

While the effects of visual stimulation do seem to play some part in this interesting effect, other sense modalities as well may be involved, Alford & Ruben (1963) point out, for instance, that the full development of hearing in the mouse occurs around two weeks of age, corresponding well to the origin of the major effect described here. The decline in activity may be a cumulative effect of the total influx of novel stimuli as sense modalities mature.

References

- ALFORD, B. R., & RUBEN, R. J. Physiological, behavioral and anatomical correlates of the developing of hearing in the mouse. *Ann. Otol. Rhinol. Laryngol.*, 1963, 72, 1-11.
- MCCLEARN, G. E. Strain differences in activity of mice: Influences of illumination. *J. comp. physiol. Psychol.*, 1960, 58, 142-143.
- THIESSEN, D. D. Behavioral and morphological features of the wabbling-lethal mouse. *Amer. Psychologist*, 1964, 19, 503. (Abstract)

Note

1. This work was supported by Grant MH-06139 from the National Institute of Mental Health.

Comment

A recent report by Schwartzbaum & Donovanick (1965) demonstrates that cross-talk between intracranial stimulation (ICS) and peripheral shock circuits can affect behavior. They found that Ss receiving low level continuous septal stimulation were impaired on passive avoidance in a food-shock conflict situation only when cross-talk which raised the ICS intensity was permitted. Their study is in some respects similar to studies by Kasper (1963; 1964) and Pellegrino (1965). The latter authors, however, do not feel that the impairments of passive avoidance they observed were the result of electrical cross-talk. Their opinion regarding the occurrence and importance of cross-talk in their experimental situations can be summarized by the following quotation.

Brain stimulation was not discontinued during the mouth shock. However, the oscilloscope was monitored carefully and the brief artifact caused by the mouth shock did not raise the intensity of the brain stimulus. Thus the mouth shock was not associated with high, possibly reinforcing, levels of septal stimulation (Kasper, 1963, p. 9).

It seems probable that the apparent difference over the importance of cross-talk between these two sets of authors can be resolved by examining major procedural differences. Schwartzbaum and Donovanick used a peripheral shock of 0.8 ma for a duration of 0.5 sec. This

is in strong contrast to the 0.10-0.15 ma and almost instantaneous shock used by Kasper and by Pellegrino (the shock duration being set by the Ss' reflex times for pulling back from an electrified water spout). The shock situation used by Schwartzbaum and Donovanick maximized cross-talk. The situation used by Kasper and by Pellegrino minimized cross-talk by the use of non-shielded short cables, separation of shock and ICS sources and cables, and an ungrounded shock circuit; spread of current between the ICS electrode and the peripheral contact was minimized by the use of a low level of peripheral shock. The lack of an impairment on passive avoidance in Schwartzbaum and Donovanick's "no cross-talk" situation as compared to the impairment found by Kasper and by Pellegrino in their "no cross-talk" situation is probably due to the difference in the intensities of punishment.

The problem of electrical cross-talk between ICS and peripheral shock circuits is extremely complex and little understood. Perhaps the use of such noxious stimuli as loud noise or air-blasts would avoid some of the problems of interpretation posed by Schwartzbaum and Donovanick's work.

Low level continuous stimulation of the septal region produces behavioral effects in situations where no peripheral shock is involved. One example is an effect on DRL responding (Kaplan, 1965). Another example is an impairment on the reversal of a position habit (Kasper, unpublished data).

Phyllis Kasper
Louis Pellegrino
McGill University

References

- KAPLAN, J. Temporal discrimination in rats during continuous brain stimulation. *Psychon. Sci.*, 1965, 2, 255-256.
- KASPER, PHYLLIS. Attenuation of passive avoidance by continuous septal stimulation. Unpublished Master's thesis, McGill University, 1963.
- KASPER, PHYLLIS. Attenuation of passive avoidance by continuous septal stimulation. *Psychon. Sci.*, 1964, 1, 219-220.
- PELLEGRINO, L. The effects of amygdaloid stimulation on passive avoidance. *Psychon. Sci.*, 1965, 2, 189-190.
- SCHWARTZBAUM, J. S., & DONOVICK, P. J. An artifact in the use of brain stimulation with shock-motivated behavior. *Psychon. Sci.*, 1965, 2, 183-184.

Reply

The more detailed specification of test conditions employed by Kasper and Pellegrino provides useful and necessary information. Differences in shock intensity and duration, rather than the other variables described, may well have minimized their artifactual electrical cross-talk. Nevertheless, in view of the complexity of the problem and the fact that the behavioral effects of such interactions are poorly understood, it would seem desirable to employ test conditions that eliminate any possible artifact—if only to determine whether brain stimulation at the time of shock is critical for the impairment in passive avoidance.

J. S. Schwartzbaum
University of Wisconsin