Table 1.

Total number of shocks received following periods of either long (10 days) or short (5 days) approach training in a straight alley.
N = 7 in each experimental group.

	Hippocampal Lesions	Cortical Lesions	Totals
Long-Training	21	12	33
Short-Training	11	7	18
Totals	32	19	51

$F=7.42$, $df=1/24$) than the long-trained cortically lesioned rats. In contrast, the short-trained animals with hippocampal lesions did not differ significantly from either the cortically lesioned short- or long-trained groups. Finally, no significant differences were obtained between the two cortically damaged groups.

Discussion

We found that prior training does, in fact, affect the degree of the passive avoidance deficiency found in animals with bilateral hippocampal lesions. Isaacson, Olton, Bauer and Swart used latency measures as their dependent variable and reported differences between cortical and hippocampal animals regardless of the number of training trials given; although the deficit did increase with an increase in the number of trials. In the Isaacson experiment, the authors "utilized a response which would easily be performed by animals with or without training." The present experiment and the work by Kimble, Kirkby, and Stein both required a formal training period, and the response measure was *not* time-dependent. Differences in experimental design, apparatus and differences in the

nature of the dependent variable may account for the differences in results. On the basis of the present study, however, we must agree with Snyder & Isaacson (1965) that "the amount of specific training in the experimental situation should not be ignored as a distinguishing characteristic between the two passive avoidance problems." (p. 1289). We might also add that the amount of specific training should not be ignored within a particular passive avoidance problem.

References

- Isaacson, R. L., Olton, D. S., Bauer, B., & Swart, P. The effect of training trials on passive avoidance deficits in the hippocampectomized rat. *Psychon. Sci.*, 1966, 5, 419-420.
- Kimble, D. P., Kirkby, R. J., & Stein, D. G. Response perseveration interpretation of passive avoidance deficits in hippocampectomized rats. *J. comp. physiol. Psychol.*, 1966, 61, 141-143.
- Kimble, D. P. The effects of bilateral hippocampal lesions in rats. *J. comp. physiol. Psychol.*, 1963, 56, 273-283.
- Snyder, D. R., & Isaacson, R. L. Effects of large and small lesions on two types of passive avoidance responses. *Psychol. Rep.*, 1965, 16, 1277-1290.
- Stein, D. G. The effects of posttrial strychnine administration and hippocampectomy on learning and memory in the rat. Unpublished doctoral dissertation, University of Oregon, 1965.
- Stein, D. G., & Kimble, D. P. The effects of hippocampal lesions and posttrial strychnine administration on maze behavior in the rat. *J. comp. physiol. Psychol.*, 1966, in press.

Notes

1. Portions of this research were supported by a USPHS Predoctoral Research Fellowship to the first author and by USPHS Research Grant MH-08545-03 to Daniel P. Kimble.
2. Present address: Department of Psychology, Clark University.
3. A figure showing maximum and minimum extent of lesions may be obtained, upon request, from the first author.

Comment

by R. L. Isaacson

In our recent report (Isaacson, Olton, Bauer, & Swart, 1966) we did not mean to imply that training was not an important variable in determining whether or not performance deficits will be exhibited by hippocampectomized animals. In fact, we had anticipated a rather strong training effect in the experiment. However, the results of the experiment, as well as of several pilot studies, led us to conclude that specific training may not be required. We proposed that training could be one variable which enhances a "total behavioral tendency" to make the response. Other things might also contribute to this tendency. I do not believe the results of the Stein and Kirkby experiment contradict this view. An interesting extension of their design would be to run subjects with both amounts of training under enhanced deprivation conditions which

should increase the "total behavioral tendency" of the licking response. I would expect the hippocampectomized animals to make relatively more responses under these circumstances.

One point is of some theoretical concern in an experiment like that of Stein and Kirkby. Training was given in a specific runway type task, but the response measured was licking from a water bottle. Surely the licking response was highly overlearned before the experiment and it is difficult to see why a few trials more or less should have much effect upon it. From the paper it is not possible to determine whether the animals' tendencies to approach the area of the water spout was altered by the training or only the licking itself. The approach response is likely to be influenced by training. If it was not, one might consider investigating variables relating to habituation or distraction and their relation to the obtained results.