Behavior Research Methods
2010, 42 (1), 363
doi:10.3758/BRM.42.1.363

ERRATUM

Nimon, K., Lewis, M., Kane, R., & Haynes, R. M. (2008). An R package to compute commonality coefficients in the multiple regression case: An introduction to the package and a practical example. *Behavior Research Methods*, **40**, 457-466.

On page 458 of the article, the final line of Table 2 contains an error: The final term should be negative rather than positive. The corrected table is shown below.

Table 2
Unique and Commonality Formulas for Four Predictor Variables

$U_{(i)}$	=	$R^2_{y.ijkl} - R^2_{y.jkl}$
$U_{(j)}$	=	$R^2_{y.ijkl} - R^2_{y.ikl}$
$U_{(k)}$	=	$R^2_{y.ijkl} - R^2_{y.ijl}$
$U_{(l)}$	=	$R^2_{y.ijkl} - R^2_{y.ijk}$
$C_{(ij)}$	=	$-R^2_{y.kl} + R^2_{y.ik} + R^2_{y.jkl} - R^2_{y.ijkl}$
$C_{(ik)}$	=	$-R^2_{y.jl} + R^2_{y.ijl} + R^2_{y.jkl} - R^2_{y.ijkl}$
$C_{(il)}$	=	$-R^2_{y.jk} + R^2_{y.ijk} + R^2_{y.jkl} - R^2_{y.ijkl}$
$C_{(jk)}$	=	$-R^2_{y.il} + R^2_{y.ijl} + R^2_{y.ikl} - R^2_{y.ijkl}$
$C_{(jl)}$	=	$-R^2_{y.ik} + R^2_{y.ijk} + R^2_{y.ikl} - R^2_{y.ijkl}$
$C_{(kl)}$	=	$-R^2_{y.ij} + R^2_{y.ijk} + R^2_{y.ijl} - R^2_{y.ijkl}$
$C_{(ijk)}$	=	$-R^2_{y.l} + R^2_{y.il} + R^2_{y.jl} + R^2_{y.kl} - R^2_{y.ijl} - R^2_{y.ikl} - R^2_{y.jkl} + R^2_{y.ijkl}$
$C_{(ijl)}$	=	$-R^2_{y.k} + R^2_{y.ik} + R^2_{y.jk} + R^2_{y.kl} - R^2_{y.ijk} - R^2_{y.ikl} - R^2_{y.jkl} + R^2_{y.ijkl}$
$C_{(ikl)}$	=	$-R^2_{y.j} + R^2_{y.ij} + R^2_{y.jk} + R^2_{y.jl} - R^2_{y.ijk} - R^2_{y.ijl} - R^2_{y.jkl} + R^2_{y.ijkl}$
$C_{(jkl)}$	=	$-R^2_{y.i} + R^2_{y.ij} + R^2_{y.ik} + R^2_{y.il} - R^2_{y.ijk} - R^2_{y.ijl} - R^2_{y.ikl} + R^2_{y.ijkl}$
$C_{(ijkl)}$	=	$R^2_{y.i} + R^2_{y.j} + R^2_{y.k} + R^2_{y.l} - R^2_{y.ij} - R^2_{y.ik} - R^2_{y.il} - R^2_{y.jk} - R^2_{y.jl} - R^2_{y.kl} + R^2_{y.ijk} + R^2_{y.ijl} + R^2_{y.ikl} + R^2_{y.jkl} - R^2_{y.ijkl}$

© 2010 The Psychonomic Society, Inc.