

Index

A

- Acanthoparyphium*, 5, 8
- Acanthoparyphium tyosenense*, 10, 152, 153
- Acris crepitans*, 76
- Adult echinostomes
 - chemoattraction and mate finding, 99
 - growth and development, 97–98
 - mating behavior, 100–101
- Amphibians
 - echinostomes
 - emergence of, 268–269
 - hosts involved in, 254–257
 - infection pathology and patterns, 265–268
 - second intermediate hosts, 265
 - vs. *Ribeiroia*, 264
 - environmental changes, macroparasite infections
 - biodiversity loss, 272–273
 - climate change, 273
 - importance of, 275
 - land use and wetland characteristics, 270
 - nutrient pollution, 270–271
 - pesticide contamination, 271
 - predation risk, 274, 275
 - Ribeiroia*
 - emerging infection, 262–263
 - frog deformities and trematode infection, 253, 258, 259
 - hosts involved in, 254–257
 - malformation associated with, 252
 - mass malformations, 262
 - Ribeiroia*-induced pathology, 259–261
 - trematodes, importance in, 250–252
 - Anas boschas fereae*, 165
 - Anas platyrhynchos*, 66
 - Anthelmintic drugs, 298–299
 - Antibody responses
 - immune proteomic approach, 197–198

- primary echinostome infections, 196–197
- Antigenic characterization, adult echinostomes, 199–200
- Antioxidant factors, 42
- Argentophilic papillae, 11
- Artificial spring water (ASW), 119, 120
- Artyfechinostomum malayanum*, 154
- Artyfechinostomum oraoni*, 154–155
- Artyfechinostomum sufrartyfex*, 154
- Austrovenus stutchburyi*, 74

B

- Babesia*, 289
- Batrachochytrium dendrobatidis*, 251
- Bellamya aeruginosa*, 22
- Benzimidazoles, 298
- Bge cells, 40, 65
- B. glabrata*, 14, 137
- Biomphalaria glabrata*, 14, 36, 162
 - embryonic (Bge), 39
 - hemocytes, effect of echinostome excretory/secretory products on, 42
- Bivalvia, 63
- Bulinidae, 14

C

- Caballerotrema*, 5
- Ca ions, 41
- Cathaemasiidae, 7
- Cercariae
 - chaetotaxy and its patterns, 3, 15
 - Echinochasmus* spp., 9
 - mortality, 63
 - Parorchis acanthus*, 10
 - second intermediate hosts studies, 18
 - survival and effects of temperature, 16
 - in vitro encystment, 123

- Cercariae and metacercariae, of *E. caproni* and *E. trivolvis*
- effects of copper sulfate toxicity on, 77
 - metolachlor and atrazine on host–parasite system, 78
 - toxicity of antifouling biocides tributyltin (TBTO) and, 77–78
- Cercarial tail of *Echinostoma* species, 15
- Cercaria yamagutii*, 152
- Cerithidea cingulata*, 152
- Cerithidea largillierti*, 152
- Chemotherapy, 296
- Cipangopaludina chinensis malleata*, 160
- Cipangopaludina japonica*, 165
- Cipangopaludina malleata*, 165
- Cipangopaludina* sp., 160
- 43- and 45- collar-spined
- Echinoparyphium*, 5
- 37-collar-spined *Echinostoma*, 6, 63
- cercariae encyst and develop to metacercariae in vitro, 63
- 37-collar-spined echinostomes, 8
- 37-collar-spined revolutum group of
- Echinostoma*, 6
- Collar spines, importance for taxonomic study, 148
- Column liquid chromatography
- methods and applications, 243–244
 - principles, 243
- Corbicula lindoensis*, 162
- Corbicula producta*, 165
- Corbicula sucplanta*, 162
- Crassostrea virginica*, 73
- Curtuteria australis*, 11
- Cytochrome c oxidase subunit 1 (CO1), 7
- Cytokine responses, 198–199
- D**
- Digenea, 36
- Digestive gland–gonad complex (DGG)
- metallic ion study, 244–245
 - sterol content, 242
- Digoniostoma pulchella*, 154
- DNA genome studies, echinostomes
- EST sequencing, 213
 - internal transcribed spacers (ITS)
 - sequence, 210–211
 - putative gene library, 212
- Dreissena polymorpha*, 64, 75
- Drepanocephalus*, 3
- Dulcolax™, 119
- E**
- Echinochasmus*, 8, 63
- Echinochasmus beleocephalus*, 156
- Echinochasmus fujianensis*, 7, 155
- Echinochasmus japonicus*, 155–156
- Echinochasmus jiufoensis*, 156
- Echinochasmus liliputanus*, 22, 63, 133
- Echinochasmus liliputanus* cercariae, 123
- Echinochasmus perfoliatus*, 157
- Echinoparyphium*, 3, 5, 8
- Echinoparyphium aconiatum*, 65
- Echinoparyphium elegans*, 7, 63
- Echinoparyphium megacirrus*, 18
- Echinoparyphium paraulum*, 165
- Echinoparyphium recurvatum*, 10, 133, 134, 158–160
- Echinoparyphium recurvatum* metacercariae, 63
- Echinoparyphium serratum*, 92
- Echinoparyphium* sp., 10
- Echinoparyphium* spp., 253, 264
- Echinostoma*, 2, 3
- wild animals
 - Asia and Europe, 132
 - North America and Central and South America, 131
 - zoonotic infections and main intermediate hosts, 133
- Echinostoma angustitestis*, 160
- Echinostoma caproni*
- cercariae, 15
 - echinostome cysts, 65
 - echinostome infections, 37
 - experimental models, 134
 - immune evasion, 40
 - infection, metabolic profiling
 - biofluids utility, 312
 - diagnostics metabolites, 307–309
 - plasma, 310
 - stool, 311–312
 - urine, 310–311
 - metacercariae pathology, 75
 - miracidium, 11, 13
 - newly hatched miracidium, 38
 - in ovo cultivation, 120–121
 - primary infection, 96
 - species-rich genera, 6–8
 - sporocysts, 42
- Echinostoma caproni* mouse model
- integrative metabonomics
 - cross cellular transport, 317–318
 - holistic influence, gut microbiota, 318
 - metabolite flow, 319

- intestinal profile changes
 - amino acids levels, 317
 - glucose concentration, 316
 - SCFAs role, 314–315
- MAS NMR spectroscopy
 - kidney, 313
 - liver, 313–314
 - spleen, 314
- metabonomic investigation
 - biomarker identification and diagnostic potential, 306–312
 - experimental protocol, 305–306
 - metabonomics studies, 302–305
- trematocidal drug discovery and development
 - drug sensitivity assay, 300
 - scanning electron microscopic (SEM) studies, 300–301
 - spectroscopy, 302
 - in vivo* evaluation, 297–299
- E. cinetorchis*
 - cercariae, 63
 - description, 160, 161
- E. echinatum*, 160–162
- E. friedi*, 12
- E. hortense*, 162, 164, 174
- E. ilocanum*, 164–165
- E. luisreyi*, 65
- E. macrorchis*, 165
- E. paraensei*, 17, 37, 65, 134
- E. revolutum*, 133, 137, 165–166, 254–257, 264, 265, 267
- Echinostoma*, sequencing ND1 and ITS genes, 7
- Echinostomatidae
 - classification, 4–5
 - cysts, 64
 - flukes, 148
 - genera of, 2
 - major and minor species, 130–134
- E. trivolvis*, 63, 112, 254–257, 264, 267
- Echinostome cercariae
 - morphology and immunocytochemistry, 15
 - second intermediate host, 63
 - survival and infectivity characteristics, 16
- Echinostome coinfection
 - helminths
 - nematodes, 287
 - tapeworms, 286–287
 - trematodes, 284–286
 - heterologous echinostomes
 - concurrent infections, 282–283
 - primary infections, 283–284
 - protozoans
 - Babesia*, 289
 - plasmodium*, 287–289
 - Trypanosomes*, 290
- Echinostome infection
 - emerging disease in green frogs, 76
 - host resistance, 172
 - metacercariae excystment, 91–93
- Echinostome life cycle maintenance
 - Dr. Bernard Fried's laboratory studies, 114–115
 - laboratories, 113
 - mammals in the wild, life cycle initiation source, 119
 - obtaining material from, 116
 - relationships between echinostomes, 117–118
- Echinostome-like digeneans, 2, 5, 8
- Echinostome-like metacercariae, *Ribeiroia*, 21–22
- Echinostome-like *Ribeiroia*, rhythmic emergence, 17
- Echinostome-like trematode group, 5
- Echinostome metacercariae
 - cysts storage, 23
 - encystment, 62–63
 - invertebrate and vertebrate hosts
 - behavioral and morphological changes, 73–74
 - marinades effects, 78–79
 - pathology of, 75–77
 - pesticides and herbicides effects, 77–78
 - second intermediate hosts, 67–72
 - second intermediate hosts
 - development stages, 65–66
 - excystation, 67
 - site selection in, 63–64
- Echinostome miracidia, 13
- Echinostome-related genera, 8
- Echinostomes
 - adult echinostomes
 - growth and development of, 97–98
 - mating behavior of, 98–99
 - column liquid chromatography
 - methods and applications, 243–244
 - principles, 243
 - definitive host establishment, 93–96
 - avian hosts, 94
 - concurrent infections with other helminths, 95–96
 - E. trivolvis*, compatibility with hamsters, 95
 - increased dosage levels' influence, 95

- Echinostomes (*cont.*)
- rats as hosts, low compatibility with
 - E. caproni*, 94
 - recovery rates, *E. paraensei*, 95
 - use of rodents as experimental hosts, 94
 - description, 186
 - emergence of, 268–269
 - excystation, 93, 121–122
 - experimental models
 - life cycle of *Echinostoma* spp. in wild, 135
 - major echinostome species, 134–137
 - minor echinostome species, 137
 - gas chromatography
 - methods and applications, 241–243
 - principles, 241
 - genomics
 - DNA and RNA studies, 210–213
 - prospects and limitations, 213–214
 - taxonomic and phylogenetic studies, 209–210
 - genomics and proteomics, 208–209
 - host specificity and infectivity, 90–91
 - human infection
 - E. hortense* infection endoscopic findings, 171
 - pathology and symptomatology, 167–171
 - immune mechanisms
 - antibody responses, 196–198
 - cytokine responses, 198–199
 - eosinophils, 193–194
 - goblet cells, 194–196
 - mast cells, 192–193
 - immunobiology
 - different strains and host species, 189–190
 - immunosuppressed mice, 189
 - rodent models, 188–189
 - immunodiagnosis
 - antibody detection, 200
 - seroantigen and coproantigen detection, 201
 - inductively coupled plasma-optical emission spectrometry
 - methods and applications, 245
 - principles, 244
 - infection pathology and patterns
 - in amphibian kidney, 263, 267
 - geographic and frequency distribution, 266
 - population level impacts, 267, 268
 - innate immunity, 44
 - Biomphalaria Glabrata*
 - antidigenean responses in, 47–49
 - immune system, 44–45
 - Echinostoma* vs. *Schistosoma*, tools for functional validation, 51
 - immune-relevant genes, predicted identity of, 50
 - invertebrate immunity, perceived simplicity of, 45–46
 - invertebrates, immunological memory in, 46–47
 - transcriptomic and proteomic studies, 49, 51
 - metacercarial excystment, 91–93
 - molecular-based studies, 6
 - natural and experimental infections, 138–139
 - pathology, 190–192
 - proteomics
 - analytical procedures, 217–220
 - immunoproteomics, 220–222
 - prospects and limitations, 222–223
 - protein identification, 214–217
 - resistance manifestations to infections
 - fecundity reduction, 188
 - host immunosuppression, 186
 - primary infections, 187
 - secondary infections, 187–188
 - Ribeiroia ondatrae* life cycles, 254–257
 - second intermediate hosts, 265
 - and snails, exploring complex interactions, 35–37
 - systematics, 2
 - thin-layer chromatography
 - amino acid analysis, 239–240
 - carbohydrates detection, 240
 - chromatogram quantification, 234–235
 - layers and mobile phases, 232–233
 - lipid studies, 235–239
 - lipophilic pigments identification, 240–241
 - principles, 230–231
 - sample application and plate development, 233
 - sample preparation, 231–232
 - statistical data analysis, 235
 - zone detection and identification, 234
 - vertebrate host fecundity, 101–104
 - vs. *Ribeiroia*, 264
 - in the wild
 - helminths co-infection, 138
 - major and minor species, 130–134
 - worm distribution and habitat location, 96–97

- Echinostomes and related genera
 description of life cycles, 10–11
 free-living stages
 cercaria, 15–19
 miracidium, 11–14
 generalized life cycle of
 echinostome, 10
 life cycle pattern, 9–11
 parasitic stages
 invertebrate hosts, 19–24
 vertebrate hosts, 24–25
- Echinostome trematodes, 62
- E. cinetorchis*, 11
- Ectosymbiotic flatworm, 18
- E. deserticum*, 3
- E. echinatum*, 6, 8, 9, 65
- E. friedi*, 2, 6, 8, 102, 103, 134
- E. hortense*, 91
- E. ilocanum*, 78
- E. japonicus*, 7
- E. jurini*, 6, 11
- E. liliputanus* cercariae, 65
- E. luisreyi*, 3
- E. malayanum*, 78, 119, 154
- Emden-Meyerhof-Parnas pathway, 42
- E. megacirrus*, 10
- Emergence pattern analysis, 17
- E. miyagawai*, 6
- Encysted metacercaria, 75–76
- Eosinophils, 193–194
- E. paraensei*, 6, 7, 8, 9, 11, 13, 14, 138
- E. paraensei* and *E. caproni* miracidia hatch,
 12, 13
- E. parvocirrus*, 6
- Episthmium caninum*, 166
- E. pseudorecurvatum*, 7
- E. recurvatum*, 7
- E. recurvatum* metacercarial cysts, 66
- E. revolutum*, 6, 7, 8, 9, 11, 13, 65
- ES proteomes, 42
- E. trivolvis*, 6, 7, 8, 9, 11, 13, 65, 91, 93, 96,
 102, 103, 134
- E. trivolvis* excystation, 92
- Euparyphium*, 3, 8
- Euparyphium albuferensis*, 3, 96
- Excretory/secretory products (ESP)
 description, 208–209
 ex vivo studies, 41
 protein characterization, 216–217
 proteomic studies
 echinostome sporocysts, 43–44
 glycolytic enzymes, 42–43
 molecular processes, 41–42
- Expressed sequence tags (EST), 208, 223
- F**
- Fasciola revoluta*, 165
- Fascioloides magna*, 40
- Flukes, 148, 152, 166
- Food-borne trematode infections, 140
- G**
- Gas chromatography
 methods and applications
 fatty acid determination, 242–243
 sterols identification, 241–242
 principles, 241
- Gastropoda, 40
- GenBank, 7, 43
- Genomics
 definition, 208
 echinostomes
 DNA and RNA studies, 210–213
 prospects and limitations,
 213–214
 ribosomal and mitochondrial markers,
 209–210
- Glossaulax didyma*, 152
- Global biodiversity, of animals, 130
- Glycerophosphocholine (GPC) activity
 cross cellular transport, 317–318
 kidney and liver, 313–315
- Glycolytic enzymes, 42
- Glycomics, 222
- Gnathopogon strigatus*, 155
- Goblet cells
 echinostome expulsion, 195–196
 in human echinostomes, 172–173
 hyperplasia, 194–195
- Gyraulus convexiusculus*, 154
- H**
- Helisoma*, 14
- Helisoma trivolvis*, 112
- Helminths, echinostome coinfection
 nematodes, 287
 tapeworms, 286–287
 trematodes
 E. caproni infections, 285–286
 Schistosoma spp, 284–285
- H. elongata* larvae, 63
- Hemocyte-mediated cytotoxicity, 42
- Heterologous echinostomes, 282–284
- Himasthla*, 5, 8
- Himasthla elongata*, 63, 121
- Himasthla escamosa*, 10
- Himasthla leptosoma*, 92

- Himasthla muehlensii*, 166
Himasthla quissetensis, 9, 133
Himasthla rhigedana, 63, 133
 Human echinostomiasis
 diagnosis, treatment, and control, 174–175
 immunology of, 172
 cytokines and Th1/Th2 immune responses, 173–174
 effector cells and antibodies, role of, 173
 goblet cells and mast cells role, 172–173
 host resistance to, 172
 symptoms and diagnosis, 140
 Human-infecting echinostomes
 egg size, 175
 geographical distribution of, 152
 human/animal infections' sources, 151
 morphological characters and life cycle, 149–150
 taxonomy, biology and epidemiology of, 150
Hypoderaeum, 3, 8
Hypoderaeum conoideum, 7, 10, 134, 166–167
Hypomesus olidus, 155
 Hypoxic environment, 39
- I**
Idiopoma javanica, 162
 Immune mechanisms, echinostome infections
 antibody responses, 196–198
 cytokine responses, 198–199
 eosinophils, 193–194
 goblet cells, 194–196
 mast cells, 192–193
 Immunobiology, echinostomes
 different strains and host species, 189–190
 immunosuppressed mice, 189
 rodent models, 188–189
 Immunodiagnosis
 antibody detection, 200
 seroantigen and coproantigen detection, 201
 Immunome, 220, 222
Indoplanorbis exustus, 154, 166
 Inductively coupled plasma-optical emission spectrometry
 methods and applications, 245
 principles, 244
 Integrative metabonomics
 cross cellular transport, 317–318
 holistic influence, gut microbiota, 318
 metabolite flow, 319
- Internal transcribed spacers 1 and 2 (ITS1 and ITS2), 7
 Intestinal profile changes, *E. caproni* infection
 amino acids levels, 317
 glucose concentration, 316
 SCFAs role, 314–315
 Intramolluskan echinostome stages, 12
 Intramolluskan parasitic stages, 21
Isthmiophora, 3, 8
Isthmiophora, chemo-orientation in, 16
Isthmiophora melis, 167
 ITS gene, 7
- K**
 Karyotyping, 6
- L**
Larus crassirostris, 153
 Larval and adult echinostomes
 maintenance of, 119
 transplantation to new sites, 121
 Larval echinostomes
 immune evasion strategies
 ex vivo studies, 41
 proteomic studies, 41–44
 in vivo studies, 40
Littorina littorea, 63, 121, 166
 Locke's ASW (1:1) medium, 123
Lunatia fortunei, 152
Lymnaea, 40
Lymnaea columella, 14, 137
Lymnaea corvus, 167
Lymnaea (Bullastra) cumingiana, 78, 154
Lymnaea peregra, 14, 63, 160, 167
Lymnaea pervia, 160
Lymnaea tumida, 166
 Lymnaeidae, 14
 Lymnaeid snails, for *H. conoideum* miracidia, 13
- M**
Mactra veneriformis, 152
 MAPK-associated signaling pathways, 51
 MAS NMR spectroscopy
 kidney, 313
 liver, 313–314
 spleen, 314
 Mass spectrometry (MS), 303–304
 Mast cells
 echinostome infections resistance, 192–193
 human echinostomes, 172–173

- Melanitta fusca stejnegeri*, 152
Melanitta nigra americana, 152
- Metabolic profiling, 297
 applications, 302–303
 chemometric tools, 304–305
E. caproni infection
 biofluids utility, 312
 diagnostics metabolites, 307–309
 plasma, 310
 stool, 311–312
 urine, 310–311
 experimental protocol, 305–306
 integrative metabolomics
 cross cellular transport, 317–318
 holistic influence, gut microbiota, 318
 metabolite flow, 319
- Metabolomics. *See* Metabolic profiling
- Metacercariae, echinostomes
 cysts viability and copper, 23–24
 excystment, 91–93
 site selection, second intermediate hosts,
 63–64
 in situ evaluation, 79
- Mical-like protein, 42–43
- Miracidia
E. caproni, 13, 39
E. friedi, 13–14
E. trivolvis, 13
Euparyphium albufereensis, 13
- Misgurnus anguillicaudatus*, 160
- Mitochondrial DNA (mtDNA), 7
- Movement patterns, of cercariae, 15
- Myofibril F-actin staining, 15
- Mytilus edulis*, 63
- N**
- N. braziliensis*, 96
Nectomys squamipes, 137, 140
- Nematodes, echinostome coinfection, 287
- Neocanthoparyphium echinatoides*, 7
- Nicotinamide adenine dinucleotide
 dehydrogenase subunit 1 (ND1), 7
- Nippostrongylus braziliensis*, 96
- Nonhost snails, 13
- N. squamipes*, 138, 139
- Nucella lapillus*, 10
- Nuclear magnetic resonance (NMR)
 spectroscopy. *See also* MAS NMR
 spectroscopy
 biofluids analysis, 303
 metabolic profiling, protocol, 305–306
Trypanosoma brucei, 302
- Nuclear ribosomal DNA (rDNA), 7
- O**
- Odontobutis obscura interrupta*, 163
Oncomelania, 40
Ondatra zibethica, 25
 Oxidative damage, 42
 Oysters, 73
- P**
- 19 papilla-like structures, 11
Parafossarulus manchouricus, 155
Parafossarulus striatulus, 157
- Parasitic stages, echinostomes
 invertebrate host, 19–24
 metacercariae, 21–24
 sporocysts and rediae, 19–21
- Parasitism, 140
- Parorchis*, 5, 8
Parorchis acanthus, 92
Paryphostomum, 3
P. combesi, 10
P. distomatosa, 10
Petasiger, 8
Petasiger encyst, 63
Petasiger variospinosus, 10
 Philophthalmidae, 5
Philophthalmus lucipetus, 10
Physa acuta, 160
Physa alexandrina, 160
Physa fontinalis, 160
Physa marmorata, 14, 137
Physastra hungerfordiana, 78
- Physidae, 14
Pila conica, 165
Pila luzonica, 78
Pila scutata, 154
Pisidium coreanum, 73
P. islandicus, 10
Planorbella, 14
Planorbella trivolvis, 264
 Planorbidae, 14
Planorbis compressus japonicus, 165
Planorbis corneus, 166
Planorbis planorbis, 160
Plasmodium, 287–289
Plasmodium falciparum, 112
- Polymerase chain reactions (PCR), 6
 Praziquantel, 119
- Principal component analysis
 (PCA), 304
- Prosobranchia, 40, 63
- Proteomics
 analytical procedures
 imaging and comparison, 219

- Proteomics (*cont.*)
- protein identification, mass spectroscopy, 219–220
 - protein separation and staining, 218–219
 - sample preparation, 217–218
- definition, 208–209
- echinostomes
- antigens identification, 220–222
 - prospects and limitations, 222–223
 - protein characterization, 214
 - protein identification
 - enzymes and ligand-binding molecules, 214–215
 - ESP characterization, 216–217
- Protozoans, echinostome coinfection, 287–290
- Babesia*, 289
 - plasmodium*, 287–289
 - Trypanosomes*, 290
- Pseudechinoparyphium echinatum*, 7
- Pseudomyicola spinosus*, 74
- Psilostomidae, 7
- R**
- Radix auricularia coreana*, 160
- Radix peregra*, 14
- Radix quadrasi*, 78
- Rana pipiens*, 64
- Rana temporaria*, 160
- Random amplification of polymorphic DNA (RAPD), 6
- Rattus rattus*, 25
- R. clamitans*, 64, 76
- R. congolensis*, 7
- Reactive oxygen species (ROS), 43
- Retractile apical papillae, 11
- Revolutum* group, 5
- Ribeiroia*
- cercariae, 258, 267, 271
 - emerging infection, 262–263
 - first intermediate hosts, 5, 9, 14
 - frog deformities and trematode infection
 - frequency distribution, 261
 - geographic distribution, 259, 260
 - limb malformations, 258
 - hosts involved in, 254–257
 - malformation associated with emerging infection, 262
 - representative malformation, 252
 - Ribeiroia*-induced pathology, 259–261
- Ribeiroia*, ITS2 sequences, 7
- Ribeiroia ondatrae*, 251, 253–257, 264
- R. marini*, 7
- RNA genome studies. *See* DNA genome studies, echinostomes
- R. ondatrae*, 7
- ROS-scavenging molecules, 43
- R. peregra*, 14
- Ruditapes philippinarum*, 152
- S**
- Scanning electron microscopic (SEM) studies, 300–301
- Schistosoma haematobium*, 37
- Schistosoma intercalatum*, 40
- Schistosoma japonicum*, 40
- Schistosoma mansoni*, 36, 37, 40, 138
- Schistosoma mansoni* sporocysts, 42
- Schistosoma mattheei*, 40
- Second intermediate hosts, echinostome metacercariae
- developmental stages
 - cysts wall and shape, 65
 - surface topography, echinostome digenean, 66
 - excystation, 67
 - invertebrate and vertebrate hosts, 67–73
 - metacercarial site selection in, 63–64
- Segmentina hemisphaerula*, 73, 160
- Segmentina nitiella*, 165
- Sensilla in cercariae, 6
- Singhia*, 2, 5
- Snail intermediate hosts, echinostomes
- host exploitation vs. host immune response, 39–40
 - miracidia host-finding behavior, 37–39
- S. nitiella*, 165
- Solen grandis*, 152
- Solen strictus*, 152
- Spectroscopy, 297, 302
- Sporocysts, 40
- Stephanoprora*, 3, 8
- Stephanoprora aylacostoma*, 10
- Stephanoprora uruguayense*, 10
- Strigeidida*, 36
- T**
- Tapeworms, 286–287
- Taxidea taxus*, 25
- Tegument, 11
- Temnocephala chilensis*, 18
- Thin-layer chromatography
- applications in echinostomes
 - amino acids, 239–240

- carbohydrates, 240
 - lipids, 235–239
 - lipophilic pigments, 240–241
 - chromatogram quantification, 234–235
 - data analysis, 235
 - layers and mobile phases, 232–233
 - principles, 230–231
 - sample application and plate development, 233
 - sample preparation, 231–232
 - zone detection and identification, 234
 - Toxic pollutants, 17
 - Trematocidal drug discovery and development
 - drug sensitivity assay, 300
 - echinostomicidal activity evaluation
 - anthelmintic studies, 297–298
 - drug screening studies, 298–299
 - scanning electron microscopic (SEM) studies, 300–301
 - spectroscopy, 302
 - Trematode infections, 296
 - Trematodes, 250, 251, 284–286. *See also* *Echinostoma*; *Ribeiroia*
 - Trematodes, exposed to ROS, 43
 - Trypanosomes*, 290
 - Trypsin-bile salts (TB) medium, 79
 - Tympanotonus microptera*, 152
-
- U**
 - Uroproctepisthmium*, 3
-
- V**
 - Valvata piscinalis*, 160
 - Venus mercenaria*, 166
 - Vertebrate host, echinostomes fecundity
 - factors affecting fecundity, 101–103
 - methods for measuring fecundity, 103–104
 - Viviparus javanicus*, 165
 - Viviparus malleatus*, 165
-
- W**
 - Wild echinostome zoonoses, potential of, 139–140
 - Wildlife conservation, impact of parasitism, 140–141
-
- Z**
 - Zebra mussels, 64, 75
 - Zoonotic disease, 134