

# Index

- A**
- Abortion, spontaneous, in Turner syndrome, 45
  - Actin, immunoreactivity of
    - with myofibromatosis cells, 60
    - with polyphenotypic small-cell tumors, 103
    - with rhabdomyoblasts of ectomesenchymomas, 103
    - with spindle cells of tufted angioma, 44, 116
    - with xanthogranuloma cells, 83
  - Acute lymphoblastic leukemia (ALL)
    - association with Down syndrome, 168
    - biopsy in, 175
    - relative incidence of, 161
  - Acute megakaryoblastic leukemia (AMKL)
    - association with Down syndrome, 164, 168
    - diagnosis of, 178
  - Acute monocytic leukemia, 194
  - Acute myeloid leukemia, 172
  - Acute myelomonocytic leukemia, diagnosis of, 176–177
  - Acute nonlymphoblastic leukemia (ANLL), association with Down syndrome, 168
  - Acute nonlymphoblastic leukemia (ANLL), incidence of, 161
  - Adenoma
    - adrenocortical, 341–346
    - hepatic, 329
    - pancreatic, 360
  - Adenomatous hyperplasia, 357–358
  - Adipose tissue tumors, 93–96
  - Adrenal cortical hyperplasia, congenital, 337–341
  - Adrenal cysts, 351
  - Adrenal neuroblastoma, 142
    - blue-cell tumors, 155–156
    - incidence of, 150
  - Adrenocortical steroids, biosynthetic pathways involving, 339
  - Adrenocortical tumors, association with adrenocortical hyperplasia, 337–352
  - Adrenogenital syndrome, 337–341
  - Airway obstruction, in Langerhans cell histiocytosis, 188
  - Aldosterone synthesis, impairment of, in 21-hydroxylase defect, 338
  - Alkaline phosphatase, placenta-like, immunoreactivity of, with embryonal carcinoma cells, 31
  - Alveolar macrophages, differentiation of histiocytes to, 181
  - Alveolar rhabdomyosarcoma, 89–92
  - Anaplasia, and prognosis, in Wilms tumor, 273, 277
  - Anaplastic astrocytoma, 204
    - characteristics of, 207
  - Anemia, in disseminated Langerhans histiocytosis, 184
  - Aneurysmal bone cyst, 415
  - Angioma, tufted, 44, 116–117
  - Angiomatoid fibrous histiocytoma, 78–81
  - Antigen, processing by macrophages, 183
  - $\alpha_1$ -antitrypsin, immunoreactivity of
    - embryonal and fetal epithelial hepatoblastoma cells, 310
    - with embryonal sarcoma of the liver, 325
    - with pancreatoblastoma cells, 360
    - with yolk sac tumors, 25
  - Apoptosis, in hemangioma involution, 42
  - Askin tumor, tumors related to
    - histogenetically, 99
  - Astrocytic tumors, of the retina, 252–255
  - Astrocytoma, 201–207
  - Atypical teratoid/rhabdoid tumor, 233–234
  - Autosomal dominant conditions
    - blue rubber bleb nevus syndrome, 117
    - sarcoma, breast and brain tumor, leukemia, laryngeal, and lung cancer, and adrenal cortical carcinoma (SBLA) syndrome, 341
    - tuberous sclerosis and neurofibromatosis, 1
  - See also* Autosomal recessive conditions; Congenital conditions; Familial disorders
  - Autosomal recessive conditions
    - Chediak-Higaski syndrome, 195
    - congenital adrenal hyperplasia, 337–341
    - infantile polycystic disease of kidney and liver, 265–266, 333
    - Omenn disease, 194
  - See also* Autosomal dominant conditions; Congenital conditions; Familial disorders
- B**
- Bean syndrome, 117–118
  - Becker nevus, 119
  - Beckwith-Wiedemann syndrome
    - adenomatous hyperplasia of islet cells associated with, 356–359
    - adrenocortical cytomegaly associated with, 350–351
    - adrenocortical neoplasms associated with, 341
    - adrenocortical tumors associated with, 337
    - hepatoblastoma associated with, 303
    - pancreatoblastoma associated with, 360
  - Birbeck granules, characteristic of Langerhans histiocytosis, 188
  - Blueberry muffin baby, 117–118
    - skin tumors in, 128, 174
    - with stage IV-S neuroblastoma metastatic to the skin, 150
  - Blue-cell tumors
    - differential diagnosis of, 89
    - in Ewing sarcoma, 420–423
    - retinoblastoma as, 248

- Blue nevus, in a congenital melanocytic nevus, 124
- Blue rubber bleb nevus syndrome, 117–118
- Bone
- involvement in disseminated Langerhans histiocytosis, 184
  - in teratomas, 18
  - tumors and tumorlike conditions of, 407–423
- Bone age, in adrenocortical adenoma, 346
- Bone marrow
- involvement in Langerhans cell histiocytosis, and prognosis, 189
  - involvement in medulloblastoma, 210
  - monocytes of, 181
- Bone marrow smear
- showing histiocytic hemophagocytosis, 192–194
  - showing neuroblastoma metastases, 150
- Botryoid form, of embryonal rhabdomyosarcoma, 83–85
- Brain tumors, 199–240
- incidence of, in newborns in the United Kingdom, 1–2
  - involvement in hemophagocytic lymphohistiocytosis, 190
  - metastatic, choriocarcinoma, 32
- Brown fat, lipoma, 94
- C**
- Café-au-lait spots, association with neurofibromas, 97, 125–126
- Calcification, and prognosis, in neuroblastoma in infants, 156
- Calcifying aponeurotic fibroma, 70
- Carcinoembryonic antigen, immunoreaction of pancreatoblastoma cells with, 360
- Carcinoma, adrenocortical, 347–350
- Cardiac tumors, 377–385
- Cardiomegaly, association with cardiac tumors, 377
- CD1a antigen, immunoreactivity to, of juvenile xanthogranulomas, 132
- CD1 antigen (T8 thymocyte antigen), immunoreactivity to, in Langerhans cell histiocytosis, 130
- Cellular congenital mesoblastic nephroma, 280, 284–287
- Cerebellar teratoma, immature, 21
- with germinal matrix-like cells, 22
  - with pigmented optic cup-like structures, 23
- Cerebral neuroblastoma, 213–217
- Cervical neuroblastoma, 149, 155
- Cervical teratoma, immature, 22
- Chediak-Higashi syndrome, 195
- Chondromesenchymal hamartoma, nasal, 415–417
- Choriocarcinoma
- clinical findings and pathological findings, 6
  - in newborns and infants, 5, 32–33, 229
- Choroidal seeding, in retinoblastoma, 250
- Choroid plexus papilloma, 201, 225–228
- Chromosomes
- 6p, defect of, in 21-hydroxylase deficiency, 337
  - 8q, defect of, in 11-hydroxylase deficiency, 337
  - 11
    - del 11p13 in Wilms tumor, 268
    - gain of, in congenital fibrosarcoma, 72
    - loss of heterozygosity, association with rhabdomyosarcoma, 83
  - 11p15, loss of heterozygosity, in adrenocortical carcinoma, 341
  - 13
    - retinoblastoma gene on, at q14, 241
  - 13q, deletion in retinoblastoma, rare form, 241
  - 20, gain of, in congenital fibrosarcoma, 72
  - 22, deletions from q11 in rhabdoid tumors, 93
- anomalies of
- association with childhood tumors, 3
  - association with leukemia, 163–165
- See also* Karyotype; Translocations; Trisomies
- Classification
- of astrocytoma, World Health Organization, 207
  - of histiocytosis, 183
  - histological, of hepatoblastoma, 305
  - of leukemia, French-American-British (FAB), 194
  - of liver tumors in the fetus and infant, 304
  - of lung tumors, 393
  - of renal tumors, fetal and infant, 268
- See also* Tumors and tumorlike conditions
- Clear-cell sarcoma, of the kidney, 291–293
- Clinical findings/features
- in adrenocortical tumors, 341
  - in brain tumors, 199–201
  - in hemophagocytic lymphohistiocytosis, 189
  - in hepatoblastoma, 303
  - in infection-associated hemophagocytic syndrome, 192–194
  - of juvenile xanthogranuloma, 196
  - of leukemia, 161–163
  - of malignant melanoma, 124
  - in mesenchymal hamartoma, hepatic, 315–318
  - in neuroblastoma, 137–153
- perinatal, of tumors, 3
  - of renal tumors and tumorlike conditions, 268–269
  - in retinoblastoma, 241–242
- Clinical group stage system, for rhabdomyosarcoma, 93
- Clinicopathological features, of germ cell tumors, 6
- Collagen, type IV, immunoreactivity of fibrosarcomas with, 75
- Complications, clinical, from capillary hemangiomas, 42
- Congenital conditions
- anomalies accompanying bilateral Wilms tumor, 276
  - bilateral retinoblastoma, 244–245
    - with metastases, 246–247
  - cellular mesoblastic nephroma, histogenetic relationship with congenital fibrosarcoma, 72
  - fibrosarcoma of bone, 423
  - generalized fibromatosis, 420
  - giant melanocytic nevi, 121–124
  - multiple fibromatosis. *See* Myofibromatosis
  - pleomorphic adenoma, 371–373
  - pulmonary lymphangiectasis, 391
  - pulmonary myofibroblastic tumor, 400–401
- See also* Autosomal dominant conditions; Autosomal recessive conditions; Familial disorders
- Congenital epulis, 70–71
- Congenital malformations
- adrenocortical cytomegaly associated with, 350–351
  - hepatoblastoma associated with, 303–304
  - tumors associated with, 1
- See also* Autosomal dominant conditions; Autosomal recessive conditions; Congenital conditions
- Congestive heart failure
- association with cardiac tumors, 377
  - fibromas, 381
  - association with hemangioma of the liver, 319
- Cortisol, biosynthesis of, defect in congenital adrenal hyperplasia, 337–341
- Craniopharyngioma, 232
- Cushing syndrome
- association with adrenocortical hyperplasia, 337
  - and insulin-resistant diabetes, association with adrenocortical hyperplasia, 338–339
- Cystadenoma, pancreatic, 360–361
- Cystic adenomatoid malformation, of the lung, 388–390, 395

- Cystic conditions  
 congenital pulmonary, 395–396  
 of the kidney, 294
- Cystic microphthalmia, 257–258
- Cystic partially differentiated  
 neuroblastoma (CPDN), 294
- Cystic Wilms tumor, 294
- Cysts, hepatic, 329
- Cytochemistry, of leukemia, acute  
 lymphoblastic and acute  
 nonlymphoblastic, 169
- Cytogenetics  
 of leukemia, 163–165  
 of retinoblastoma, 241
- Cytokeratin  
 immunoreactivity of  
 with atypical teratoid/rhabdoid  
 tumors, 233  
 with cytotrophoblasts and  
 syncytiotrophoblasts, 32  
 with embryonal carcinoma cells, 31  
 with embryonal hepatoblastoma  
 cells, 307  
 with hepatoblastoma cells, 313  
 with melanotic neuroectodermal  
 tumor of infancy, 99  
 with polyphenotypic small-cell  
 tumors, 103  
 with yolk sac tumors, 25  
 markers, in choroid plexus papilloma,  
 225
- Cytokines, secretion of, by macrophages,  
 183
- Cytomegaly, adrenocortical, 350–351
- Cytotrophoblasts, of choriocarcinomas, 32
- D**
- Delivery, rupture of a tumor at, 1
- Dendritic cells, 183
- Dendrocyte, dermal, origin of juvenile  
 xanthogranuloma in, 196
- Dermal structures, in teratomas, 20
- Desmin, immunoreactivity of  
 with fibrous hamartoma cells, 65  
 with rhabdomyoblasts of  
 ectomesenchymomas, 103
- Desmoplastic cerebral astrocytoma,  
 characteristics of, 207
- Desmoplastic infantile ganglioglioma, 218
- Diagnosis  
 of calcifying aponeurotic fibroma, from  
 chondroid area presence, 70  
 of embryonal rhabdomyosarcoma, 325  
 of fibrous hamartoma of infancy,  
 histological findings, 68–69  
 of malignancy, in young children, 1  
 of neuroblastoma, 156–157
- Differential diagnosis  
 of abdominal masses in the newborn, 2  
 of cranial fasciitis, 107  
 of cutaneous blue nodules, 118  
 of cutaneous infiltrates in the newborn,  
 188  
 of Ewing sarcoma, 420–423  
 of hemophagocytic  
 lymphohistiocytosis, 191–192  
 of leukemia, 165–166  
 of retinoblastoma, 242–243  
 of small blue-cell malignant tumors, 89
- Dopamine  $\beta$ -hydroxylase,  
 immunoreaction of, with melanotic  
 neuroectodermal tumor of infancy,  
 99
- Down syndrome  
 acute megakaryoblastic leukemia  
 (AMKL) associated with, 164  
 leukemia associated with, 163–164  
 congenital, 168–170  
 in fetuses, 161
- Dysgenetic gonads, gonadoblastomas in,  
 29–30
- Dysgerminoma, 32
- Dystocia, maternal, as an indication of a  
 large congenital tumor, 1
- E**
- Ectodermal components, of teratomas, 16
- Ectomesenchymoma, 102–103
- Electron microscopy  
 for diagnosis of brain tumors, 201  
 for diagnosis of neuroblastoma, 156
- Embryoid bodies, association with yolk  
 sac tumors, 28
- Embryonal carcinoma, 31  
 clinical findings and pathological  
 findings, 6
- Embryonal tumors  
 epithelial hepatoblastoma, 307–313  
 medulloepithelioma, 235  
 rhabdomyosarcoma, 83–93  
 of the liver, 325–327  
 sarcoma of the liver, 325–327
- Embryonic disk, teratoma tissues derived  
 from, 16
- Enchondromatosis (Ollier disease),  
 418–419
- Endodermal components, of teratomas, 16
- Enolase, neuron-specific, immunoreaction  
 of, with polyphenotypic small-cell  
 tumors, 103
- Enzymatic defects, related to congenital  
 adrenal hyperplasia, 337
- Eosinophilic granuloma, 188
- Ependymal rosette, in a cerebellar  
 teratoma, 21
- Ependymoma, 201, 222–224
- Epidermal birthmarks, 119–120
- Epidermal nevus/epidermal nevus  
 syndrome, 119
- Epignathus, solid teratoma, 10
- Epithelial membrane antigen,  
 immunoreactivity with, of atypical  
 teratoid/rhabdoid tumors, 233
- Epstein-Barr virus infection, association  
 with lymphomatoid  
 granulomatosis of the lung, 392
- Erythroblastosis, from Rh incompatibility,  
 167
- Erythroleukemia  
 diagnosis of, from glycophorin A  
 analysis, 174  
 trisomies associated with, 164
- Erythrophagocytosis, in hemophagocytic  
 lymphohistiocytosis, 189
- Esterase, immunoreaction of  
 pancreatoblastoma cells with, 360
- Ewing antibody, in primitive  
 neuroectodermal tumors of  
 thoracopulmonary origin, 401–402
- Ewing sarcoma  
 involving bone, 420–423  
 tumors related to histogenetically, 99
- Exocrine tumors, of the pancreas,  
 360–363
- Exophthalmos, in Hand-Schuller-Christian  
 syndrome, 188
- Eye, tumors of, 241–260
- F**
- Factor VIII, immunoreactivity of, with  
 endothelial cells of tufted  
 angioma, 44, 116
- Factor XIIIa, expression of  
 in fibrous histiocytomas, 75  
 in xanthogranuloma cells, 83
- Failure to thrive, in disseminated  
 Langerhans cell histiocytosis, 184
- Familial disorders  
 association of, with brain tumors, 199  
 Chediak-Higaski syndrome, 195  
 hemophagocytic lymphohistiocytosis,  
 189  
 infantile polycystic disease of kidney  
 and liver, 265–266, 333  
 Omenn disease, 194–195  
 retinoblastoma, 241  
 von Hippel-Lindau disease, 38–39  
*See also* Autosomal dominant  
 conditions; Autosomal recessive  
 conditions; Congenital conditions
- Family cancer syndrome,  
 rhabdomyosarcoma in, 83
- Fasciitis, cranial, 105–107
- Fetal alcohol syndrome, association with  
 cystic hygroma, 45
- Fetal rhabdomyomatous Wilms tumor,  
 277
- Fetiform teratoma, 20

- $\alpha$ -Fetoprotein  
 immunoreactivity of, with yolk sac tumors with, 25  
 levels of, in hepatoblastoma, 303  
 in malignant tumor cells, 23  
 normal serum levels of, by age, 29
- Fetus-in-fetu, 20–24  
 clinical findings and pathological findings, 6  
 intrahepatic, 328  
 retroperitoneal, 24
- Fibrohistiocytic tumors, 75–83
- Fibroma  
 calcifying aponeurotic, 70  
 cardiac, 381
- Fibromatosis, 55–60  
 congenital generalized, 420  
 digital, 59–60  
 involving the lung, 395  
 involving the pancreas, 364
- Fibrosarcomas  
 congenital, 72–75  
 differentiating from fibromatoses, 58  
 in the first year of life, 37, 54
- Fibrous connective tissue, neoplasms originating in, 37
- Fibrous connective tissue tumors (myofibroblastic tumors), 54–75
- Fibrous dysplasia, of bone, 418
- Fibrous hamartoma of infancy, 65–69
- Fibrous histiocytoma, 75–81
- Flexner-Wintersteiner rosette, in retinoblastoma, 248, 251
- Focal nodular hyperplasia, of the liver, 329
- French-American-British (FAB) classification, of leukemia, 194
- Fryns syndrome, association with cystic hygroma, 45
- G**
- Ganglioglioma, 218–221
- Ganglioneuroma  
 adrenal, 159  
 maturation of neuroblastoma to, 152
- Gangliorhabdomyosarcoma, 102–103
- Gastric teratoma, 14
- Gaucher disease, lethal, 182
- Genitalia, ambiguous, in congenital adrenal cortical hyperplasia, 337–339
- Genitourinary tract anomalies, association with neuroblastoma in situ, 158
- German Soft Tissue Sarcoma Study Group, 89
- Germ cell tumors, 5–36  
 intracranial neoplasms, 229–231  
 of the liver, 328
- Giant cell astrocytoma, subependymal, 206  
 characteristics of, 207
- Gigantism, of an extremity, in cavernous lymphangioma, 48–49
- Glaucoma, association with Sturge-Weber syndrome, 116
- Glial fibrillary acidic protein (GFAP)  
 antibodies to, in central nervous system neoplasms, 201  
 immunoreactivity with, of atypical teratoid/rhabdoid tumors, 233  
 markers in choroid plexus papilloma, 225  
 neural marker in teratomas, 20
- Glial filament protein, neural marker in teratomas, 20
- Glioblastoma multiforme, 205  
 characteristics of, 207
- Gliomas, predisposition to, association with neurocutaneous syndromes, 1
- Glycophorin A, finding in diagnosis of erythroleukemia, 174
- Gonadal tumors, non-germ cell, 34
- Gonadoblastoma, 29–30  
 clinical findings and pathological findings, 6
- Grading, of neuroblastic tumors, 156. *See also* Staging
- Granular cell tumor (congenital epulis), 70–71
- Granulosa cell tumor, juvenile, 35–36
- Growth, rate of, in the embryo, 1
- H**
- Hamartomas  
 chest wall, 410–415  
 chondromesenchymal nasal, 415–417  
 cutaneous, 120–121  
 myocardial, 384  
 retinal, 252–253
- Hand-Schuller-Christian syndrome, diagnostic criteria for, 188
- Hashimoto-Pritzker syndrome, 188–189
- Hemangioendothelioma, infantile kaposiform, 44, 116
- Hemangioma, 37–39, 113–115  
 capillary, 39–42  
 of the salivary gland, 367–368  
 cardiac, 384  
 cavernous, 42–43  
 of the liver, 303, 319–324  
 lobular capillary, 116  
 pancreatic, 364  
 pulmonary, 395
- Hemangiomas, diffuse, 37
- Hemangiopericytoma, 51–54  
 infantile, comparison with myofibromatosis, 60
- Hematological disorders, in disseminated Langerhans histiocytosis, 184
- Hematopoiesis, during development, 161
- Hematuria, in ossifying renal tumor of infancy, 297
- Hemihypertrophy  
 adrenocortical tumors associated with, 337, 341  
 hepatoblastoma associated with, 303
- Hemophagocytic lymphohistiocytosis, 189–192  
 secondary, 192–194
- Hemophagocytic syndromes, 189–194
- Hepatic tumors, miscellaneous, 329–333
- Hepatoblastoma, of the liver, 303–315
- Hepatocellular carcinoma, 329–330  
 differentiating from embryonal epithelial hepatoblastoma, 307–312
- Histiocyte, differentiation of, 181
- Histiocyte Society, 183
- Histiocytic Disorders, German/Austrian Registry for, 189
- Histiocytic lymphoma, 194
- Histiocytoses, 181–198
- Histiocytosis X. *See* Langerhans cell histiocytosis
- Histogenesis, of retinoblastoma, 248
- Histological findings  
 in cardiac fibromas, 381  
 in embryonal rhabdomyosarcoma, 325  
 in hyperinsulinemic hypoglycemia of infancy, 353–356  
 in pancreatoblastomas, 360  
 in sialoblastoma, 369–370  
 in soft tissue tumors, 38–39
- Histological patterns, in clear-cell sarcoma of the kidney, 291
- Histological types  
 of congenital brain tumors, 200  
 of germ cell neoplasms, World Health Organization classification, 5–7  
 of rhabdomyosarcomas, 83–93
- Histopathology  
 of neurocutaneous melanosis, 124–125  
 of teratomas, 16–20
- HMB45, immunoreactivity of, with melanotic neuroectodermal tumor of infancy, 99
- Human chorionic gonadotropin, immunoreactivity of, with syncytiotrophoblasts, 32
- Human placental lactogen, immunoreactivity of, syncytiotrophoblasts, 32
- Hydrocephalus  
 in astrocytoma, 202  
 in ependymoma, 222
- Hydronephrosis, 262–263, 294
- Hydrops fetalis  
 association of  
 with cardiac teratomas, 381–384  
 with cystic hygroma, 45

- with neuroblastoma, 137
  - as an indication of a congenital tumor, 1
  - 21-Hydroxylase, deficiency of, in
    - congenital adrenal hyperplasia, 337–338
  - Hygromas, 44–50
    - of the neck, 46
  - Hyperinsulinemic hypoglycemia of
    - infancy, defined, 353
  - Hypertriglyceridemia
    - in hemophagocytic lymphohistiocytosis, 189
  - Hypofibrinogenemia, in hemophagocytic lymphohistiocytosis, 189
  - Hypogammaglobulinemia, association with lymphomatoid granulomatosis of the lung, 392
- I**
- Imaging, for antenatal diagnosis of tumors, 1
  - Immune defect, association with sinus histiocytosis, 194
  - Immunocompromise, in hemophagocytic lymphohistiocytosis, 192–194
  - Immunohistochemical features
    - of atypical teratoid/rhabdoid tumors, 233
    - of brain tumors, 201–202
    - of familial erythrophagocytic lymphohistiocytosis, 190
    - of primitive neuroectodermal tumors, 99
    - of retinoblastomas, 248
    - of yolk sac tumors, 25
  - Immunoperoxidase studies, to confirm diagnosis of neuroblastoma, 156
  - Immunophenotypes, in congenital leukemia, 163
  - Immunoreactivity
    - of choriocarcinomas, 32
    - of fibrosarcomas, with vimentin and collagen type IV, 75
    - of giant melanocytic nevi, 121
    - of juvenile xanthogranulomas, 132
    - in Langerhans cell histiocytosis, 130
    - of neuroblastoma skin nodules, 129
    - of tufted angiomas, 116
  - Incidence
    - of brain tumors, 199
    - of hemangiopericytoma, in children relative to adults, 51
    - of neuroblastoma, 137
    - of torticollis, 60
    - of tumors of the newborn, in the United Kingdom, 1–4
  - Infantile polycystic disease, 329
    - of the kidneys and liver, 333
  - Infantile polycystic disease of kidney and liver, 265–266
  - Infection-associated hemophagocytic syndrome, 192–194
  - Inflammatory pseudotumor, of the lung, 393–394
  - Insulin-resistant diabetes, association with adrenocortical hyperplasia and Cushing syndrome, 338–339
  - Interferon, 183
  - Intergroup Rhabdomyosarcoma Study
    - association of congenital malformations with rhabdomyosarcoma found in, 83
    - classification of tumors by, 89–93
  - International Staging System Working Party, 153
  - Intestinal adenomatous polyposis, hepatoblastoma associated with, 303
  - Involuting phase, of capillary hemangiomas, 39–42
  - Islet cells
    - adenoma of, 359–360
    - types of, in pancreatic hyperplasia, 356–359
- J**
- Juvenile xanthogranuloma, 81–83
    - cutaneous, 131–132
    - origin in dendrocytes, 196
- K**
- Kaposiform hemangioendothelioma
    - association with hemangioma of the liver, 319
    - infantile, 116
  - Karyotype
    - in congenital leukemia, 163
    - mosaic, 21 in transient leukemia, 169
    - 45,X0/46,XY in gonadoblastoma, 29
    - in patients with gonadoblastomas, 29–30
    - in patients with gonadoblastomas, 29–30
    - trisomy 11, in cellular congenital mesoblastic nephroma, 285
    - X0, in patients with Turner syndrome, 45–50
    - See also* Chromosomes; Translocations; Trisomies
  - Kasabach-Merritt syndrome, 42
    - association of
      - with infantile kaposiform hemangioendothelioma, 44
      - with hemangioma of the liver, 319, 322
      - with tufted angioma, 116
  - Keratin, immunoreaction of
    - pancreatoblastoma cells with, 360
  - 17-Ketosteroids, secretion of, in adrenocortical carcinoma, 341
  - Kidney
    - development of, 270
    - multicystic dysplastic, 264
  - Kidney tumors, 261–301
  - Klippel-Trenaunay-Weber syndrome, generalized hemangioma accompanying, 42, 116
  - Kupper cells, differentiation of histiocytes to, 181
- L**
- Langerhans cell histiocytosis, 183–189
    - bone involvement in, 408–409
    - skin lesions in, 130
  - Langerhans cells
    - characteristic appearance in Letterer-Siwe disease, 186–189
    - differentiation of histiocytes to, 181
  - Laryngotracheal papilloma, 396–397
  - Lectin I, *Ulex europaeus*, immunoreactivity of, with endothelial cells of tufted angioma, 44
  - Leptomeningeal hemangiomatosis, association with Sturge-Weber syndrome, 113–116
  - Leptomeningeal lymphohistiocytic infiltration, in hemophagocytic lymphohistiocytosis, 189–190
  - Leptomeningeal melanoma, association with neurocutaneous melanosis, 124–125
  - Letterer-Siwe disease, 130, 183–189
  - Leukemia, 161–180
    - differential diagnosis of, 89
    - incidence of, in newborns in the United Kingdom, 1–2
    - See also* Acute . . . leukemia entries
  - Leukemia cutis, 127
    - in myelomonoblastic leukemia, with translocations, 164
  - Leukemoid reaction, association with infection and trisomies 18 and 21, 364
  - Leukocoria, in retinoblastoma, 241
  - Leukocytophagocytosis, in sinus histiocytosis, 194
  - Li-Fraumeni syndrome, adrenocortical tumors associated with, 337
  - Lipoblastoma, 94–96
  - Lipoma, 94–95
  - Liver
    - enlargement of, in hemophagocytic lymphohistiocytosis, 189
    - fetal, hematopoiesis in, 162
    - metastasis of yolk sac tumor to, 28
    - sacroccygeal teratoma with, 19
    - tumors of, 303–335
      - antenatal diagnosis with ultrasonography, 1
      - See also* Hepat-entries
  - Lobular capillary hemangioma, 117–118

- Location  
 of fetus-in-fetu, 20  
 of lymphangiomas, 45–50  
 of neuroblastomas, primary and metastatic, 150  
 of teratomas, 5
- Loss of heterozygosity  
 chromosome 11, association with rhabdomyosarcoma, 83
- Lung  
 metastasis of yolk sac tumor to, 28  
 tumors of, 387–405
- Lymphadenopathy, sinus histiocytosis with, 194
- Lymphangiectasis, 51
- Lymphangioma, 44–50, 118
- Lymph node biopsy, in disseminated Langerhans histiocytosis, 184
- Lymphocytes, infiltration of the pancreas, 364
- Lymphokines, effect of macrophages on, 183
- Lymphoma, histiocytic, 194
- Lymphomatoid granulomatosis, 392
- Lysozyme, immunoreaction of, with embryonal sarcoma of the liver, 325
- M**
- Macrophage antigens, expression of, in xanthogranuloma cells, 83
- Macular stains, port wine and nevus flammeus, 113
- Maffucci-Hast syndrome  
 enchondromatosis of the bone in, 418  
 vascular tumors associated with, 39
- Magnetic resonance imaging  
 to detect optic nerve gliomas, 253  
 of kaposiform hemangioendotheliomas, 44
- Malformations, vascular, 43–44
- Malignancy  
 differentiating from immature normal structures, 1  
 of fibrosarcoma, 54  
 of soft tissue tumors, 37  
 of yolk sac tumor, 25
- Malignant histiocytosis, 194
- Malignant melanoma, 124–125
- Malignant peripheral nerve sheath tumor (malignant schwannoma), in triton tumors, 104
- Malignant tumors, metastasis to the pancreas, 364
- Masculinizing adrenal cortical tumor, 346
- Mast cell disease, 132–133
- McCune-Albright syndrome, polyostotic fibrous dysplasia in, 418
- Mediastinum  
 lymphangioma of, 50
- neuroblastoma of, 146–148  
 teratoma of, 13, 17
- Medulloblastoma, 201, 210–212
- Medulloepithelioma, 235  
 of the eye, 252
- Melanocytosis, 124–125
- Melanotic neuroectodermal tumor of infancy, 99–102
- Meningeal tumors, 235
- Mesenchymal cystic hamartoma, of the lung, 396
- Mesenchymal hamartoma, of the liver, 303, 315–318
- Mesoblastic nephromas  
 antenatal diagnosis of, using ultrasonography, 1  
 congenital, 261, 280–287  
 multilocular cysts associated with, 294
- Metanephric blastema, differentiation of embryonic nephrogenic tissue into, 270
- Metastasis  
 of adrenocortical carcinomas, 350  
 of germ cell tumors, to the liver, 328  
 of hemangiopericytomas, 51  
 of hepatoblastomas, common sites of, 315  
 to the liver, in newborns and infant, 303  
 of neuroblastoma, to the placenta, 137–141  
 to the pancreas, of malignant tumors, 364  
 of retinoblastomas, 248–251  
 of yolk sac tumors, 25
- Microphthalmia, cystic, 257–258
- Mitosis-karyorrhexis index (MKI), of neuroblasts, and prognosis in neuroblastoma, 156
- Monophagocytic system, ontogeny of, 181–183
- Monosomy, 7, association with leukemia, 163
- Morphological features, of leukemia, acute lymphoblastic and acute nonlymphoblastic, 169
- Mortality, from non-malignant tumors, neonatal, 1
- Mosaic. *See* Karyotype, mosaic
- Mucoepidermoid tumor, of the salivary gland, 374
- Multicystic kidney, 294
- Multilocular cyst (MLC), cystic partially differentiated nephroblastoma (CPDN), 294
- Multilocular cystic nephroma (MLCN), 294
- Multiple vascular leiomyomas. *See* Myofibromatosis
- Myelin basic protein, neural marker in teratomas, 20
- Myofibroblastic tumor, 54–75  
 congenital pulmonary, 400–401
- Myofibroblasts, in fibromatoses, 58
- Myofibromatosis, 60–65  
 infantile, 420
- Myxoma, of the heart, 381
- N**
- Nasopharyngeal dermoid, 12  
 differentiating from congenital pleomorphic adenoma, 371–373
- National Cancer Survey, on malignant liver tumors in the first year of life, 303
- National Wilms Tumor Study (NWTs), 261, 278  
 clear-cell sarcoma of the kidney found in, 291
- Nephroblastomatosis, renal, prognosis in, 278–279
- Nephrogenic rests, 278–279
- Nephrogenic tissue, in teratomas, 20
- Nerve growth factor receptor, neural marker in teratomas, 20
- Nerve tumors, malignant peripheral, association with neurocutaneous syndromes, 1
- Nesidioblastosis, 353–359
- Nesidiodyplasia, 353–359
- Neural crest, as the origin  
 of giant congenital melanocytic nevi in, 121–124  
 of neuroblastoma in, 137–161  
 of peripheral nervous system tumors in, 95  
 of primitive neuroectodermal tumors in, 207
- Neural elements, of giant melanocytic nevi, 124
- Neural markers, of teratomas, 20
- Neuroblastoma, 137–161  
 antenatal diagnosis of, using ultrasonography, 1  
 cerebral, 213–217  
 differential diagnosis of, 89  
 incidence of, in newborns in the United Kingdom, 1–2  
 skin manifestations of, 128
- Neuroblastoma in situ, 157–158
- Neurocutaneous melanosis, 124–125
- Neurocutaneous syndromes, neoplasias associated with, 1
- Neuroectodermal tumor, peripheral, of thoracopulmonary origin, 401–403
- Neurofibroma, 97–99
- Neurofibromatosis, 125–126
- Neurofibromatosis I  
 central nervous system findings in, 200  
 of the eye, 253–255  
 neurofibromas associated with, 97

- rhabdomyosarcoma associated with, 83  
triton tumor associated with, 104
- Neurofilament protein (NFP)  
antibodies to, in central nervous system neoplasm, 201  
immunoreactivity of, with neuroblastoma, 156  
neural marker in teratomas, 20
- Neuroglial tissue, of teratomas, 17
- Neurological signs, in hemophagocytic lymphohistiocytosis, 189
- Neuron-specific enolase (NSE)  
immunoreactivity of  
with melanotic neuroectodermal tumor of infancy, 99  
with neuroblastomas, 129, 152, 156  
neural marker in teratomas, 20
- Nevus  
melanocytic, 121–125  
sebaceous, 119–120
- Nevus flammeus, 39, 113–114
- Nevus venosus (port wine stain), 39, 113–114
- Noonan syndrome, association with cystic hygroma, 45
- O**
- Oligohydramnios, association with cystic hygroma, 45
- Ollier disease, enchondromatosis, 418–419
- Omenn disease, 194–195
- Ontogeny, of the monophagocytic system, 181–183
- Optic glioma, 203, 253
- Orbital tumors, 256–258
- Organomegaly, in leukemia, 171–173
- Ossifying renal tumor of infancy, 297–299
- Osteoid, in hepatoblastomas, 313
- Ovary, follicular cyst of, 34
- P**
- Pancreas  
developing, 354  
normal, nesidioblastosis in, 355  
sacrococcygeal teratoma with, 19
- Pancreatic tumors, 353–365
- Pancreatoblastoma, 353, 360–363  
defined, 356–359
- Papilloma  
choroid plexus, 225–228  
laryngotracheal, 396–397
- Papillomatosis, of the larynx, 396–397
- Papillomavirus, human, 396–397
- Pathology  
of brain tumors, 201–202  
of congenital mesoblastic nephroma, 280–287  
of Cushing syndrome associated with adrenal cortical nodular hyperplasia, 338  
of disseminated Langerhans cell histiocytosis, 186–189  
of hemophagocytic lymphohistiocytosis, 189–192  
of leukemia, 171–174  
of neuroblastoma, 153–157  
of retinoblastoma, 243–251  
of rhabdoid tumor of the kidney, 288–290  
of Wilms tumor, 271–278
- Pericardial effusion, association with pericardial teratoma, 381–384
- Pericardial teratoma, 377
- Peripheral nervous system tumors, 95–107
- Peripheral neuroepithelioma, tumors related to histogenetically, 99
- Petechiae, conditions causing, in the newborn, 163
- P-glycoprotein transmembrane protein, in rhabdomyosarcoma, prognosis from, 83
- Phagocytic cells, 183
- Phakomatoses, neoplasia associated with, 1
- Pharyngeal teratoma, 11
- Pierre-Robin syndrome, association with nephrogenic rests, 279
- Pigmented congenital epulis. *See* Melanotic neuroectodermal tumor of infancy
- Pigmented neuroectodermal tumor. *See* Melanotic neuroectodermal tumor of infancy
- Pilocystic astrocytoma, grade I, characteristics of, 207
- Pineal, involvement of, along with bilateral retinoblastoma, 252
- Pleomorphic adenoma, salivary gland, 372–374
- Pleuropulmonary blastoma, arising from lung cysts, 395, 398–400
- Plexiform neurofibroma, 97–98, 125  
involving the salivary gland, 367–368
- Polycystic disease, infantile, of kidney and liver, 265–266
- Polyembryoma, 32  
clinical findings and pathological findings, 6
- Polyphenotypic small-cell tumor, 103–104
- Polysialic acid, neural marker in teratomas, 20
- Port wine hemangioma, in Sturge-Weber syndrome, 235–237
- Port wine nevus, 113–114
- Pregnancy-specific  $\beta_1$ -glycoprotein, immunoreactivity of syncytiotrophoblasts and cytotrophoblasts with, 32
- Presacral teratoma, with embryonal carcinoma and yolk sac tumor, 31
- Primitive gut, sacrococcygeal teratoma with, 19
- Primitive neuroectodermal tumor (PNET), 99, 207–209  
association with brain tumors, 199  
cerebellar, 210–212  
cerebral, 213–217  
differential diagnosis of, 89  
differentiating from pulmonary blastoma, 399–400  
in the first year of life, 37  
retinoblastoma as, 248  
of thoracopulmonary origin, 401–403
- Prognosis  
in adrenocortical carcinoma, 347, 350  
in atypical teratoid/rhabdoid tumor, 233  
in botryoid embryonal rhabdomyosarcoma, 83–84  
in brain tumors, 238  
survival rates, 210  
in cerebral neuroblastoma, 213  
in cervical cystic hygroma, prenatal, 45–50  
in choriocarcinoma, 32  
in choroid plexus papilloma, 225  
in clear-cell sarcoma of the kidney, 291  
in congenital and childhood hemangiopericytoma, 51  
in cystic adenomatoid malformation of the lung, 395  
in desmoplastic infantile ganglioglioma, 218  
in ganglioneuroma, 159  
in generalized fibromatosis with pulmonary involvement, 395  
in hemangioma of the liver, 319  
in hemophagocytic lymphohistiocytosis, 189  
in hepatoblastoma, 315  
in immature teratoma, 20  
in infantile fibrosarcoma, 75  
in infantile kaposiform hemangioendothelioma, 116  
in intracranial teratoma, 229  
in Langerhans cell histiocytosis, 189  
in leukemia, 161  
in medulloepithelioma, 252  
in melanotic neuroectodermal tumor of infancy, 99  
in meningeal tumors, 235  
in multicentric myofibromatosis, 60–65  
in neuroblastoma, 153, 159  
mitosis-karyorrhexis index for evaluating, 156  
in Omenn disease, 195  
in panrenal nephroblastomatosis, 278  
in polyphenotypic small-cell tumor, 103  
in primitive neuroectodermal tumors, 99, 401  
in retinoblastoma, 248, 252

- Prognosis (*Continued*)  
 in rhabdoid tumor, 93  
 in rhabdomyosarcoma, 83, 89, 93  
 in sialoblastoma, 369  
 in triton tumor, 104  
 in Wilms tumor, 273  
 in xanthogranuloma, 83
- Proliferative myositis, 105
- Prune belly syndrome, 267
- Pulmonary involvement, in Letterer-Siwe disease, 184–185
- Purpura, conditions causing, in the newborn, 163
- Pyogenic granuloma, 117–118
- R**
- Recurrence  
 of calcifying aponeurotic fibroma, 70  
 of fibromatosis, clinical features associated with, 58–60  
 of lipoblastomas, 94–95  
 of melanotic neuroectodermal tumor of infancy, 99  
 of sialoblastomas, 369
- Reese-Ellsworth staging system, for retinoblastoma, 242
- Regression  
 of congenital hemangiomas, 37–39, 113  
 of the liver, 319–324  
 of congenital hemangiopericytomas, 51  
 of granular cell tumors, 70  
 of juvenile xanthogranulomas, 81, 83, 131  
 of myofibromatosis, 60  
 of nephrogenic rests, 278  
 of neuroblastomas, 150–151, 157  
 of retinoblastoma, 243  
 of skin lesions in Hashimoto-Pritzker disease, 188–189
- Renal tumors, 261–301
- Rendu-Osler-Weber disease, lung involvement in, 395
- Retina, astrocytic tumors of, 252–255
- Retinal anlage tumor. *See* Melanotic neuroectodermal tumor of infancy
- Retinal detachment, in retinoblastoma, 243
- Retinoblastoma, 241–252  
 congenital, spontaneous regression of, 157
- Retroperitoneal neuroblastoma, 144–145
- Retroperitoneal teratoma, 15
- Rhabdoid tumor  
 association with brain tumors, 199  
 differential diagnosis of, 89  
 in the first year of life, 37, 93  
 of the kidney, 261, 288–290  
 of the skin, 129
- Rhabdomyoblasts  
 as the source of rhabdomyosarcoma, 83  
 in triton tumors, 104
- Rhabdomyoma, 83  
 cardiac, 377–380
- Rhabdomyomatosis, of the lung, 401
- Rhabdomyosarcoma  
 botryoid form, 83–85  
 differential diagnosis of, 89  
 embryonal, 83–93  
 of the liver, 325–327  
 of the eyelid and orbit, 256–258  
 in the first year of life, 37  
 origin of  
 in lung cysts, 395  
 in mesenchymal cystic hamartoma of the lung, 396  
 of the skin, 129
- Rosai-Dorfman disease, 194
- Rubeosis iridis, in retinoblastoma, 241
- S**
- S-100 protein, immunoreactivity of  
 with chondromesenchymal hamartoma cells, 415  
 with fibrous hamartoma cells, 65  
 in Langerhans cell histiocytosis, 130  
 with neuroblastoma cells, 156
- Sacrococcygeal teratoma  
 cystic, 8  
 immature, yolk sac tumor cells in, 23  
 mature, with pancreas, 19  
 with small bowel and cartilage components, 18  
 solid, 7  
 yolk sac tumor cells in, 28
- Salivary gland, tumors of, 367–375
- Salivary gland anlage tumor, 371–373
- Salt-wasting type, of congenital adrenal hyperplasia, 338, 340
- Sarcoma, breast and brain tumor, leukemia, laryngeal, and lung cancer, and adrenal cortical carcinoma (SBLA) syndrome, 341
- Sarcomas  
 incidence of, in newborns in the United Kingdom, 1–2  
 primary, of the liver, 325–327
- Schwannian stroma, for categorizing neuroblastomas, 156
- Sex  
 and incidence of clear-cell sarcoma of the kidney, 291  
 and incidence of fibromatosis, 55  
 and incidence of islet cell adenoma, 359–360  
 and incidence of meningeal tumors, 235
- and incidence of pleuropulmonary blastoma, 398–400  
 male, yolk sac tumors of, 5
- Sialoblastoma, 367, 369–370  
 differentiating from salivary gland anlage tumor, 371
- Sinus histiocytosis, with massive lymphadenopathy, 194
- Sinusoidal lining cells, differentiation of histiocytes to, 181
- Skeletal malformations, association with neurofibromatosis type I, 99
- Skeletal muscle tumors, 83–93  
 of the lung, 401
- Skin  
 involvement of  
 in congenital leukemia, 174  
 in leukemia, 161–163  
 lesions of, in Letterer-Siwe disease, 183–184  
 tumors and tumorlike conditions of, 113–135
- Small blue-cell malignant tumors, differential diagnosis of, 89
- Smooth muscle hamartoma, 120–121
- Soft tissue tumors, 37–111
- Sonography  
 for diagnosis of diffuse nephroblastomatosis antenatally, 278–279  
 for diagnosis of intracranial teratomas, 229  
 prenatal, appearance of neuroblastoma on, 137
- Spider cells, in cardiac rhabdomyoma, 378–379
- Spindle cell neoplasms, fibromatoses as, 58
- Spindle cell rhabdomyosarcoma, 89–90
- Spleen, enlargement of, in hemophagocytic lymphohistiocytosis, 189
- Staging  
 of adrenal cortical carcinoma, 350  
 of hepatoblastoma, 315  
 of neuroblastoma, 153  
 of retinoblastoma, 242  
 of rhabdomyosarcoma, 93  
 of Wilms tumor, 278
- Stillbirth, in antenatal leukemia, 161
- Strabismus, in retinoblastoma, 241
- Strawberry hemangioma, 115
- Sturge-Weber syndrome, hemangiomas associated with, 37, 113–116, 235–237
- Subcutaneous fat, necrosis of the newborn, 126
- Surface marker analysis, to characterize cell types in leukemia, 174

- Symptoms, of neonatal brain tumors, 200
- Synaptophysin  
immunoreactivity of  
with ectomesenchymoma cells, 103  
with melanotic neuroectodermal tumor of infancy, 99  
with neuroblastoma, 156  
neural marker in teratomas, 20
- Syncytiotrophoblasts, of  
choriocarcinomas, 32
- Syndromes, associated with Wilms tumor, 269
- T**
- T cells, presentation of antigens to, by  
Langerhans cells, 183
- Telangiectasia, hereditary, 395
- Teratoid/rhabdoid tumor, atypical, 233–234
- Teratoma, 5–25  
antenatal diagnosis of, using  
ultrasonography, 1  
cardiac, 381–384  
clinical findings and pathological findings, 6  
gastric, 14  
immature, defined, 20  
intracranial, 201, 229–231  
of the liver, 328  
pericardial, 377  
presacral, with embryonal carcinoma and yolk sac tumor, 31  
skin, neuroglial tissue, and choroid plexus of, 16  
testicular, trabeculae of bone from, 18  
*See also* Sacrococcygeal teratoma
- Testis  
granulosa cell tumor of, 35–36  
neonatal tumors of, 34  
teratoma of, 16  
yolk sac tumor of, 26–28
- Tetralogy of Fallot, gonadoblastoma in a patient with, 30
- Thyrocervical teratoma, 9
- Torticollis, 60–61
- Touraine syndrome, 124–125
- Touton giant cells, in juvenile xanthogranulomas, 82–83, 132, 196
- Tracheoesophageal fistula, association with neuroblastoma in situ, 158
- Transient leukemia, in Down syndrome, 168
- Translocations  
in acute megakaryoblastic leukemia {t(1;22)(p13;q13)}, 164  
in acute monoblastic leukemia {t(2;22)(p13;q13)}, 165  
{t(8;16)(p11;q13)}, 164  
{t(9;11)(p21;q23)}, 164  
in acute nonlymphoblastic leukemia {t(9;11)(p22;q23)}, 164  
in alveolar rhabdomyosarcoma {t(2;13)(q35;q14)}, 83, 89–93  
in anaplastic hepatoblastoma t(10;22), 313  
in congenital leukemia, 163–165  
in embryonal sarcoma of the liver, 19q, 325–327  
in Ewing sarcoma, {t(11;22)(q24;Q12)}, 420–423  
in neuroectodermal tumor of thoracopulmonary origin, t(11;22), 401  
in primitive neuroectodermal tumors {t(11;22)(q24;q11-12)}, 99  
*See also* Chromosomes; Karyotypes; Trisomies
- Trilateral retinoblastoma, 252
- Trisomies  
7, in congenital pigmented nevi, 124  
8, erythroleukemia associated with, 164  
11  
fibrosarcomas associated with, 75  
mesoblastic nephroma associated with, 280, 285  
13, cystic hygroma associated with, 45  
18  
cystic hygroma associated with, 45  
leukemoid reaction associated with, 364  
neuroblastoma in situ associated with, 158  
19, erythroleukemia associated with, 164  
21  
acute megakaryoblastic leukemia associated with, 164  
cystic hygroma associated with, 45  
bilateral, 48  
erythroleukemia associated with, 164  
leukemoid reaction associated with, 364  
*See also* Chromosomes; Karyotype; Translocation
- Triton tumor, 104
- Tuberous sclerosis, 126, 206  
age of appearance of lesions of, 200  
association with cardiac rhabdomyomas, 377–378  
retinal, 252–253
- Tumors and tumorlike conditions  
of the adrenal cortex, 338  
of bone, 409  
of the heart, 378  
of the pancreas, 354  
of the salivary glands, 369
- Turner syndrome  
bilateral cystic hygromas associated with, 48  
lymphangiomas associated with, 45–50
- Twinning, structures resulting from  
abortive attempts at, 5. *See also* Fetus-in-fetu
- U**
- Ultrasonography  
for identification of retinoblastoma in utero, 241  
prenatal  
for fetus-in-fetu detection, 20  
for hepatoblastoma detection, 303  
for Wilms tumor detection, 272
- Umbilical cord, single-vessel, association with cystic hygroma, 45
- Undifferentiated carcinoma, of the submaxillary gland, 374
- Unilocular cyst, of the liver, 332
- Urticaria pigmentosa, 132–133
- V**
- Vascular conditions  
of the lungs, 395  
malformations associated with syndromes in newborns, 113–116  
malformations of the brain associated with, 235–237  
of the skin, 113–119
- Vascular invasion, in choriocarcinoma, 32–33
- Vascular tumors, 37–54
- Venous angiomas, as vascular malformations, 43–44
- Verocay body pattern, of clear-cell sarcoma of the kidney, 291
- Vimentin immunoreactivity  
with congenital fibrosarcomas, 77–78  
with atypical teratoid/rhabdoid tumors, 233  
with chondromesenchymal hamartoma cells, 415  
with embryonal hepatoblastoma cells, 307  
with embryonal sarcoma of the liver, 325  
with fibromatosis, 58  
with fibrosarcomas, 75  
with fibrous hamartoma cells, 65  
with fibrous histiocytomas, 75–77  
with granular cell tumors, 70  
with juvenile xanthogranulomas, 132  
with melanotic neuroectodermal tumor of infancy, 99  
with myofibromatosis cells, 60  
with ossifying renal tumor of infancy cells, 297

Vimentin immunoreactivity (*Continued*)  
with polyphenotypic small-cell  
tumors, 103  
with rhabdomyosarcoma cells, 89  
with xanthogranuloma cells, 83  
Viral infections, infection-associated  
hemophagocytic syndrome  
associated with, 192–194  
Virilization  
in adrenocortical carcinoma, 347  
in hydroxylase deficiency defects, 338

von Hippel-Lindau disease, hemangiomas  
associated with, 37–39  
Von Recklinghausen disease, 97, 125–126

**W**

Wilms tumor, 261, 268–278, 303  
association with brain tumors, 199  
cystic, 294  
World Health Organization (WHO)  
classification of astrocytoma by, 207  
classification of germ cell tumors by, 5–7

**X**

Xanthogranuloma, juvenile, 81–83, 196

**Y**

Yolk sac tumor (endodermal sinus tumor),  
5, 25  
clinical findings and pathological  
findings, 6  
elements resembling, in a teratoma, 23  
of the liver, 328