

# Index

## A

- Acrophobia
  - clinical findings, 177–178
  - course and therapy
    - behavioral therapy, 181
    - physiological visual height imbalance, 179, 180
    - subjective visual height intolerance, 178, 180
  - pathogenesis, 178
  - prevalence, 178
- Alternobaric vertigo, 149
- Anterior canal benign paroxysmal positioning vertigo (ac-BPPV), 65

## B

- Basilar type migraine
  - clinical aspects and course, 132–134
  - differential diagnosis, 136–138
  - pathophysiology, 134–135
  - patient history, 131–132
  - pragmatic therapy, 135–136
- Benign paroxysmal positioning vertigo (BPPV)
  - anterior canal (ac-BPPV), 65
  - central positional vertigo/nystagmus, 67–68
  - clinical features and course, 54–55
  - differential diagnosis, 67
  - hc-BPPV
    - canalolithiasis and cupulolithiasis, 64
    - persistent hc-BPPV, 65
    - vs. posterior BPPV, 63–64
    - therapy, 65–66
  - pathomechanism
    - course of attacks, 57
    - direction of nystagmus, 57–58

- duration, 57
- latency, 55
- patient history, 53
- pragmatic therapy
  - additional medication, 63
  - Epley maneuver, 61–62
  - ineffective measures, 63
  - physical liberatory maneuvers, 58–60
  - recurrences after, 62
  - self-treatment, 62
  - Semont maneuver, 60–61
  - surgery, 63
- Semont liberatory maneuver, 56
- Bilateral vestibulopathy
  - causes of, 95
  - cervical vestibular-evoked myogenic potentials, 41
  - clinical aspects and natural course, 90–91
  - differential diagnosis, 94
  - etiology, 92
  - pathophysiology
    - oscillopsia and blurred vision, 91
    - postural vertigo, 91
    - posture and gait unsteadiness, 91
    - spatial memory and navigation deficits, 92
  - patient history, 89–90
  - pragmatic therapy
    - patient education, 94
    - physical therapy, 93
    - recovery, 93
    - vestibular function
      - prevention, 93
    - therapeutic principles, 92
  - Brandt-Daroff maneuver, 58, 60

**C**

- Caloric testing, 35–36, 90
- Car sickness, 175
- Central vestibular syndromes
  - central vestibular structure
    - ascending pathways, 112–114
    - descending pathways, 114
    - pathological stimuli, 114
    - vestibulo-ocular reflex, 112, 113
  - clinical aspects, 114
  - course and pathophysiology, 114–115
  - differential diagnosis, 115
  - higher vestibular functional disorders, 111–112
  - vs. peripheral vertigo, 115–116
  - vestibular migraine
    - clinical aspects and course, 132–134
    - differential diagnosis, 136–138
    - pathophysiology, 134–135
    - patient history, 131–132
    - pragmatic therapy, 135–136
  - vestibulo-ocular reflex
    - horizontal (yaw) plane, 118–120
    - neuronal pathways, 112, 113
    - sagittal (pitch) plane, 119–125
    - vertical (roll) plane, 126–131
- Cervical vestibular-evoked myogenic potentials (cVEMPs)
  - bilateral vestibulopathy, 41
  - evaluation criteria, 40
  - Menière's disease, 42
  - superior canal dehiscence syndrome, 41
  - vestibular neuritis, 41
  - vestibular schwannoma, 41
  - vestibulo-colic reflex, 39
- Cervicogenic vertigo, 172–173
- Childhood vertigo
  - benign paroxysmal vertigo
    - basilar-type migraine, 170
    - episodic vertigo, 165, 169–170
    - therapy, 170
  - central vestibular signs, 165
  - differential diagnoses
    - attacks of vertigo, 168–169
    - stance and gait imbalance, 169
    - sustained vertigo, 169
  - episodic ataxias
    - acetazolamide and 4-aminopyridine, 170–171
    - symptoms, 170
    - types, 170

- motion sickness, 171
- therapy, 166–168

- Cogan's syndrome, 76
- Computed tomography (CT)
  - perilymph fistula, 96
  - petrous bone, 45–46
- Convergence reaction, 19–20
- Cover tests
  - alternating cover test, 16
  - one-eye cover test, 15
  - uncover test, 15–16

**D**

- Dix-Hallpike positioning
  - maneuver, 29–30
- Dizziness. *See* Vertigo/dizziness
- Downbeat nystagmus syndrome (DBN)
  - 3,4-diaminopyridine, 126
  - idiopathic DBN, 123–124
  - therapy, 124–125
  - visual and vestibulocerebellar ataxia, 122
- Drug-induced vertigo, 171–172

**E**

- Electronystagmography (ENG)
  - electrode placement, 36, 38
  - rotatory chair and rotatory drum, 36, 39
- Epley maneuver, 61–62

**F**

- Finger-pointing test, 32

**H**

- Head-impulse test, 12, 90, 119
  - bilateral vestibulopathy, 90
  - horizontal VOR, 25–27
  - stroke, 116
  - vestibular neuritis, 76
  - video-oculography, 36, 37
- Head tilt measurement, 14
- Height vertigo. *See* Acrophobia
- Horizontal canal benign paroxysmal positioning vertigo (hc-BPPV)
  - canalolithiasis and cupulolithiasis, 64
  - persistent hc-BPPV, 65
  - vs. posterior BPPV, 63–64
  - therapy, 65–66

**L**

- Labyrinthine failure
  - clinical manifestations, 147–148
  - corticosteroids, 148
  - hemorrhage and petrous bone fracture, 146–147
  - vestibular training program, 148

**M**

- Magnetic resonance imaging (MRI)
  - fourth nerve palsy, 131
  - petrous bone and cerebellopontine angle, 46, 48
  - upbeat nystagmus syndrome, 125
  - vestibular pseudoneuritis, 119
- Mal-de-debarquement syndrome, 175–176
- Menière's disease
  - cervical vestibular-evoked myogenic potentials, 42
  - clinical problems, 83–84
  - clinical syndrome and course
    - certain Menière's disease, 77
    - definite Menière's disease, 77
    - possible Menière's disease, 78–79
    - probable Menière's disease, 77
  - differential diagnosis, 83
  - etiology, 79
  - pathophysiology, 79
  - patient history, 77
  - pragmatic therapy
    - acute attack, 82
    - prophylactic treatment, 82
    - vestibular drop attacks, 82–83
  - therapy
    - betahistine, 81
    - gentamicin, 80
    - glucocorticoids, 80–81
- Mismatch theory, 174
- Modified Epley liberatory maneuver, 59–60
- Motion sickness
  - course and therapy
    - antivertiginous drugs, 176
    - extreme acceleration stimuli, 175
    - mal-de-debarquement syndrome, 175–176
    - vestibular training, 176
  - drug prophylaxis for, 180
  - pathogenesis, 174–175
  - physical maneuvers, 177
  - pragmatic therapy, 176–177
  - symptoms of, 173–174
  - types, 175

**O**

- Ocular vestibular-evoked potentials (oVEMPs), 42–43
- One-eye cover test, 15
- Ophthalmoscope, 24, 25
- Optokinetic drum, 24–26
- Oscillopsia, 159, 165
- Otolithic vertigo, 149

**P**

- Perilymph fistula
  - clinical aspects and course
    - computed tomography, 96, 97
    - incidence and prevalence, 96–97
    - provocation tests, 96
    - superior semicircular canal dehiscence, 97–100
    - traumatic, 148–149
    - Tullio phenomenon, 96
  - differential diagnosis, 96
  - pathophysiology, 98, 100–101
  - patient history, 94–95
  - pragmatic therapy, 101–102
- Peripheral vestibular forms of vertigo
  - benign paroxysmal positioning vertigo (*see* Benign paroxysmal positioning vertigo (BPPV))
  - bilateral vestibulopathy (*see* Bilateral vestibulopathy)
  - Menière's disease (*see* Menière's disease)
  - perilymph fistulas (*see* Perilymph fistula)
  - vestibular neuritis (*see* Vestibular neuritis)
  - vestibular paroxysmia (*see* Vestibular paroxysmia)
- Phobic postural vertigo
  - cardinal symptoms and features, 157–158
  - clinical aspects and course, 158–159
  - differential diagnosis, 162
  - organic syndromes, 163
  - pathophysiology, 159, 160
  - patient history, 157–158
  - posturographic analyses, 159–161
  - pragmatic therapy, 161–162
- Physical therapy, 49
  - bilateral vestibulopathy, 93
  - central compensation, 73
  - stance and gait, 93
- Politzer balloon, 96
- Posttraumatic BPPV, 146
- Posttraumatic psychogenic vertigo, 151
- Posturography, 44

**S**

- Seasickness, 175
- Semont maneuver, 60–61
- Somatoform vertigo
  - frequency
    - predictive value, 154
    - psychiatric comorbidity, 153–154
    - vestibular symptoms, 154
  - incidence, 154, 155
  - pathogenetic mechanisms, 156
  - pathogenetic model, 154
  - patient history, 155–156
  - phobic postural vertigo
    - cardinal symptoms and features, 157–158
    - clinical aspects and course, 158–159
    - differential diagnosis, 162
    - organic syndromes, 163
    - pathophysiology, 159, 160
    - patient history, 157–158
    - posturographic analyses, 159–161
    - pragmatic therapy, 161–162
  - treatment, 156–157
- Space sickness, 175
- Superior canal dehiscence syndrome,
  - cVEMP, 41

**T**

- Traumatic vertigo
  - central vestibular forms, 150
  - cervicogenic vertigo, 150
  - peripheral vestibular forms
    - alternobaric vertigo, 149
    - labyrinthine failure, 146–148
    - otolithic vertigo, 149
    - perilymph fistula, 148–149
    - posttraumatic BPPV, 146
  - posttraumatic psychogenic vertigo, 151
- Tullio phenomenon, 96

**U**

- Upbeat nystagmus (UBN)
  - etiologies, 124
  - magnetic resonance tomography, 124, 125
  - pathoanatomical findings, 124
  - therapy, 125

**V**

- Vehicle simulator sickness, 175
- Vergence test, 19–20

**Vertigo/dizziness**

- acrophobia
  - behavioral therapy, 181
  - clinical findings, 177–178
  - pathogenesis, 178
  - physiological visual height imbalance, 179, 180
  - prevalence, 177
  - subjective visual height intolerance, 178, 180
- central vestibular forms (*see* Central vestibular syndromes)
- cervicogenic vertigo, 172–173
- childhood (*see* Childhood vertigo)
- drug-induced vertigo, 171–172
- frequent forms, 5
- laboratory examinations
  - acoustic-evoked potentials, 44
  - computed tomography, 45–46
  - electronystagmography, 36, 38, 39
  - imaging, 44
  - magnetic resonance imaging, 46, 48
  - neuro-orthoptic and psychophysical examination, 37–38, 40–41
  - posturography and gait analysis, 44
  - pure-tone audiogram, 44
  - vestibular-evoked myogenic potentials (*see* Vestibular-evoked myogenic potentials (VEMPs))
  - video-oculography, 34–37
- motion sickness (*see* Motion sickness)
- neuro-ophthalmological and otological examinations, 33
  - convergence reaction, 19–20
  - Dix-Hallpike positioning maneuver, 29–30
  - eye, gaze positions, 16–19
  - finger-pointing test, 32
  - Frenzel's glasses and Politzer balloon, 23–24
  - head-impulse test, 25–27
  - head tilt measurement, 14
  - ophthalmoscope, 24, 25
  - optokinetic drum, 24–26
  - procedures, 13
  - saccades, 20–21
  - smooth pursuit eye movements, 21–22
  - stance and gait examination, 30–31
  - vergence test, 19–20
  - visual fixation suppression test, 27–29
- patient history, 5
  - brainstem/cerebellar symptoms, 10
  - duration of vertigo, 6–8

- headache, 10
- head movements, 10
- oscillopsia with head stationary, 9
- stance and gait control disturbances, 10–11
- trigger and exacerbation, 8
- type of vertigo, 6
- vestibular and audiological symptoms, 9
- peripheral vestibular forms of vertigo
  - benign paroxysmal positioning vertigo (*see* Benign paroxysmal positioning vertigo (BPPV))
  - bilateral vestibulopathy (*see* Bilateral vestibulopathy)
  - Menière's disease (*see* Menière's disease)
  - perilymph fistulas (*see* Perilymph fistula)
  - vestibular neuritis (*see* Vestibular neuritis)
  - vestibular paroxysmia (*see* Vestibular paroxysmia)
- peripheral vestibular vertigo, 2
- physical examination, 12
- physiological and pathological, 1, 2
- somatoform (*see* Somatoform vertigo)
- traumatic (*see* Traumatic vertigo)
- treatment
  - drug therapy, 46–49
  - physical therapy, 49
  - psychological/psychiatric and behavioral, 50
  - surgery, 50
- vestibulo-ocular reflex, 1, 3
- Vestibular-evoked myogenic potentials (VEMPs), 42
  - cervical vestibular-evoked myogenic potentials
    - bilateral vestibulopathy, 41
    - evaluation criteria, 40
    - Menière's disease, 42
    - superior canal dehiscence syndrome, 41
    - vestibular neuritis, 41
    - vestibular schwannoma, 41
    - vestibulo-colic reflex, 39
  - ocular vestibular-evoked potentials, 42–43
- Vestibular migraine
  - clinical aspects and course, 132–134
  - differential diagnosis, 136–138
  - pathophysiology, 134–135
  - patient history, 131–132
  - pragmatic therapy, 135–136
- Vestibular neuritis
  - causal therapy, 73, 75
  - central vestibular compensation, 73–75
  - cervical vestibular-evoked myogenic potentials, 41
  - clinical problems, 72
  - diagnosis, 69–70
  - differential diagnosis, 75–76
  - dynamic dysfunction, 71
  - etiology, 72
  - incidence, 70
  - pathophysiology, 71
  - patient history, 69
  - pragmatic therapy, 74–75
  - static symptoms, 71
  - symptomatic therapy, 73, 75
  - symptoms and clinical signs, 69, 70
- Vestibular paroxysmia
  - clinical aspects and course, 84–85
  - clinical problems, 89
  - diagnostic criteria, 86
  - differential diagnosis, 89
  - etiology, 87
  - pathophysiology, 86, 88
  - patient history, 84
  - pragmatic therapy
    - ineffective treatment, 89
    - medical treatment, 88
    - surgical treatment, 88–89
- Vestibular schwannoma, cVEMP, 81
- Vestibulo-colic reflex, cVEMP, 39
- Vestibulo-ocular reflex
  - horizontal (yaw) plane
    - clinical symptoms, 118–119
    - head-impulse test, 119
    - ischemic infarctions, 119–120
    - lesions, 118
    - neuronal pathways, 112, 113
  - planes of action, 1–2
  - sagittal (pitch) plane
    - brainstem–cerebellar network instability, 120
    - downbeat nystagmus syndrome, 123–125
    - lesion sites, 119–120
    - upbeat and downbeat nystagmus, 120
    - upbeat nystagmus, 124–125
  - vertical (roll) plane, 126–131
- Video-oculography (VOG)
  - caloric testing, 35–36
  - eye movement recording, 34–35
  - head-impulse test, 36, 37
- Visual fixation suppression test, 27–29