

Index

A

- Analgesics
 - adjuvant, 36
 - NSAID, 35
 - opioids, 35–36
- Anesthetic agents, 261–262
 - amide-linked group, 33
 - bupivacaine, 34
 - ester group, 33
 - lidocaine, 33–34
 - ropivacaine, 34
- Aneurysmal bone cysts (ABCs), 322–323
- Angina pectoris, 383–384
- Antibiotics, 261
- Aortogram, 327
- Arachnoiditis, 381–382
- Arteriovenous malformations (AVMs), 299
- Asymptomatic hemangiomas, 322
- Autonomic nerve blockade
 - celiac plexus blockade
 - anatomy, 235–237
 - contraindications, 239
 - indications, 236
 - injection technique, 236–238
 - common sympathetic blockades, 230
 - hypogastric plexus blockade, 243–245
 - impar ganglion blockade, 243–244, 246
 - lumbar sympathetic blockade
 - anatomy, 239, 240
 - indications, 239
 - injection technique, 239
 - radiofrequency block technique, 239–242
 - radiofrequency ablation, 230
 - stellate ganglion blockade
 - anatomy, 230, 231
 - complications and side effects, 232
 - indications, 230
 - injection technique, 230–232
 - radiofrequency-induced blockade, 232
 - sympathetic nerve blockade, 230
 - thoracic and splanchnic sympathetic blockades
 - anatomy, 233, 234
 - complications and side effects, 235
 - indications, 233

- injection technique, 233–234
- radiofrequency block technique, 235

B

- Balloon assisted vertebroplasty. *See* Kyphoplasty
- Bones, anatomy
 - parapedicular approach, 6
 - percutaneous vertebroplasty (PV)/kyphoplasty (KP), 1
 - scanning, 46–47
 - straight sagittal (anterior-to-posterior) orientation, 5
 - venous system, 6
 - vertebra size, 1

C

- Cavernous malformations, 313–314
- Celiac plexus blockade
 - anatomy, 235–237
 - contraindications, 239
 - indications, 236
 - injection technique, 236–238
- Cement
 - cortoss, 258–259
 - injection
 - cannula, 267
 - pain relief, 267–269
 - small syringes, 266–267
 - leak risk, 250
 - polymethylmethacrylate (PMMA)
 - antibiotics, 258, 261
 - bone reconstruction and augmentation, 249
 - disadvantages, 258–259
 - precaution, 258
 - selection and preparation, 257–258
- Cerebrospinal fluid leak, 292
- Cervix
 - epidural injections
 - lumbar, 168
 - procedure, 168–170
 - selective nerve root blocks, 170–171
 - facet joints, 210–213
- Chronic back pain, 378–379
- Cobb's syndrome, 317

Complex regional pain syndrome (CRPS), 379–381
 Computed tomography (CT)
 guidance, sacroplasty
 antibiotics, 363
 imaging, 366
 puncture sites, 363–364
 trans-SI approach, 364
 scanning, 45
 Constant flow rate infusion pump, 280
 Continuous epidural/intrathecal infusion trials, 285–286
 Conus medullaris AVM, 309–310
 Conventional radiographs, 44–45
 Corticosteroids
 betamethasone, 32–33
 methylprednisolone acetate, 32
 CPT coding device, 390
 Craig system, 97
 CRPS. *See* Complex regional pain syndrome (CRPS)

D

Dekompressor®, 151, 154
 Demographics, 158
 Diagnostic angiography, 323
 Discography
 cervical
 anatomy, 131
 anterior paratracheal approach, 134–135
 oblique approach, 135–138
 post-procedure care, 138
 procedure, 131–133
 surgical planning, 139
 complications, 108–109
 conscious sedation, 109–110
 disc morphology, 120, 121
 discogram report, 110
 indications, 108
 lumbar
 anatomy, 111–112
 functional anesthetic discography, 122–124
 manometry, 121–122
 positive predictive value (PPV), 110–111
 post-procedure care, 120–121
 procedure, 112
 surgical outcomes, 124
 surgical management, 58–61
 thoracic
 anatomy, 125
 post-procedure care, 130
 procedure, 125–130
 vertebrogenic pain, 141
 Dorsal root ganglion (DRG)-PRF procedure
 billing, 197
 complications, 191
 contraindication, 185
 efficacy, 195–196
 equipment and drugs, 185–186
 indications, 184–185
 laboratory, 186
 mechanism of action, 180–182

 patient history, 186
 patient positioning, 187–188
 physical examination, 186
 post treatment care, 191
 pre-operative medication, 186
 PRF lesioning history, 179–180
 radicular pain, 177–179
 surgical anatomy, 182–184
 technique, 188–193
 Dural arteriovenous malformations, 301–302

E

Electromyography (EMG), 47
 Endovascular treatment
 ABCs, 322–323
 aortogram, 327
 diagnostic angiography, 323
 microcatheter, 331
 neoplastic/metastatic lesions, 323
 neoplastic vascular lesions
 cavernous malformations, 313–314
 hemangioblastomas, 314
 neurophysiologic monitoring, 331
 spinal vascular diseases
 classification and clinical presentation, 300
 conus medullaris AVM, 309–313
 dural arteriovenous malformation, 301–303
 extradural AV shunts, 299–301
 extradural/intradural AVM, 307
 intradural (pial) arteriovenous fistula, 303–307
 intramedullary AVM, 307–308
 isolated spinal aneurysms, 313
 systemic syndromes
 Cobb's syndrome, 317
 Klippel–Trenaunay (KT) syndrome, 317
 neurofibromatosis type-1, 317–318
 Osler–Weber–Rendu Syndrome, 316–317
 Parkes–Weber (PW) syndrome, 317
 vertebral hemangiomas
 asymptomatic hemangiomas, 322
 characteristics, 318
 lesions causing cord compression, 319–321
 painful lesions, 321–322
 Epidural arteriovenous fistulas (AVS), 301
 Epidural injections
 cervical
 lumbar, 168
 procedure, 168–170
 selective nerve root blocks, 170–171
 demographics, 158
 fluoroscopic guidance, 161–162
 historical perspectives, 157–158
 patient selection, 158–159
 coagulation, 159–161
 contraindications, 161
 physical examination, 159
 steroid, 64–65
 techniques
 catheter, 164–165

- caudal, 163–165
 - intralaminar, 162–163
 - transforaminal, 166
 - Extradural AV shunts, 299–301
 - Extradural/intradural AVM, 307
 - Extrapedicular approach, 340
- F**
- Facet blocks, 61–62
 - Facet joint injections
 - anatomy
 - cervical facet joints, 210–213
 - lumbar facet joints, 209–210
 - sagittal and axial diagram, 208–209
 - thoracic facet joints, 213
 - clinical presentation, 207–208
 - contraindications, 214
 - diagnostic criteria, 213
 - pathophysiology, joint pain, 210
 - radiofrequency ablation, 216
 - techniques, 214–216
 - lateral cervical facet approach, 219
 - lumbar intra-articular, 216–217
 - lumbar periarticular, 217
 - medial branch block, 217–218
 - posterolateral cervical facet approach, 219
 - thoracic facet block, 219–220
 - Failed back spinal syndrome (FBSS), 378–379
 - Fine-needle aspiration techniques, 76–77
 - Fluoroscopic guidance, sacroplasty, 369
 - Fluoroscopy, 198–199
- G**
- Gate control theory, 376–378
 - Gentamicin, 125–126
 - Gout and headache, 376
- H**
- Headache. *See* Gout and headache
 - Hemangioblastomas, 314
 - Hereditary hemorrhagic telangiectasia. *See* Osler–Weber–Rendu syndrome
 - Hypogastric plexus blockade, 243–245
- I**
- Imaging equipments
 - biplane fluoroscopic equipment, 29–30
 - computed tomography (CT), 31–32
 - fixed-base C-arm fluoroscopic arrangement, 29–31
 - Impar ganglion blockade, 243–244, 246
 - Implanted drug delivery systems
 - advantages, 282
 - constant flow rate infusion pump, 280
 - continuous epidural/intrathecal infusion trials, 285–286
 - device-related complications
 - allergic reaction, 294
 - catheter-related problems, 294
 - pump-related problems, 292–293
 - SynchroMed® II, 293
 - infusate-related complications
 - bridge bolus calculation, 295
 - drug compounding errors, 295
 - programming errors, 294–295
 - technician errors, 295–296
 - surgical complications
 - bleeding, 290
 - cerebrospinal fluid leak, 292
 - infection, 291–292
 - neurological injury, 289–290
 - seroma formation, 290–291
 - Intraspinal drug delivery clinic, 282–283
 - Intrathecal drug delivery
 - device-related complications
 - allergic reaction, 294
 - catheter-related problems, 294
 - pump-related problems, 292–293
 - SynchroMed® II, 293
 - infusate-related complications
 - bridge bolus calculation, 295
 - drug compounding errors, 295
 - programming errors, 294–295
 - technician errors, 295–296
 - surgical complications
 - bleeding, 290
 - cerebrospinal fluid leak, 292
 - infection, 291–292
 - neurological injury, 289–290
 - seroma formation, 290–291
 - Intercostal neuralgia, 382–383
 - Interstitial cystitis/pelvic pain, 384
 - Intradural (pial) arteriovenous fistula (AVF), 303–307
 - Intramedullary AVM, 307–309
 - Intraspinal drug delivery clinic, 282–283
 - Intraspinal opioids, 284
 - Intrathecal drug delivery
 - device-related complications
 - allergic reaction, 294
 - catheter-related problems, 294
 - pump-related problems, 292–293
 - SynchroMed® II, 293
 - infusate-related complications
 - bridge bolus calculation, 295
 - drug compounding errors, 295
 - programming errors, 294–295
 - technician errors, 295–296
 - surgical complications
 - bleeding, 290
 - cerebrospinal fluid leak, 292
 - infection, 291–292
 - neurological injury, 289–290
 - seroma formation, 290–291
 - Isolated spinal aneurysms, 313
- K**
- Kirschner wire (K-wire), 342
 - Klippel–Trenaunay (KT) syndrome, 317
 - Kyphoplasty (KP), 67–68, 271–273
 - biomechanical investigations, 351
 - lateral radiograph demonstration, 338
 - patient selection

Kyphoplasty (KP) (*cont.*)

- relative contraindications, 339
- selection and exclusion criteria, 339

results

- complication rate, 350
- lateral radiograph, 349
- vs. PV, 350

technique

- anteroposterior image, 347
- cannula and stylet, 344
- extrapedicular approach, 340
- inflation end points, 346
- Kirschner wire (K-wire), 342
- lateral radiograph, 349
- materials required, 341
- posterolateral approach, 342
- technically demanding procedure, 349

VCFs, 337

L

Low back pain (LBP), 57

Lumbar discectomy. *See* Percutaneous lumbar discectomy

Lumbar discography

- anatomy, 111–112
- functional anesthetic discography, 122–124
- manometry, 121–122
- positive predictive value (PPV), 110–111
- post-procedure care, 120–121
- procedure
 - anterior, posterior tear and posterior bulge, 118, 119
 - curved tipped needle, 115, 116
 - disc morphology, 120–121
 - fluoroscope, 114
 - lateral and AP imaging, 118
 - superior articular process (SAP), 114, 115
- surgical outcomes, 124

Lumbar facet joints, 209–210

Lumbar sympathetic blockade

- anatomy, 239, 240
- indications, 239
- injection technique, 239
- radiofrequency block technique, 239–242

M

Magnetic resonance imaging (MRI), 45–46, 251–253, 256–257

Microcatheter, 331

Minimally invasive intradiscal therapy, 65–66

Motor evoked potentials (MEP), 331

Myelography, 45

N

Neoplastic/metastatic lesions, 323

Neoplastic vascular lesions

- cavernous malformations, 313–314
- hemangioblastomas, 314

Nerve conduction studies (NCVs), 48

Neurofibromatosis type-1, 317–318

Neurolytic (cytotoxic) agents, 37

Neuromodulation/neurodestructive procedures, 54–55

Neuropathic pain treatment algorithm, 41–42

Neurostimulation history, 375–376

Nonsteroidal anti-inflammatory (NSAID), 35

Nuclear medicine (NM), 252–253, 255

O

Occipital neuralgia, 383

Opioids, 35–36

Osler–Weber–Rendu syndrome, 316–317

P

Pain relief, 273–274

Parkes–Weber (PW) syndrome, 317

Patient evaluation

- ablative techniques, 40
- diagnostic nerve blocks
 - central nerve blocks, 50
 - false positive results, 48–49
 - peripheral nerve blocks, 49–50
- electrodiagnostics
 - electromyography (EMG), 47
 - nerve conduction studies (NCVs), 48
- imaging studies
 - bone scanning, 46–47
 - computed tomography scanning (CT), 45
 - conventional radiographs, 44–45
 - magnetic resonance imaging (MRI), 45–46
 - myelography, 45
 - thermography, 47
 - ultrasound, 46
- laboratory tests, 48
- medical history, 42–43
- physical examination, 43–44
- procedure selection criteria
 - algorithmic approach, 52–53
 - neuromodulation/neurodestructive procedures, 54–55
 - pain generators, 51–52
 - radiofrequency lesioning techniques, 54–55
 - psychological evaluation, 50–51

Percutaneous lumbar discectomy

- anatomy, 147–148
- complications and consent, 152
- historical perspective, 149–150
- intervertebral discs, spinal pain, 149
- needle placement
 - bulged disc, 150, 151
 - Dekompressor®, 151, 154
 - lateral view, 150, 153
 - P–P proper needle placement, 150, 152

Percutaneous sacroplasty (PS), 357

Percutaneous spine biopsy

- complications, 76–78
- destructive/space-occupying lesions, 76

- equipment requirements
 - biopsy needle systems, 80–81
 - choice, 79
 - fluoroscopy-guided biopsies, 79–80
- fine-needle aspiration techniques, 76–77
- hemorrhage/hematoma formation, 76, 77
- indications, 76
- patient preparation
 - axial CT image, 78–79
 - laboratory parameters, 78
- postoperative care, 101
- soft tissue
 - dysphagia, 98, 99
 - presacral abscess, 101–105
 - sacroiliac osteomyelitis and abscess, 100, 101
- techniques
 - approaches, 81–82
 - Craig system, 97
 - disc and vertebral endplate, 84
 - hematoma formation, 76, 77
 - hypervascular lesions, 97
 - lesion location and lesion size approach, 81–82
 - microbiological analysis, 94
 - needle excursions, 91, 93, 94
 - parapedicular approach, 83, 84
 - posterior approach, 83
 - procedure-related complication, 94, 95
 - routes, 83
 - sclerotic/osteoblastic lesions, 94, 96
 - skin entry site, 85, 86
 - suprahyoid and infrahyoid compartments, 88, 89
 - transpedicular approach, 88, 92
 - vertebral body lesion and posterior element lesion, 82–83
- Percutaneous vertebroplasty (PV)
 - anesthesia, 261–262
 - antibiotics, 261
 - cement injection, 266–269
 - cement selection and preparation, 257–259
 - complications, 272–273
 - image guidance, 259–261
 - indications, 250
 - informed consent, 259
 - laboratory evaluations, 261
 - needle introduction and placement, 262–266
 - pain relief, 273–274
 - patient selection and workup, 250–257
 - postoperative care, 268, 270, 271
 - results, 270–272
 - venography, 266
 - vertebral compression fracture (VCF), 249–250
- Peripheral nerve PRF procedure
 - complications and efficacy, 201–202
 - peripheral nerve damage, 197, 198
 - post surgical pain treatment, surface landmarks, 199, 201
 - suprascapular nerve treatment, 198–200
 - technique, 197–198
- Peripheral nerve stimulator, 383
- Peripheral stimulation, technical factors, 384–385
- Peripheral vascular disease, 383–384
- Personal therapy manager (PTM), 282
- Phantom limb pain, 382
- Pharmacological agents
 - analgesics
 - adjuvant, 36
 - NSAID, 35
 - opioids, 35–36
 - anesthetic agents
 - amide-linked group, 33
 - bupivacaine, 34
 - ester group, 33
 - lidocaine, 33–34
 - ropivacaine, 34
 - antibiotics, 34–35
 - corticosteroids
 - betamethasone, 32–33
 - methylprednisolone acetate, 32
 - neurolytic (cytotoxic) agents, 37
 - radiographic contrast agents, 36–37
- PHN. *See* Post-herpetic neuralgia
- Polymethylmethacrylate (PMMA)
 - antibiotics, 258, 261
 - bone reconstruction and augmentation, 249
 - disadvantages, 258–259
 - precaution, 258
 - selection and preparation, 257–258
- Posterolateral approach, 342
- Post-herpetic neuralgia (PHN), 381
- Post surgical pain treatment, 199, 201
- Pulsed radiofrequency (PRF)
 - application, 175
 - DRG procedure
 - billing, 197
 - complications, 191
 - contraindication, 185
 - efficacy, 195–196
 - equipment and drugs, 185–186
 - indications, 184–185
 - laboratory, 186
 - mechanism of action, 180–182
 - patient history, 186
 - patient positioning, 187–188
 - physical examination, 186
 - post treatment care, 191
 - pre-operative medication, 186
 - PRF lesioning history, 179–180
 - radicular pain, 177–179
 - surgical anatomy, 182–184
 - technique, 188–193
 - pain medicine, 176–177
 - peripheral nerve procedure
 - complications and efficacy, 201–202
 - peripheral nerve damage, 197, 198
 - post surgical pain treatment, surface landmarks, 199, 201
 - suprascapular nerve treatment, 198–200
 - technique, 197–198
 - radiofrequency current effect, 175–176

- R**
- Radiofrequency lesioning techniques, 54–55
 - Radiographic contrast agents, 36–37
 - Reflex sympathetic dystrophy (RSD), 379
 - Retrocural approach. *See* Splanchnic nerve denervation
 - Retroneural approach. *See* Dorsal root ganglion (DRG)-PRF procedure
 - RSD. *See* Reflex sympathetic dystrophy (RSD)
- S**
- Sacral fractures
 - anatomy, 361
 - diagnosis, 358–361
 - incidence, 357–358
 - morbidity and mortality, 358
 - patient selection, 361–362
 - Sacroiliac joint (SI) injections, 62–63
 - anatomy, 220–223
 - clinical presentation, 222
 - diagnosis, 222–226
 - innervation, 221–222
 - techniques, 226
 - Sacroiliac osteomyelitis, 100, 101
 - Sacroplasty
 - complete clinical picture, 369–371
 - complications, 369
 - conventional treatment, 355–356
 - CT guidance
 - antibiotics, 363
 - CT imaging, 366
 - puncture sites, 363–364
 - trans-SI approach, 364
 - diagnosis, 355
 - fluoroscopic guidance, 369
 - outcomes, 369
 - percutaneous sacroplasty (PS), 357
 - sacral fractures
 - anatomy, 361
 - diagnosis, 358–361
 - incidence, 357–358
 - morbidity and mortality, 358
 - patient selection, 361–362
 - typical patient presentation, 355
 - Selective nerve root blockade, 63–64
 - SI injections. *See* Sacroiliac joint (SI) injections
 - Single intrathecal bolus dosing, 285
 - Somatosensory evoked potentials (SSEP), 331
 - Spinal cord stimulation
 - angina pectoris and peripheral vascular disease, 383–384
 - arachnoiditis, 381–382
 - chronic back pain, 378–379
 - complications, 386–387
 - CPT coding device, 390
 - CRPS, 379–381
 - device manufacturers, 391–394
 - dorsal roots, 378
 - FBSS, 378–379
 - gate control theory and implantable stimulators, 376–378
 - gout and headaches, painful condition, 376
 - intercostal neuralgia, 382–383
 - interstitial cystitis/pelvic pain, 384
 - neurostimulation history, 375–376
 - occipital neuralgia, 383
 - peripheral stimulation, technical factors, 384–385
 - phantom limb pain, 382
 - PHN, 381
 - technical factors, 385–386
 - techniques and implantation pearls, 387–390
 - Spinal vascular diseases
 - classification, 299
 - conus medullaris AVM, 309–313
 - dural arteriovenous malformation, 301–303
 - extradural AV shunts, 301
 - extradural/intradural AVM, 307
 - intradural (pial) arteriovenous fistula (AVF), 303–307
 - intradural AVM, 307–309
 - isolated spinal aneurysms, 313
 - Spine anatomy
 - anatomical spaces, 14–15
 - bones
 - parapedicular approach, 6
 - percutaneous vertebroplasty (PV)/kyphoplasty (KP), 1
 - straight sagittal (anterior-to-posterior) orientation, 5
 - venous system, 6
 - vertebra size, 1
 - intervertebral discs and joints, 8
 - macrocirculation
 - embryological development, 15
 - radicular arteries, 18
 - radiculomedullary arteries, 18
 - radiculopial arteries, 18
 - segmental arteries, 17
 - microcirculation
 - centrifugal system, 19
 - centripetal system, 21
 - nerves
 - lumbar foramina, 12
 - neural foramina, 7
 - parasympathetic nerves and sympathetic nerves, 13
 - somatic arterial supply
 - angiography, 22
 - intervertebral disc, 8–11
 - venous
 - epidural veins, 26
 - intrinsic venous system, 23
 - lumbar level, 19
 - multiple longitudinal venous channels, 25
 - thoracic level, 19
 - Splanchnic nerve denervation, 235
 - Splanchnic sympathetic blockade. *See* Thoracic and splanchnic sympathetic blockade
 - Stellate ganglion blockade
 - anatomy, 230, 231
 - complications and side effects, 232
 - indications, 230
 - injection technique, 230–232
 - radiofrequency-induced blockade, 232

- Suprascapular nerve treatment, 198–200
- Surgical management
- discography, 58–61
 - epidural steroid injections, 64–65
 - facet blockade, 61–62
 - low back pain (LBP), 75
 - minimally invasive intradiscal therapy, 65–66
 - sacroiliac joint (SI) injections, 62–63
 - selective nerve root blockade, 63–64
 - vertebroplasty and kyphoplasty, 66–68
- Sympathetic nerve blockade, 230
- Systemic syndromes
- Cobb's syndrome, 317
 - Klippel–Trenaunay (KT), 317
 - neurofibromatosis type-1, 317–318
 - Osler–Weber–Rendu syndrome, 316–317
 - Parkes–Weber (PW) syndrome, 317
- T**
- Thermography, 47
- Thoracic and splanchnic sympathetic blockade
- anatomy, 233, 234
 - complications and side effects, 235
 - indications, 233
 - injection technique, 233–234
 - radiofrequency block technique, 235
- Thoracic discography
- anatomy, 125
 - post-procedure care, 130
 - procedure
 - disc annulus, 126, 128
 - fluoroscopic views, 127, 129–130
 - gentamicin, 125–126
 - lamina, 126
- Thoracic facet joints, 213
- Transcostovertebral approach, 79, 83, 85, 88
- Transpedicular approach, 88, 92, 263–265
- Treatment of spine related pain syndromes.
See Epidural injections
- U**
- Ultrasound, 46
- V**
- Venography, 266
- Ventral intradural AVF. *See* Intradural (pial) arteriovenous fistula (AVF)
- Vertebral compression fracture (VCF), 249–250, 337
- Vertebral hemangiomas
- asymptomatic hemangiomas, 322
 - characteristics, 318
 - lesions causing cord compression, 319–321
 - painful lesions, 321–322
- Vertebroplasty, 66–67
- W**
- Waddell's signs, 44
- Z**
- Zygapophyseal joints, 49, 61