

Index

- Acantholinqua* 1
- adaptive strategies 31, 32
- adult trout
 - age and size 186–8
 - preferred habitat 56–63
- age
 - adults 186–8
 - determination 154–6
 - feeding rhythms and 98
 - at maturity, brown trout 23, 25
 - sea trout 156–9
 - smolts 184–5
- agonistic behaviour in fry 119–23
 - bite 122
 - charge 120
 - chase 122
 - flight 122–4
 - frontal and lateral threat displays 122
 - submission posture 122
- aluminium 40
- alevins, preferred habitat of 62
- anadromization in brown trout 198
- angling 6
- Anguilla anguilla* (eel) 16
 - competition with 67
- aquaculture 7
- area of distribution 41
- bank, stream *see* stream bed and banks
- bank effect 39
- benthic population, effects on 105
- biogenic capacity of environment 31–2, 278, 280
- biological cycle of brown trout 7–9, 28–30
- biological reference flow 72
- biotypological system 50–1
- Brachymystax* 1
- breeding areas 54–6
- calcium 40
- calcium index 23, 31
- caryology 206
- Chironomus* 128
- circadian rhythms, territoriality dynamics and 135–6
- commercial fishing 7
- competition *see* interspecific competition;
intraspecific competition
- conservation of genetic diversity 216–17
- Continuous Under Threshold Habitat Duration 75
- Coregonus lavaretus* (whitefish) 147
- Cottus gobio* (bullhead) 16
 - competition with 67–8
- dams, effect of 71–2
- displacement, downstream, of fry 128–33
 - in artificial streams 129–33
 - in natural environment 128–9
- distribution of brown trout
 - current 4–6
 - original 2–4
- drainage order 47
- E.V.H.A. 75
- ecological sector 41
- electrophoretic analysis 206–7, 211–12, 220
- energy budget 100–4, 105
 - accumulation 100–2
 - calculation 103
 - losses through excretion 102
 - metabolic losses 102–3
- enzyme polymorphism analysis 206–7
- Esox lucius* (pike) 16, 147
- exploitation *see* fishing
- fecundity
 - brown trout 26
 - sea trout 164–7, 193, 199
- feeding behaviour
 - brown 94–5
 - sea trout 161–3
- feeding rhythms 96–8

284 Index

- fishing 195–6
 - commercial 7
 - effects on genetic diversity 217–18
 - sort 6
- flow (current) 38
- fluvial hydrosystem 46
- food consumption rate 98–100
- food consumption, study methods 94
- food sources 92
- food stock estimation 92–3
 - spatial variability 93
 - temporal variability 93–4
- fry
 - agonistic behaviour 119–23
 - early downstream displacement 128–33
 - early social and territorial behaviour 117–19
 - life under gravel and learning to swim 119
 - site attachment 119–24
- Gastrocnemius aculeatus* (stickleback) 115
- gene flow between domestic and natural stocks 213–214
- genetic diversity
 - comparison with other salmonid species 213
 - conservation of 216–17
 - ecological differentiation 211–12
 - effects of fishing 217–18
 - effects of restocking 213–16, 218–20
 - genetic populations 205–6
 - geographic differentiation 207–11
 - significance of differentiation 212–13
- genetic management 217–20
 - long-term 216–17
 - short-term 217–20
- genetic variability, description of 206–7
- geographical isolation 207–11, 212
- Gobio gobio* (gudgeon) 68
- granulometry 39
- growth
 - brown trout 20–3, 24
 - habitat and 53–4
 - juveniles 153
 - sea trout 159–60, 187–8
 - spawners 159–61
- habitat of brown trout
 - characteristic feature 38–41
 - definition 37
 - growth and 53–4
 - preferred 56–63, 77
 - spatial integration of results 77
 - study objectives 77–8
 - temporal variations 77
- Habitat Quality Index (HQI) 73
- Habitat Suitability Index (HIS) 75
- HABSCORE 76
- 'half-blood' strategy 219–20
- heavy metals 40
- home range 46
- homing 167–8
- Hucho* 1
- hydropeaking 75
- interspecific competition 64–8
 - with Atlantic salmon (*Salmo salar*) 65–6
 - with bullheads 67–8
 - with eel 67
 - with minnow 67
 - with other salmonids 65–7
 - with rainbow trout, brook charr and coho salmon 66–7
- intraspecific competition 63–4
- juveniles
 - mortality 154
 - populations 150–3
 - preferred habitats 56–62
 - see also alevins; fry
- Lepomis cynellus* (green sunfish) 115
- LESLIE matrix 76
- light 39
 - feeding rhythms and 97
- longitudinal structure of trout zone 52–4
- longitudinal zonation 47–51
- Lotta lotta* (burbot) 147
- macrobenthos productivity 92–3
- macrohabitat 41, 43, 44–6
- macrophytes 40–1
- management
 - a priori* 279–80
 - ecological study and 278–9
 - natural populations in France (1660–1986) 227–59
 - methodology 277–8
 - temporal variation 72
- mean age at smoltification (M.S.A.) 185
- mean sea age at first reproduction (M.R.A.) 188
- microhabitat 43
- migration
 - energy costs 105
 - migratory rhythms 180–3
 - phases of 27
 - reproductive, brown trout 27–8
 - in the sea 183–4
 - sea trout 180–4
 - up stream 163–4
- mortality of juveniles 154
- Nemacheilus barbatulus* (stone loach) 16, 68
- nitrates 40
- nitrite 40
- Oncorhynchus* 1, 2
 - homing in 167
- Oncorhynchus clarkii* 73
- Oncorhynchus gairdneri* 73

- Oncorhynchus gorbuscha* 137
Oncorhynchus keta 137
Oncorhynchus kisutch (coho salmon),
 competition with 66–7
Oncorhynchus mykiss (rainbow trout),
 competition with 66–7
Oncorhynchus nerka 137
 overpopulation, effect of, on territoriality 124
 oxygen, dissolved 40
- panmixia 206
 partial territories 134
Perca fluviatilis (perch) 147
 pH 40
 PHASBIM method 72, 73, 76
 phosphates 40
Phoxinus phoxinus (minnow), competition with 67
 phylogeny, brown trout 1–2
 population dynamics in brown trout 28–30
 population estimation 72–6
 prediction models for macrohabitats 73–5
 prediction models for trout zone 73
 populations
 brown trout 16–19
 genetic 205–6
 genetic management of 216–20
 sea trout 199
Potamopyrgus jenkinsi 95, 96
 prediction models
 for macrohabitats 73–5
 for trout zone 73
 prey
 chase after 94–5
 effect on territoriality 124–8
 prey capture rate 95–6
 intensity of consumption 95–6
 prey selection 95
 prey size 95
- redds
 characterization 56
 location with macrohabitats 55–6
 reproduction
 in sea trout 163–8
 rate, in sea trout 188–93
 reproductive migration in brown trout 27–8
 restocking 169–70, 280
 effects on genetic diversity 218–20
 rheotaxis 117, 119
 river continuum concept 91
 rotating territories 134
Rutilus rutilus (roach) 147
- Salmo (Trutta) carpio* 1
Salmo ferox 211
Salmo gairdneri (rainbow trout; steelhead) 1
Salmo (Trutta) marmoratus 1
Salmo salar (Atlantic salmon) 1, 16, 116
 competition with 65–6
 homing in 167
Salmo trutta macrostigma 211
Salmothymus 1
Salvelinus 1, 2
Salvelinus alpinus (Arctic charr) 147
Salvelinus fontinalis (brook charr) 4, 73, 122
 competition with 66–7
 scale growth 155, 278
 seasonal rhythms, territoriality dynamics and 135–6
 segment 41
 sequence 1
 sex ratio
 brown trout 23
 sea trout 156–9, 191–2, 193
 sexual maturation
 brown trout 23–6
 sea trout 156–9
 size
 adults 186–8
 fecundity and 164–6
 at maturity in brown trout 23, 26
 sea trout 180, 198
 smolts 184–5
 slope width index 23, 31
 spawning
 brown trout 27–8
 multiple 164–7
 sea trout 164
 spawning sites
 brown trout 28
 sea trout 163–4
 spawning zones 54–5, 57–8
 sport fishing 6
 static swimming 117
 stocking capacity 280
 stream bed and banks
 cleaning 70
 cutting macrophyte vegetation 70–1
 gravel extraction 71
 modifications of flow and dams 71–2
 physical changes 70–1
 restoration 70
 straightening and widening 70
 water abstraction 71
 water discharge 71–2
 stream bed morphology 39
 stream flow 71–2
 stream order 48
 submission posture 122
 substrate particle size (granulometry) 39
 survival rates
 brown trout 28, 29
 sea trout 193–6
 suspended matter 40
 sympatry 211–12

286 **Index**

- temperature 39–40
 - feeding and 97
- temporal variations 68–72
 - daily 68
 - habitat management 72
 - human activity 69–72
 - natural 68–9
 - seasonal 69
 - physical changes to bed and banks 70–1
 - modifications of flow and dams 71–2
- territorial behaviour 115–17
 - in fry 117–24
 - during ontogeny 133–6
- territory
 - attachment in fry 123–4
 - density and 124–6
- toxins, feeding rhythms and 98
- trout zone 41
 - longitudinal structure 52–4
 - longitudinal zonation 47–51
- vegetation, bank-side 40–1
- Von Bertalanffy model 22
- water abstraction 71
- water discharge 71–2
- water quality 30, 50
- Weighted Suitable Area 75
- zone of influence 122