

# Index

## A

- Adenoids, prominent, 31
- Anatomy, developmental, 1–11
- Aneurysmal bone cysts, 125–127
- Angulation, of C2 on C3, 20–21
- Anomalies, 39–56
  - of the dens, range of, 58
- Apophyseal joints
  - abnormalities of, in flexion-induced injuries, 83–85, 91–92
  - normal, in the older child, 26–28
  - obliteration of, in rheumatoid arthritis, 123–124
- Arachnoid cysts, 128
- Arch defects
  - anterior C1, 53
  - posterior
    - C1, 46–47
    - C2, 47–51
    - lower vertebrae, 51–53
  - See also* Neural arches
- Arnold-Chiari I malformations, 128
- Astrocytomas, spinal cord, 128
- Atlantoaxial instability, in rheumatoid arthritis, 123–125
- Atlanto-occipital dissociation, 115–117
- Atlanto-occipital instability, 132–133
- Atlas (C1), embryonic development of, 1
- Avulsion fracture
  - differentiating ring epiphysis from, 30–32

of the spinous tips, 87, 89

*See also* Teardrop fracture

## Axis (C2)

- embryonic development of, 1
- synchrondoses of, 6–7

## B

- Basilar invagination, 44, 48
- Battered child syndrome,
  - hangman's fractures in, 103
- Bone cysts, aneurysmal, 125–127
- Brachial plexus injuries, 117–119
- Butterfly vertebra, 39–40

## C

- Calcific diskitis, 124–126
- Central veins, misinterpreting as fractures, 36–37
- Cervical line, normal posterior, 22
- Cervical spine
  - lower
    - injuries to, 79–95
    - vertebral fusion at, 41
  - normal motion of, causing pseudoabnormalities, 15–16
  - upper, injuries to, 95–118
- Chromosomes
  - trisomy 13–15, coronally cleft vertebrae associated with, 39–40
  - trisomy 21, atlanto-occipital instability associated with, 133

- Chondrofication centers, in development of cervical vertebrae C3–C7, 2–4
- Clay shoveler's fracture, 87–89
- Cleft vertebrae, anomalies associated with, 39
- Collagen vascular diseases, cervical spine instability in, 123–124
- Computed tomography  
axial, for Jefferson fracture evaluation, 113–115  
for rotation force injury evaluation, 94–95  
for subluxation or dislocation evaluation, 110–112  
in trauma to the cervical spine, 77
- Coronal hemivertebra, 39–40
- Cysts  
arachnoid, 128  
bone, aneurysmal, 125–127
- D**
- Dens  
abnormal development of, associated with C1–C2 fusion, 42–43  
absence of, 69  
anomalies and normal variation of, 57–73  
bifid, 71–72  
as the body of C1, 70  
embryonic development of, 1  
fracture of, 115–116  
in anterior force injury, 99–100  
differentiation from os odontoideum, 64–66  
in extension force injury, 107  
in hyperflexion injuries, 96–98  
pseudo-os odontoideum with, 68  
ossification centers of, 3  
tilting of, 70–71  
in extension force injury, 106  
from hyperextension forces, 103
- Disk, disruption of  
in extension force injuries, 104–105  
in rotation force injuries, 92–94
- Dislocation  
anterior, of C2 on C3, 21–23  
C1–C2, due to rotational force, 110  
of C2 on C3, differentiating from hangman's fracture, 99–102  
posterior, of vertebral bodies, 19–20
- E**
- Embryonic development, of the spine, 2  
lack of dens in, 69
- En face pedicle sign, in rotational injury, 92–94
- Ewing's sarcoma, 127–128
- F**
- "Fat" C2 sign, 113
- Flexion curve, apex of, change in location with normal growth, 16
- Forces, in cervical spine injury, 75–76. *See also* Flexion forces
- Fractures  
central veins mistaken for, 36–37  
compression (axial load) injuries in the lower cervical spine, 92–95  
in the upper cervical spine, 112–115  
computed tomography for differentiating from wedging, 24–26  
extension injuries in the lower cervical spine, 87–91  
in the upper cervical spine, 97–107  
flexion injuries in the lower cervical spine, 79–87, 88, 89  
in the upper cervical spine, 95–97  
forces in, 75–76  
in the lower cervical spine, 79–95  
occult, of the cervical spine and base of the skull, 77  
pars, in hypoextension injury, 89–90  
posterior C1, in hyperextension, 101

- rotation injuries
  - in the lower cervical spine, 91–92, 93
  - in the upper cervical spine, 108–112
  - in the upper cervical spine, 95–115
- Fusion-segmentation anomalies, 39
  - at the C2–C3 level, 41
- H**
- Hangman's fracture
  - anterior dislocation of C2 on C3 in, 21
  - with C2–C3 disk disruption, 104–105
  - differentiating from posterior C2 arch defects, 47–51, 99–102
  - in infants, 107–108
- Hemangiomas, 125–127
- High anterior arch of C1, 16–17
- Histiocytosis X, 130–132
- Hypertrophy, of intervertebral disks, with vertebral fusion, 42
- Hypoplasia, dens, associated with posterior C1 arch defects, 46–47, 51
- I**
- Infection leading to cervical spine instability, 123–125
- Inflammation leading to cervical spine instability, 123–125
- Iniencephaly, 42–43, 46
- Injuries, to the cervical spine, differences between infants and adults, 16
- Instability of the cervical spine, plain film signs of, 78–79
- Interspinous distance, C1–C2, exaggerated, 17
- Intervertebral disks, hypertrophied, vertebral fusion with, 42
- J**
- Jefferson fracture, 112–115
  - differentiating from synchondrosis, 4
  - versus lateral mass pseudooffset, 34
- K**
- Klippel-Feil syndrome, fusion anomalies in, 42, 45
- Kyphosis, at the C3–C4 level, associated with a coronal hemivertebra, 40
- L**
- Langerhans cell histiocytosis, 130–132
- Lateral flexion injuries, 91
- Lateral mass, C1, bilateral and unilateral offset of, 34–36
- Lateral view, for evaluating cervical spinal injuries, 76–77
- Leukemia, bony involvement of the cervical spine in, 128
- Ligaments, underdevelopment of, in association with a missing dens, 69
- Locked facette sign, in flexion-induced injuries, 84, 87, 91–92
- Lymphoma, bony involvement of the cervical spine in, 128
- M**
- Magnetic resonance imaging
  - in acute trauma, 78
  - for differentiating hemangiomas from aneurysmal bone cysts, 127
  - for evaluation
    - of injury in SCIWORA syndrome, 116–117
    - of neurofibromatosis, 129
    - of soft tissues, 29, 91–95
    - of vertebral fusion, 42
- Malignant tumors, primary, of the cervical spine, 127–128
- Motion artifact, in computed tomography, 118–119
- Motion of the cervical spine, causing pseudoabnormalities, 15–16
- N**
- Neoplastic diseases, 125–128
- Neural arches
  - development of, 2–4
  - dysplastic, 55

- Neural arches *cont.*  
 fractures of, 88–99  
 fusion of, C1 and C2, 43–44  
 normal variations of, 13–15
- Neurofibromatosis  
 plexiform, 130–131  
 type I, 129–131
- Neurological deficit, in SCIWORA syndrome, 116–117
- Normal synchondroses, 4–10
- Normal variations, 13–38
- O**
- Occipital condyles, articulation with the lateral masses of C1, 1
- Occipitalization, 13, 43–47
- Open facette sign, 91–94
- Open-mouth odontoid view for evaluating cervical spinal injuries, 76–77
- Os odontoideum  
 acquired, 68, 97  
 configurations of, 64–66  
 development of, with a hypoplastic dens, 59
- Ossicles  
 abnormal  
 association with a hypoplastic dens, 67  
 in the area of the os terminale, 62
- Ossification centers, of the dens, 3–4
- Osteogenesis imperfecta, basilar invagination associated with, 46
- Osteogenic sarcoma, 129
- Osteoid osteomas, 127
- Osteomyelitis-diskitis of the cervical spine, 123–125
- Os terminale  
 complex with the os odontoideum, 57–69  
 embryonic derivation of, 1, 57  
 normal, 58–59  
 variable appearance of, 60–61
- P**
- Pars, fractures of, in extension force induced injuries, 88–89
- Perched facette sign, in flexion-induced injuries, 84, 86, 91–92
- Platybasia, of the skull, 44
- Posticus ponticus, 36–37
- Predental distance, in infants, 17–19
- Pseudodislocation, anterior, at the cervical spine, 23–24
- Pseudo-offsetting, lateral masses of C1 in infancy, 34–36
- R**
- Rheumatoid arthritis, cervical spine instability in, 123–124
- Rhizomelic punctate epiphyseal dysplasia, coronally cleft vertebrae associated with, 39
- Ring epiphysis  
 differentiating from avulsion fractures, 30–32  
 normal, 32
- S**
- SCIWORA syndrome, 116–117, 118
- Segmentation anomalies, 40
- Soft tissues  
 injury to, magnetic resonance imaging for delineating, 78  
 prevertebral  
 evaluating, 26–30  
 pathologic thickening of, 31
- Spina bifida, 7, 11
- Spinal canal, cervical, pseudowidening of, 31–32
- Spinal cord  
 compression of, from instability at the C1–C2 level, 62  
 injury to  
 in hyperextension injuries, 89–91  
 magnetic resonance imaging for delineating, 78  
 SCIWORA syndrome, 118  
 tumors of, 128–129
- Spinal stenosis, 134
- Spinous processes, normal variations of, 13–14
- Spinous tips, avulsion fractures of, 87, 89
- Sprengel's deformity, 42, 45

Still's disease, cervical spine  
instability in, 123–124

Synchondroses, 4–11

Syringomyelia, 128

## T

Teardrop fracture

in anterior force fracture of the  
lower cervical spine, 81–82

in anterior force fracture of the  
upper cervical spine, 96–  
97

in extension force injury to the  
lower cervical spine, 88–91

in flexion force injury at C2–C3,  
84

Transverse processes, 30–32

Trauma, 75–122

Trisomy

13–15, coronally cleft vertebrae  
in, 39–40

21, atlanto-occipital instability  
associated with, 133

## U

Upper cervical spine, injuries to,  
95–118

## V

Vertebrae, C3–C7, synchondroses  
of, 7, 10

Vertebral bodies

development of, ossification

center development, 4

dysplastic, in neurofibromatosis,  
129

fusion anomalies of, 39–46

normal variations of, 13–15

## W

Wedging, of C3 and C4, 23–28

Wry neck, 109–111