

Index

A

- AA. *See* Ascorbic acid (AA)
- ABA. *See* Abscisic acid (ABA)
- Abortion
- agent, 391
 - foetus/cause, 395
 - prostaglandin, 395
- Abscisic acid (ABA), 242
- ACE. *See* Angiotensin I-converting enzyme (ACE)
- ADP. *See* Allele discovery platform (ADP)
- AFLP. *See* Amplified fragment length polymorphism (AFLP)
- African Caricaceae
- neotropical clade, 85, 87
 - species, 83, 87
 - statistical area reconstruction, 88
 - strict clock model, 87
- Alanine aminotransferase (ALT), 402
- Allele discovery platform (ADP)
- allergen reduction, 415
 - antioxidant levels, 416
 - breeding system, 409
 - clonal propagation system, 410
 - and CODDLE, 412
 - cyanide reduction, 415
 - detection, mutations, 412
 - disease and pest resistance, 417
 - and DNA, 411
 - drought resistance, 416
 - Eco-ADP, 410
 - efficient crop nutrition, 415
 - family Caricaceae, 409
 - flowering and fruiting, 409
 - flower types, 413–414
 - fruit crops, 410
 - fruit quality, 414
 - fruits consumed and nutritional profile, 409
 - hand pollination, 409–410
 - ionizing radiation, 411
 - LI-COR-based systems, 413
 - life cycle, crop/early-bearing crop, 416
 - medicinal use, 416
 - mutant population using EMS, 411
 - nematode resistance, 415
 - non-transgenic, 418
 - papain production, 415
 - papaya genomics, 413
 - PCR products, 411, 412
 - phenotypic database, 417
 - and PRSV, 414–415
 - reverse genetic functional genomics tool, 410–411
 - root architecture, 416
 - and SNP, 410
 - source and sink, 417
 - and TILLING, 410
 - and VAD, 409
 - and WHO, 409
- ALT. *See* Alanine aminotransferase (ALT)
- Amplified fragment length polymorphism (AFLP)
- cluster analysis, 39
 - cpDNA intergenic spacer region, 44
 - gene content, 148
 - genetic variation, 41
 - genome sequence, 147
 - high-density genetic map, 147
 - morphological characteristics, 362
 - papaya genetic map, 150, 360
 - PRSV-P coat protein marker, 359
 - RFLP, 358

- Amplified fragment length polymorphism (AFLP) (*cont.*)
 segregation distortion, 148
 sex co-segregating markers, 149
- Angiotensin I-converting enzyme (ACE), 402
- Anthelmintic activity
 alkaloid carpaine and benzyl thiourea, 392
Ascaridia galli, 392–393
 BITC, 392
 dried papaya seeds, 392
Heligmosomoides polygyrus, 393
in vitro and infected animals, 392
 proteolytic enzymes, 392
- Anti-amoebic activity, 401
- Anticancer and chemoprotection
 BITC, 400
 cell death, 400
 CYP2E1, 400
 enzymes act, 400
GSTM1, 401
 isothiocyanates, 400–401
 monocytes and memory T lymphocytes, 400
 nitrosamines, 400
- Antifertility properties
 females
 abortifacient, 394
 administration, purified papain, 395
 animal bioassays, 395
 aqueous papaya seed extracts, 395
 ethanolic papaya seed extracts, 395
 papaya latex/unripe fruit extracts, 394
 plant preparations, 394
 ripe papaya fruit safety, 395
 teratogenicity studies, 395
 uterine tissues, 395
 males
 animal trials, 396
 biochemical parameters and contractile pattern, 396
 chloroform-and water extracts, 396
 folklore, 395
 short-term administration
 manifestation, 396
- Antifungal activity
Candida albicans, 397
 chitinases, 397
 gas chromatographic analysis, ripening
 papaya fruit, 397
 green fruit, 396–397
Phytophthora parasitica, 397
- Anti-hypertension, 402
- Antimalarial activity, 397–398
- Antimicrobial activity
 antibiotic-resistant genes, 399
 BITC, 398
 cup plate method, 398
 cyanide-producing glycosides, 399
 cyanogenic glycosides, 398
 ethanol extracts, 398
 gram-positive and negative bacteria, 398
 horizontal gene transfer, 399
 pharmacological properties, 399
- Applications, molecular markers
 BAC library, 360–361
 Brassicales, 359
 chromosome-specific cytogenetic markers, 361
 codominant marker, 360
 genetic map, 359
 major and minor linkage groups, 360
 physical map, 361
 SSR markers, 359–360
- Ascorbic acid (AA)
 antioxidant capacity, 377
 genotype selection and glucose, 382
 green fruit, 381
 hydrophilic antioxidant activity, fresh
 papayas, 380
 solo-type varieties, 384
 water-soluble antioxidant, ripe papayas, 378
- B**
- BAC end sequences (BES), 333
- BACs. *See* Bacterial artificial chromosomes (BACs)
- Bacterial artificial chromosomes (BACs)
 AU9 male and DNA, 334
 characteristics, 171
 end sequences, 333, 337
 and FISH, 316
 hermaphrodite plant, 171
 from hermaphrodite SunUp, 345
 and HSY, 333
 library, 332, 333, 360–361
 low and reiterative, 312
 male-specific markers, 332
 map-based cloning, chromosome walking,
 333–334
 physically map and sequence, 313
 positive mapping, 334
 Sanger method and X physical map, 335
 sex types, 172
 SH95B12 and SH85B24, 335
 SH86B15 and SH85C03, 335
 SH52F06 and SH58C24, 337
 SunUp hermaphrodite, 336
- Benzyl-isothiocyanate (BITC)
 anthelmintic effect, 392
 antifungal activity, 397

- CYPE1, 400
 enzymatic hydrolysis, benzyl
 glucosinolate, 398
 induce apoptosis (cell death), 400
 isothiocyanates, 400–401
 seed constituent response, 395
 treat intestinal worms, 403
- Biogeography, Caricaceae
 Africa and Eocene arrival, 87–89
 Mexico/Guatemala, 85–87
 morphological and molecular studies, 82–85
- Biology, papaya
 branching, 18–19
 dehydration postponement, 28–29
 description, 17
 flowers (*see* Papayas)
 fruits, 26–27
 growth and development, 20–21
 laticifer, 22–23
 leaf epidermis and palisade parenchyma, 19
 leaf lamina and petiole, 18, 19
 leaf-level photosynthesis, 28
 nutrient economy, 29
 PPF, 28
 root system, 21, 23
 seeds, 27
 sex expression, 23
 sexual forms, 19
 stem, 18, 19
 sugar content, 28
 support and transport systems, stem, 21–22
 tropical fruits, 28
 vegetative parts, 17–18
 xylem and transpiration rates, 28
- BITC. *See* Benzyl-isothiocyanate (BITC)
- Breeding, solo papaya
 economic realities, 107–108
 reconfiguration, plant architecture
 biological strategy, 108–109
 dormant papaya seed bank, 108
 fertilizer application, 109
 productive genotypes, 108
 resistances, 110
 true-breeding hermaphrodite, 109
 tapping hybrid vigor, 110
 true-breeding hermaphrodites, 110–111
- C**
- Cancer
 antioxidants provide protection, 343
 and cardiovascular disease, 377
 cell death of, 400
 lycopene-rich diets, 380
- Carbon balance, 28
- Carica papaya* L. (Papayas). *See also*
 Solo papaya
 Asian countries, production, 14
 CPFC1, 386
 cytogenetics, Molecular cytogenetics
 genetic variation, 384
 germplasm, 385
 harvested area, 9, 10
 map (*see* Physical mapping)
 production, 9–10
 provitamin A pigments range, 384
 trade, 11–13
 tropical fruits, 11, 12
 and *Vasconcellea*, 383
 yields, 9, 10
- Carica papaya* sex-linked markers
 (CPSMs), 333
- Carica* with *Vasconcellea*
 breeding, introgression (*see* Introgression
 breeding, *Vasconcellea* species)
 cytogenetics and interspecific
 hybridization, 54
 description, 48
 genetic divergence (*see* Genetic
 divergence, *Carica* with
 Vasconcellea)
 genetic diversity (*see* Genetic diversity,
 Vasconcellea)
 germplasm, breeding, 58–59
 morphology and floral biology, 53
 observations and data, 47–48
 reproductive barriers, 60–61
 taxonomy, 52–53
- β -Carotene
 chloroplasts and peppers, 342
 color difference, 343
 cultivar “Sunrise”, 383
 lycopene, 343, 345, 347, 348
 non-esterified and esterified, 381
 papaya carotenoids, 377
 vitamin A, 382
 and xanthophylls, 385
- ζ -Carotene, 377
- Carotene biosynthetic pathway
 BACs, 347
 characterization, 343
 chloroplasts, 342, 344
 chromoplasts, 344
 citrus plants and *CpLCY-B*, 346
 color complementation and full-length
 coding regions, 348
 consumer’s choice, 349
CpPDS and *CpZDS*, 350–351
Cs β -LCY2 and dual regulation, 343
 enzymatic reactions and plastids, 344

- Carotene biosynthetic pathway (*cont.*)
 flesh color gene, 345–346
 IPP and isomer, 344
lcy-β1 and *lcy-β2*, 349, 350
 lycopene and mapping genes, 345
 map-based cloning, 346–347
 nucleotide and amino acid sequence,
CpCYC-b, 347–348
 PCR amplification and lycopene β-cyclase
 genes, 349
PDS and *ZDS*, 350
 phytoene, 344–345
 plants and health benefits, 343
 PSY, 351
 qPCR and tomato, 346
 SNP, 349–350
 SunUp papaya BAC (SH18O09) and AU9
 BAC (DM105M02), 348–349
- Carotenoid cleavage dioxygenase (CCD), 242
- Carotenoids
 active retinol, 379
 antioxidant capacity and vitamin C, 377
 biosynthesis, 250–251
 biosynthetic pathway, 381, 384–386
 chilling temperatures, 382–383
 β-cryptoxanthin, 381
 human health and vitamin C, 380
 provitamin A, 378
 synthesis, 382
 types, 377
 vitamin A precursors and antioxidants, 378
 yellowed-fleshed types, 384
- CCD. *See* Carotenoid cleavage dioxygenase
 (CCD)
- Cell wall synthesis and degradation
 carbohydrate hydrolases, esterases and
 lyases, 247, 248
 cellulose-related structural genes, 246
 galacturonosyltransferases, 247
 hemicellulose degradative enzymes, 249
 matrix polysaccharides and pectins, 245
 plastic and elastic extension, 247
 starch degradation, 249
 transferases, carbohydrates, 246
 triplicate replication, 246
- Centromeres
 coevolution, 164
 locations, papaya chromosomes, 162
 pachytene chromosome, 160
- Chlorophyll, skin degradation, 249–250
- Chymopapain
 nucleus pulposus, 394
 proteolytic enzymes, 392, 393
- Climacteric fruit, 27
- Cloning
 anthocyanins, 342
 carotene biosynthetic pathway (*see*
 Carotene biosynthetic pathway)
 fruit color (*see* Fruit flesh color)
 fruit ripening, 341, 342
 lycopene and peppers, 342
- CNS. *See* Conserved noncoding sequences
 (CNS)
- CODDLE. *See* Codons to optimize
 discovery of deleterious lesions
 (CODDLE)
- Codons to optimize discovery of deleterious
 lesions (CODDLE), 412
- Coexistence, nontransgenic papaya for Japan,
 132–133
- CoGe
 dotplot, papaya and peach, 209
 high resolution, syntenic genomic regions,
 216–217
 web-based comparative genomics
 platform, 206
- Cold resistance, 66, 67, 74
- Comparative genomics
 genome assemblies, 205
 molecular cytogenetic tools, 158
 physical maps, 169
 sex chromosome evolution, papaya, 201
- Conserved noncoding sequences (CNS)
 comparison, syntenic set of gene, 221
 DNA-binding factors, 220–221
 orthologous syntenic genes, 222
 protein-coding sequences, 221
- CpCYC-b* gene
 chloroplast-specific version, 342
 description, 347
 ORFs, 344
 polymorphisms, 347
 tomato *CYC-b*, 347
- CPSMs. *See* *Carica papaya* sex-linked
 markers (CPSMs)
- Crop improvement
Carica and *Vasconcellea*, 36
 clear-cut traits, 35
 genetic polymorphism, 35
 magnitude and structure, genetic diversity,
 35–36
- β-Cryptoxanthin, 377–381, 383, 384
- Cultivar development
 “Kapoho” x “Sunrise” hybrid, 104
 “Sunrise” and “Sunset”, 102–103
 transgenic “SunUp” and “Rainbow”,
 104–106
 “Waimanalo” and “Kamiya”, 103–104

D

DAPI. *See* 4',6-Diamidino-2-phenylindole (DAPI)
Dengue fever, 399

Deregulation, Hawaii transgenic papaya
China, 139

and commercialization

FDA and EPA, 126–127
field trial, Kapoho, 124–126
license agreements, 128
marketing, 127
PAC, 127
“red zone”, 124

and export

Canada, 135
Japan, 135–136
MAFF, 136
MHLW, 136–139

Diabetes

ALT, 402
blood glucose and blood lipids, 401
hypoglycaemic agents, 402
unripe papaya fruit, 401

4',6-Diamidino-2-phenylindole (DAPI)

cytological distribution, 161
meiotic pachytene chromosomes, 160, 161
pericentromeric heterochromatin., 162

Dietary reference intake (DRI), 377, 378, 379

Disease resistance, marker assisted breeding

DNA, 368
mapping, 368
Opa11_5R and Opk4_1r, 369

prsv-1, 368

PRSV-P (*see* Papaya ringspot virus
type P (PRSV-P))

Vasconcellea spp., 367

Diversity

crop improvement, 35–36
genetic, 39–43
phenotype, 36–39

DNA polymorphism

AFLP (*see* Amplified fragment length
polymorphism (AFLP))
RAPD markers (*see* Randomly amplified
polymorphic DNA (RAPD))
RFLP, 39, 143–144, 358, 362

DNA profiling method, 226–227

Domestication

chichihualtzapotl, 8
cultivars and papaya trees, 8
germination rates, 8
maize and *Agave fourcroydes*, 6, 7
rain and temperatures, 8
well-developed agricultural systems, 7

DRI. *See* Dietary reference intake (DRI)

E

Early agriculture, 7

Environmental Protection Agency (EPA),
126, 127

EPA. *See* Environmental Protection Agency
(EPA)

EST sequencing and gene prediction

ab initio gene prediction software, 193
abiotic stress responses, 195
cDNA clones, 193
DAPI, 192
global protein analysis, 194
papaya *vs.* sequenced plant genomes, 194
plant processes, 195
repetitive elements, 194
rice draft genome, 194
TF, 195

Ethylene synthesis

aminotransferases I and II domains, 255
Arabidopsis proteins, 255
copper cofactor, 257
protein prediction, 254–255
receptor genes, 256
response pathway, 257
SAMS, 254
signal transduction pathway, 257

Evolution and fruit ripening

gene clusters, 263
gene-controlled process, 263
genome-wide changes, 264
molecular and biological processes, 263
plant biology, 264
variation, species and cultivars, 263

Expressed sequence tagged sites (ESTs).

See EST sequencing and gene
prediction

F

FDA. *See* Food and Drug Administration
(FDA)

FISH. *See* Fluorescence in situ hybridization
(FISH)

Flesh color. *See* Fruit flesh color

Fluorescence in situ hybridization (FISH)
based chromosome identification, 159
centromere-specific sequences, 162
genetic maps, 150
high-resolution pachytene, 160
karyotyping, 160
mitotic and meiotic chromosomes, 163
positive BACs to Y^h chromosome, 334
probes, 159

Food and Drug Administration (FDA), 126, 127

- Foreign competition, PRSV-resistant varieties, 107, 108
- Fractionation, gene loss polyploidy
 annotated plant genomes, 220
 gene content, Arabidopsis, 218–219
 multiple syntenic genomic regions, 219
 orthologous sequences, Papaya vs. Arabidopsis, 220
- Fruit crops
 in Andes, 48–49
 babaco production, 50
 grafting techniques, 50
V. cundinamarcensis, 49–50
V. stipulata, 50
V. x heilbornii “Babaco”, 49
- Fruit flesh color
 acrylamide sequencing gels, 179
 AFLP, 178
 animals and green skin, 341
 carotenoid antioxidants, 177
 commercial value and molecular markers, 349
C. papaya, 361
CpLCY-B and *CrtL-b*, 346
 degenerate primers, 345–346
 de novo synthesis, color pigments, 343
 dinucleotide insertion mutation, 351
 219 F₂ (KD3 x 2H94), 370
 fruit ripening and genes regulation, 346
 high-density AFLP genetic map, 178
 high-density genetic map and SSR markers, 345
 linkage groups, 360
 lycopene β -cyclase gene, 347
 map-based cloning, 346–347
 mechanism and regulation, 345
 Mendelian fashion, 351
 papaya marketing, 370
 red and yellow, 342
 SCAR marker, 179
 and sex type, 359
 steroid hormones, 177
 SunUp papaya BAC (SH18O09) and AU9 BAC (DM105M02), 348–349
 yellow-fleshed allele, 178
- Fruit growth
 ABA, 242
 carotenoid cleavage, 242
 CCD, 242
 cell wall synthesis and degradation, 245–249
 climacteric fruits, 242
 ethylene production, 241
 expansins, 244–245
 genetic characteristics, 241, 242
 mesocarp carotenoid development, 243
 plasticity, papaya size, 243
 respiration, 241
 ripening, 249–262
 shape, 243–244
- Fruit quality traits
 Australian papaya industry, 370
 cloning and characterization, 370
CpCHY-b, 370
CpCYC-b, 370
 DNA markers, 370, 371
 flesh color, 370
- Fruit ripening
CpLCY-B, 342
Cs β -LCY2, 343
 flesh color gene, 346
 red and yellow papaya, 347
 symptoms, 341, 342
- Fruits
 climacteric, 27
 female and hermaphroditic plant, 26–27
 hermaphroditic flower, 26
 laticifers, 22–23
 photosynthesis, carbon gain, 27
 stamens, 25
 sugar content, 28
 tropical pioneer, 18
- Functional genomics, 410–411
- G**
- GBIF. *See* Global Biodiversity Facility (GBIF)
- Gene pairs
 dotplot visualization, 210, 211
 evolutionary relationships, 211, 212
 papaya vs. peach, 212–214
 synonymous substitution values, 212–213
- Genetic divergence, *Carica* with *Vasconcellea*, 58
- Genetic diversity
 advantage, molecular methods, 39, 40
 AFLP, 39
 breeding lines, 39
Caricaceae and molecular markers (*see* Molecular markers, *Caricaceae*)
C. papaya and *Caricaceae* species, 41
 description, 43
 genetic similarity, papaya cultivars, 41, 42
 Hawaiian “Solo” papayas, 43
 microsatellite markers, 41
 pair-wise comparisons, 43
 papaya lines, Costa Rica, 41
 RAPD, 39

- USDA germplasm collection, 43
- Vasconcellea*
- Colombian samples, 57–58
 - description, 73
 - hybridizations and introgressions, 54
 - nuclear and chloroplast datasets, 55
 - specimens, taxa, 57
 - staminate flowers, 54
 - “syngameons”, 55
 - V. chilensis*, *V. candicans*, 56–57
 - V. pulchra* and *V. longiflora*, 57
 - V. quercifolia* and *V. glandulosa*, 56–57
 - V. stipulata* and *V. cundinamarcensis*, 57
 - V. stipulata* and *V. parviflora*, 55–56
 - V. x heilbornii*, *V. weberbaueri*, 55–56
- Genetic engineering
- Hawaii’s papaya cultivars, PRSV resistance, 105
 - papaya germplasm, 109
- Genetic erosion and *Vasconcellea* conservation
- ex situ* conservation, 72
 - in situ* conservation, 71
- Genetic mapping
- BLASTN search, genome sequence, 150
 - characteristics, papaya from 1939–2009, 145
 - codominant markers, 360
 - and cytogenetic
 - chromosome-specific markers, 175
 - karyotyping methods, 174
 - metaphase stage, 174
 - NOR, 176
 - cytological data, 150
 - and genome sequence, 361
 - genome sequence data, 150
 - high-density, 143
 - hypermethylation, 147
 - integrated physical maps, 150
 - LGs, 151–152
 - marker-assisted selection, 147
 - molecular marker development, 358–359
 - molecular marker types, 150
 - morphological markers, 359
 - papaya chromosome characteristics, 153
 - papaya comparison, 170
 - PCR amplification, 144
 - physical and genome sequence, 173–174
 - PRSVCO, 146
 - PRSV-P resistance gene(s), 368
 - RAPD, 144
 - RFLP, 144
 - SDR, 146
 - segregation distortion, 148
 - sequence-tagged, 171
 - sex chromosome evolution, 145
 - SSR, 171
 - whole genome sequence, 149
- Genetic recombination
- FISH, 177
 - restriction enzyme fingerprinting data, 176–177
 - suppression, 177
- Genetic resources, 35
- Genetics of sex, 98
- Genomics
- comparative (*see* Comparative genomics)
 - web-based comparative genomics platform, CoGe, 206
- Genotypic variation, nutritional composition
- C. papaya*, 383–384
 - lycopene, 384–385
 - in nutrient quality, 384
 - production characteristics, 384
 - provitamin A pigments ranges, 384
 - red-fleshed varieties contain lycopene, 384
 - vitamin C content ranges, 384
- Global Biodiversity Facility (GBIF), 4, 5
- H**
- Hawaii’s transgenic papaya
- biotechnology, 139–140
 - coexistence, 132–133
 - cultivars grown, 131
 - deregulation and commercialization (*see* Deregulation, Hawaii transgenic papaya)
 - development, 117–119
 - environmental impact, 132
 - nontransgenic papaya, 131–132
 - pathogen-derived resistance approach, 116–117
 - production, Puna, 128–131
 - PRSV (*see* Papaya ringspot virus (PRSV))
 - resurgence, 132
 - R1 plants, line 55–1, 119–120
 - R0 plants, line 55–1, 120
 - “SunUp” and “Rainbow”, Oahu, 123–124
 - transgene flow, “Rainbow” and “Kapoho”, 133–135
- Hermaphrodite-specific Y chromosome (HSY)
- and BACs (*see* Bacterial artificial chromosomes (BACs))
 - BES and CPSMs, 333
 - characteristics, 332
 - description, 333
 - and linkage group 1 (LG1), 337
 - male-specific/sex-co-segregating markers, 333

- Hermaphrodite-specific Y chromosome (HSY) (*cont.*)
 map-based cloning, chromosome walking, 334
 non-recombining sex determination region borders, 335
 physical mapping and FISH, 334
 SH86B15 and SH85C03, 335
 SH94G23 and SH68N07, 335
 SSR markers, 335, 337
 and X chromosomes (*see* X chromosomes)
 Y^h chromosome, 332
- Hermaphroditic flowers
 staminate flowers (sf), 20, 24
 stresses, 24
- Heterochromatin
 cytological distribution, DAPI, 161
 definition, 160
 knob-like structures, 162
 pericentromeric regions, pachytene chromosomes, 160, 161
 types, repetitive DNA, 160
 Y chromosomes, 226
- Highland papayas
 description, 48
 distribution and climatic preferences, 51–52
 fruit crops, 48–50
 medicinal uses, 51
 sources, papain, 50–51
 wild and feral populations, 51
- HSY. *See* Hermaphrodite-specific Y chromosome (HSY)
- Hybridization and cytogenetics
 chloroplast, 55
 cytogenetics and interspecific, 54
 Fluorescence in situ, 73
 fruit and papain production, 58
 seed germination, 60
V. cauliflora, 62–63
V. parviflora, 68
V. quercifolia, 57
V. x heilbornii, 49, 56
- Hypertension, 391, 402
- I**
- Internal transcribed spacer (ITS), 84
 Inter simple sequence repeat (ISSR) technique, 367
- Introgression breeding, *Vasconcellea* species
 description, 73
 traits, interest, 59
V. cauliflora, 61–63
V. cundinamarcensis, 66–67
V. glandulosa, 71
V. goudotiana, 69–70
V. microcarpa and *V. horovitziana*, 70
V. monoica, 68–69
V. parviflora, 67–68
V. quercifolia, 63–66
V. stipulata, 69
- ITS. *See* Internal transcribed spacer (ITS)
- K**
- Kapoho field trial, 124–126
- Karyotypes
 chromosome-specific cytogenetic markers, 159
 cytological preparation techniques, 159
 FISH-based, 160
 heterochromatin, 158
 integration, genetic and cytological maps, 159
 primary constriction, 159
- L**
- Lander–Waterman theory, 207
- LGs. *See* Linkage groups (LGs)
- Linkage groups (LGs)
 chromosome number assignment, 151–152
 mitotic and meiotic chromosomes, 159
 telomeric regions, 150
- Lycopene
 carotenoid biosynthetic pathway, 384
 and carotenoids (*see* Carotenoids)
 fruit, 381, 383
 levels range, 380
 red-fleshed varieties, 377, 380, 384
 ripe “Maradol” and “Golden” fruit, 384
 yellow-fleshed cultivars, 385
- Lycopene β-cyclase
arabidopsis and *lcy-β1*, 349
 chromoplast-specification (*see* *CpCYC-b* gene)
CpLCY-B and degenerate primers, 346
 fruit-specification and *CYC-b*, 347
 functioning, leaf and fruit, 346
 gene encoding enzyme, 351
- M**
- MADS-box family
 developmental functions, 195
 evolutionarily distinct lineages, 195
 flower development, 195
 functionality, 196

- ovule and seed development, 196–197
 - transcription factor families, 196
 - type I and II, 196
 - MAFF. *See* Ministry of Agriculture Fisheries and Forestry (MAFF)
 - Male-specific Y chromosome (MSY)
 - and HSY, 332
 - and YY/WW genotype, 331
 - Marker assisted breeding
 - disease resistance, 367–369
 - fruit quality traits, 370–371
 - hybrid identification, 369–370
 - Medicinal plant
 - anthelmintic activity, 392–393
 - anti-amoebic activity, 401
 - anticancer and chemoprotection, 400–401
 - antifertility properties, 394–396
 - antifungal activity, 396–397
 - anti-hypertension, 402
 - antimalarial activity, 397–398
 - antimicrobial activity, 398–399
 - antiviral activity, 399
 - diabetes, 401–402
 - “folk” remedies efficacy, 391
 - plant tissue, 391
 - tropical and subtropical regions, 391
 - wound healing, 393–394
 - Mexican papaya clade, 89
 - MHLW. *See* Ministry of Health Labor and Welfare (MHLW)
 - Microsatellite markers, 174
 - Ministry of Agriculture Fisheries and Forestry (MAFF), 136
 - Ministry of Health Labor and Welfare (MHLW), 136–139
 - Model fruit trees
 - characteristics, 188
 - cysteine protease, 188–189
 - draft genome, 189
 - genome sequencing project, 187
 - hermaphrodite plants, 188
 - nutritional value, 188
 - papaya consumption, 188
 - Molecular clock dating, *Caricaceae*, 87–89
 - Molecular cytogenetics
 - classical, 158
 - cytological structure, 162–164
 - distribution, heterochromatin, 160–162
 - karyotype, 158–160
 - locations, centromeres, 162
 - sex chromosome evolution, 158
 - techniques, 158
 - Molecular markers
 - AFLP, 358
 - breeding program, 355
 - Caricaceae*
 - AFLP, 362
 - isozyme and RAPD techniques, 362
 - microsatellite-enriched library, 363
 - PCR techniques, 362
 - phenotypic, morphological and horticultural characteristics, 361–362
 - SSR, 363
 - taxonomy studies, 363
 - Vasconcellea*, 364
 - DNA markers, 355–357
 - isozymes, 358
 - microsatellite, 358
 - PCR, 358
 - RAPD, 358
 - RFLPs, 358
 - sex determination
 - DAF, 365
 - decamer primers, 366
 - DNA markers, 364
 - genetic linkage map, 365
 - hermaphrodites, 364–365
 - ISSR technique, 367
 - male and female plants, 364
 - microsatellites, 367
 - OPC09 and OPE03, 366–367
 - OPT1C and OPT12, 365
 - PCR assay, 366
 - primer BC210, 366
 - primer OPA06, 365
 - RAPD techniques, 366
 - SCAR technique, 365
 - single dominant gene with alleles, 364
 - SSRs, 355, 358–359
 - Monoecy, *Vasconcellea monoica*, 53, 68, 69, 73
 - MSY. *See* Male-specific Y chromosome (MSY)
- N**
- Nontransgenic papaya, Japan
 - coexistence, 132–133
 - production, Puna, 131–132
 - NOR. *See* Nucleolus organizing regions (NOR)
 - Nucleolus organizing regions (NOR), 176
 - Nutritional composition
 - AA (*see* Ascorbic acid (AA))
 - antioxidant capacity, 380
 - carotenoids (*see* Carotenoids)
 - environmental effects, 382–383
 - flesh color, 385

- Nutritional composition (*cont.*)
 and fruit development
 AA and carotenoid, 381
 climacteric-type fruit, 380
 “Maradol” and vitamin A, 381
 sugar content and TSS, 380
 vitamin C, 380–381
 genetic characterization, ripening and color
 development genes, 386
 genotypic variation, 383–385
 lycopene (*see* Lycopene)
 transgenic techniques, 385
 value ranges, 378, 379
 vitamin A (*see* Vitamin A)
 vitamin C (*see* Vitamin C)
- O**
 Open reading frames (ORF), 254
 ORF. *See* Open reading frames (ORF)
 Origin, papaya
 analysis, morphometric characters, 6, 7
 Caricaceae family, 3–4
 description, 3
 GBIF, 4
 Herbarium specimens, 4
 Mesoamerican, 5
 morphometric measurements, 5, 6
 subtropical and tropical environments, 3
 wild papaya populations, Yucatan
 Peninsula, 5, 6
- P**
 PAC. *See* Papaya Administrative Committee
 (PAC)
 Paleopolyploidy
 angiosperms, 206
 collinear signal, 206
 definition, 206
 Papain
 and chymopapain, 393
 and ficin, 393
 Food and Drug Administration, 395
 formulations, 395
 intradisc injection to treat lumbar
 osteochondrosis patients, 394
 proteolytic enzymes, 392
 wound healing, 403
 Papaya Administrative Committee (PAC),
 127, 128, 136
 Papaya repeatome
 analysis, high copy number genes, 236
 annotation, TEs, 236
 architectonic function, 226
 biological analysis, 226
 centromeres, 233–234
 classes, repetitive sequences, 225
 constitutive heterochromatin, 226
 C-value paradox, 225
 data access and retrieval, 236–237
 DNA profiling method, 226–227
 gene families, 232
 genetic and physical map, 227
 PCR, 227
 phylogenetic relationships, 227
 plant genome evolution, 226
 sex-chromosome repeatome, 234–235
 sources, 235–236
 tandem repeats, 229–232
 telomeres, 232–233
 TEs, 227–229
 Papaya ringspot virus (PRSV)
 and ADP, 414–415
 coat protein gene, 117
 and cross protection, 116
 PDR, 117
 in Puna
 cultivars grown, 131
 detection and spread, 120–121
 environmental impact, 132
 farmer adoption, 128
 infected “Kapoho” field, 130
 nontransgenic papaya, 131–132
 production and industry, 122–123
 “Rainbow”, 128, 129
 resurgence, 132
 transgenic field trial, 124–128
 transgenic papaya, 131
 varieties grown, 129, 130
 R0 plants, line 55–1, 120
 Papaya ringspot virus coat protein marker
 (PRSVCO), 146
 Papaya ringspot virus type P (PRSV-P)
 DNA markers, 369
 genetic mapping, 368
 GMOs plants, 368
 RAF markers, 368
 single dominant gene, 368
 transgenic papayas resistant, 369
 Papayas (*Carica papaya* L.) *See also* *Carica*
papaya L. (Papayas)
 ADP (*see* Allele discovery platform
 (ADP))
 flowers
 female, 20, 25, 26
 hermaphroditic, 24–25
 male, 19, 20, 26

- fruit flesh color (*see* Fruit flesh color)
- genome
 - syntenic sequences (*see* Syntenic sequence conservation)
- medicinal plant (*see* Medicinal plant)
- molecular markers (*see* Molecular markers)
- nutritional analysis (*see* Nutritional composition)
- physical mapping (*see* Physical mapping)
- sex chromosomes (*see* Sex chromosomes)
- Papaya transgene flow, 133–135
- PCR. *See* Polymerase chain reaction (PCR)
- PGR. *See* Plant growth regulator (PGR)
- Phenotype diversity
 - flower and fruit characteristics, 36, 37
 - fruit shape, 38
 - Hawaiian Solo lines, 38
 - hermaphrodite plants, 38
 - inflorescence peduncles, 36
 - morphological and isozyme idiotypes, 38–39
 - sex expression, 38
- Phenotypic plasticity, 17, 18, 23, 29
- Phenotypic traits, 35
- Photosynthetic photon flux density (PPFD), 28
- Phylogeny, Caricaceae
 - Africa and Eocene arrival, 87–89
 - and Moringaceae, Brassicales, 81
 - morphological and molecular studies
 - Carica papaya*, 83
 - description, 84, 85
 - distribution, Africa and Neotropics, 82
 - economic importance, 84
 - ITS, 84
 - Jacaratia*, 82
 - Jarilla*, 82–83
 - Vasconcellea*, 82, 84
 - papaya origin, Mexico/Guatemala
 - ancestral area reconstruction and molecular clock dating, 86, 87
 - ecological/morphological studies, 86
 - nucleotides, 85
 - Vasconcellea x heilbornii*, 85–86
- Physical mapping
 - BAC-anchored markers, 169
 - dioecious species, 329, 330
 - draft genome sequence, 179
 - flowering plants, 329
 - fruit flesh color, 177–179
 - genetic map and genome sequence, 173–174
 - genetic recombination, 176–177
 - genome coverage, 176
 - heterochromatic, 338
 - high-density genetic map, 170
 - hybridization, overgo probes, 169
 - integration, genetic and cytogenetic maps, 174–176
 - monoecious species, 329
 - MSY (*see* Male-specific Y chromosome (MSY))
 - overgo and single-copy probe hybridization, 173
 - papaya BAC libraries, 171–172
 - QTLs, 179
 - and sequencing
 - BACs (*see* Bacterial artificial chromosomes (BACs))
 - HSY (*see* Hermaphrodite-specific Y chromosome (HSY))
 - knob structures, 316
 - molecular position, 315
 - pseudo-autosomal regions, 314–315
 - repetitive element accumulation and expansion, 319–322
 - SSR markers and MSY, 315
 - sequencing technologies, 169
 - sex chromosomes (*see* Sex chromosomes)
 - sex determination (*see* Sex determination)
 - unisexual flower development, 330
- Phytoene synthase (PSY), 344, 351
- Plant growth regulator (PGR)
 - auxin-related genes, 262
 - brassinosteroid-related genes, 263
 - ethylene receptors, 262
- Polymerase chain reaction (PCR)
 - assay, 366
 - description, 358
 - techniques, 362
- Polyploidy
 - Eurosid phylogeny, 207
 - gene loss problem, 207
 - genetic features, 206
 - sequential, 207
 - statistical distribution, 206
- PPFD. *See* Photosynthetic photon flux density (PPFD)
- Production
 - description, 14
 - harvested area and yields, 9–10
 - tropical fruits, 11, 12
- PRSV. *See* Papaya ringspot virus (PRSV)
- PRSVCO. *See* Papaya ringspot virus coat protein marker (PRSVCO)
- PRSV-P. *See* Papaya ringspot virus type P (PRSV-P)
- PSY. *See* Phytoene synthase (PSY)
- Public Sector GMO, 135

Q

- qPCR. *See* Quantitative real-time PCR (qPCR)
 QTLs. *See* Quantitative trait loci (QTLs)
 Quantitative real-time PCR (qPCR), 346
 Quantitative trait loci (QTLs), 371

R

- “Rainbow” and “SunUp”, R0 field trial on
 Oahu
 Canada, 135
 F1 hybrid, 131
 Japan, 135–136
 Kapoho field trial, 128
 and “Laie Gold” papaya, 132
 PRSV-infected “Kapoho” field, 130
 PRSV-infected nontransgenic papaya, 126
 reclamation, Puna papaya lands, 129
 resistance, 126
 R0 field trial, Oahu, 123–124
 transgene flow, 133–135
 US Embassy, 138
 virus-resistance performances, 125
 Randomly amplified DNA fingerprinting
 (RAF) markers, 368
 Randomly amplified polymorphic DNA
 (RAPD)
 fragmentation, genomic DNA, 144
 genetic mapping, 144
 Hawaiian Solo cultivars and papayas, 39
 inflated map distances, 146
 and isozyme, 44
 markers, 358
 PCR amplification, 144
 QTLs, 153
 techniques, 366
 RAPD. *See* Randomly amplified polymorphic
 DNA (RAPD)
 Repetitive accumulation and expansion
 Arabidopsis genome, 321
 DNA transposons and LTR, 320
 elements, papaya sex chromosomes, 320
 embryo abortion, Y^hY^h genotype,
 321–322
 hermaphrodite papaya fruits, 321
 organelle genome integration, 321
 palindromic inverted repeats, 321
 rapid and dramatic, 319
 retrotransposons and Ty3-gypsy
 elements, 320
 X and HSY chromosomes, 321
 Repetitive DNA
 accumulation, 234
 centromere-specific sequences, 233
 classification, 227

- phylogeny, 227
 Reproductive barriers, *Carica* and
 Vasconcellea
 fruit and seed development, 60
 hybrid embryos and endosperms, 60
 sesquidiploids, 61
 species, 60
 sterility, 60–61
 zygotic embryos, 60
 Respiration
 carbon metabolism and electron transport,
 253–254
 monocots, 253
 ORF, 254
 plant mitochondria, 254
 Restriction fragment length polymorphisms
 (RFLPs)
 cpDNA intergenic spacer region, 39
 fragmentation, genomic DNA, 144
 genetic base, 144
 molecular markers, genetic mapping,
 143–144
 in papaya genomic study, 358
 phylogenetic relationships, 362
 RFLPs. *See* Restriction fragment length
 polymorphisms (RFLPs)
 Ripening, fruit
 carotenoid biosynthesis, 250–251
 ethylene
 response, 256–258
 synthesis, 254–256
 and evolution, 263–264
 laticifers and protease, 258–259
 PGR, 262–263
 and plant growth regulator related genes,
 262–263
 plant respiration, 253–254
 skin chlorophyll degradation, 249–250
 sugar accumulation, 251–253
 volatile production, 259–262
 Root systems, 21, 23

S
 S-adenosyl-L-methionine synthases
 (SAMS), 254
 SAMS. *See* S-adenosyl-L-methionine
 synthases (SAMS)
 Satellite DNA
 centromeres, 233
 diverse repetitive sequences, 234
 phylogenetic relationships, 227
 SDR. *See* Sex-determining region (SDR)
 Sequence-characterized amplified region
 (SCAR) technique, 365

- Sex-chromosome repeatome
 - chromosome walking, 234
 - evolution, 235
 - evolutionary features, 234
 - MSY explosion, 235
 - plant-sex chromosomes, 235
 - sex determination, 234
- Sex chromosomes
 - cytological features, 158
 - draft genome, 189
 - early stage evolution, 164
 - evolution
 - BAC approach and segregation ratios, 313
 - characteristics and mechanisms, 312
 - Chimpanzee Y chromosome, 312
 - co-segregating markers, 314
 - cytologically and karyotype analysis, 314
 - DAPI chromosome staining, 314
 - dioecious plants, 310
 - disease and abiotic stresses, 312
 - female and male sterile mutations, 311
 - flower morphology, trioecious papaya, 310–311
 - genome sequencing projects, 312
 - identification, stamen-promoting gene, 312
 - knob structures pair, 314
 - male/female heterogametics, 309
 - maleness, 312
 - map-based approach, 324–325
 - physical mapping and sequencing (*see* Physical mapping)
 - sex determination genes, 311
 - single gene, 309
 - size and degeneracy, 309–310
 - syntenic analysis, 324
 - true-breeding hermaphrodite, 313
 - evolutionary history, 189
 - female genome sequence, 335
 - HSY (*see* Hermaphrodite-specific Y chromosome (HSY))
 - initiation and degeneration, 189
 - MSY-specific heterochromatin distribution, 163
 - pachytene chromosome lengths, 159
 - pseudo-autosomal region, 189
 - SH53E18 and SH61H12, 334–335
 - X chromosome (*see* X chromosomes)
- Sex determination
 - angiosperm species and chromosomal, 330
 - BAC library, 332
 - characteristics, 332
 - chromosome pairs, 332
 - genetic analysis, 331–332
 - genotyping, 337
 - high-density genetic map, 332
 - individual plants, sex types, 331
 - and molecular markers, DAF (*see* Molecular markers, sex determination)
 - non-recombining, 335
 - papaya flowers, 331
- Sex-determining region (SDR), 146
- Shotgun sequencing
 - high-density genetic map, 191, 200
 - papaya genome
- Simple sequence repeats (SSR)
 - AU9 x SunUp, 360
 - FISH and BAC clones, 361
 - genetic distance, 147
 - genome sequence project, 363
 - libraries, 359–360
 - mapping population, 148
 - microsatellite markers, 358–359
 - and *NBS* gene families, 371
 - physical map and whole genome shotgun sequences, 150
 - primers, 367
 - sequence based, 144
- Single-nucleotide polymorphism (SNP), 349–350
- Sister group of papaya, 81
- Skin chlorophyll degradation, 249–250
- SNP. *See* Single-nucleotide polymorphism (SNP)
- Solo papaya
 - breeding (*see* Breeding, solo papaya)
 - cultivar diversification
 - “Kapoho” x “Sunrise” hybrid, 104
 - “Sunrise” and “Sunset”, 102–103
 - transgenic “SunUp” and “Rainbow”, 104–106
 - “Waimanalo” and “Kamiya”, 103–104
 - geographic dispersal, 96
 - germplasm deficiencies and genetic solutions
 - fruit carpellody and carpel abortion, 99
 - objectives, 99
 - sex segregation, 98
 - growth, export industry
 - fruit fly disinfestation, 100
 - gross production, 100
 - “Kapoho” papaya, 100–101
 - tourism connection, 101–102
- Hawaii
 - arrival, 97
 - characteristics, 97–98
 - description, 96
 - gynodioecious papayas, 97
 - origin and domestication, 95–96
 - taxonomy, 95

- SPS. *See* Sucrose phosphate synthase (SPS)
- SSR. *See* Simple sequence repeats (SSR)
- Stamen carpellody, 99, 109, 110
- Sucrose phosphate synthase (SPS), 253
- Sucrose synthase (SUS), 252
- Sugar accumulation
- hexose transporter genes, 253
 - invertase (INV), 252
 - photosynthates, 251
 - SPS, 253
 - SUS, 252
 - symplast pathway, 251
- “SunUp” and “Rainbow”, R0 field trial on Oahu, 123–124, 126, 131, 137
- SunUp variety, papaya
- Agrobacterium-mediated and biolistic approaches, 200
 - chloroplast sequences, 200
 - genetic engineering, 199
 - papaya genome sequencing project., 189
 - PRSV, 189
 - segregation patterns, 199
 - variable integration site, 199
- Suppression of recombination
- homologous pairing, 314
 - HSY, 316, 320, 324
 - large-scale inversions, 324
 - Meullers ratchet, 322
 - sex determination loci, 309
 - sex specific regions, 317
 - Y^h chromosomes, 322
- Syntenic sequence conservation
- autopolyploidy, 206
 - CNS, 220–222
 - computational techniques, 223
 - Eurosid phylogeny and polyploidy, 207
 - evolutionary history, 207
 - fractionation, gene loss polyploidy, 218–220
 - gene pairs, 211–214
 - high-resolution analysis
 - collinear arrangement, homologous genes, 215, 216
 - dotplot, CoGe, 216–218
 - parallel line coordinates visualization scheme, 214
 - intragenomic syntenic regions, 206
 - morphological differences, 223
 - paleopolyploidy events, 206
 - per-base sequencing cost, 223
 - syntenic path assembly, 210–211
 - whole-genome comparison, 207–209
- Synteny analysis
- alignment and subalignment, 198
 - alignment, collinear regions, 198
 - alignment of collinear regions, 198
 - genome duplication events, 199
 - phylogenetic dating and collinearity, 199
 - polyploidization, 198
- T**
- Tandemly-arrayed sequences. *See* Tandem repeats
- Tandem repeats
- angiosperms, 230
 - detection, 236
 - gene density, 229
 - gene regulation/inactivation, 230
 - physical quantity, DNA, 230
 - SSR sampling
 - EST sequences, 231
 - marker-assisted selection, 232
 - repetitive motif, 230
 - types, 230–231
 - WGS, 229
- Targeting induced local lesion in genome (TILLING), 410, 414–415
- TEs. *See* Transposable elements (TEs)
- TF. *See* Transcription factors (TF)
- TILLING. *See* Targeting induced local lesion in genome (TILLING)
- Transcription factors (TF), 195
- Transgenic papaya
- in Hawaii (*See* Hawaii’s transgenic papaya)
 - in Japan, 131, 135–136
 - genome
 - BAC-end sequences, 190
 - characterization, SunUp, 199–200
 - EST sequencing and gene prediction, 192–195
 - genetic mapping and scaffold anchoring, 191–192
 - MADS-box family, 195–197
 - model fruit tree, 188–189
 - physical mapping, 190
 - QTL, 187
 - repeat composition, 197–198
 - SunUp variety, 189
 - synteny analysis and genome-wide duplications, 198–199
 - whole shotgun sequencing approach, 189
- Translational biotechnology, 139–140
- Transposable elements (TEs)
- phylogenetic analysis, 228, 229
 - plant genomes, 227
 - sex chromosomes, 228
 - types, 228
 - WGS data, 227

U

USDA. *See* U.S. Department of Agriculture (USDA)

U.S. Department of Agriculture (USDA)
germplasm collection, Hawaii, 43
marketing order agreement, 127
nutritional values, papaya fruit, 378, 379

V

VAD. *See* Vitamin A deficiency (VAD)

Vasconcellea species

AFLPs, 39
breeding lines, 39
description, 36
developments, Caricaceae family, 74
and genetic erosion (*see* Genetic erosion
and *Vasconcellea* conservation)
genetic resources, 47
highland papayas (*see* Highland papayas)
RAPD, 39, 44
relationship with *Carica* (*see Carica*
with *Vasconcellea*)
sexual incompatibility, 44
taxonomical confusion, 47

V. cauliflora

aneuploids, 63
characterization, 62
climatic requirements, 61
genetic resistance, PRSV-P, 61
hybridization, 62–63
papaya mosaic virus, resistance, 59
planting, 62
sex expression, 62
variation, vigor, 62

V. quercifolia

backcross populations, 63–64
cpDNA level, 56
cross *C. papaya*, 63
genotypes, 64
hybrids and backcross plants, 64, 65
PRSV-P resistance, 65–66
seed germination, 64
sib-crossing, 66
viral symptoms, 64
wide adaptation, 63

Vitamin A

β -carotene and IOM, 379
and carotenoid content, 377
carotenoid synthesis and accumulation, 381
DRI and VAD, 377
environmental factors, 382
genotypic variation, 383
precursors and antioxidants, 378

ranges, 378

yellow-fleshed papayas, 381

Vitamin A deficiency (VAD), 409

Vitamin C

AA synthesis, 382
and carotenoids, 380
DRI and carotenoids, 377
fruits and vegetables, 378
gene encoding, δ -galacturonic acid
reductase, 386
nutritional benefit, 378
ranges, 384
ripened on-tree vs. off-tree, 381
seasonal variations in climate, 382
solo-type cultivars grown in Hawaii, 378

Volatile production, papaya

cleavage, carotenoids, 262
cytoplasmic terpene synthesis, 262
ecological interactions, 259–260
esters, 260
flavor and aroma perception, 260
linalool production, 260
mesocarp alcohol dehydrogenase
activity, 260
straight chain ester biosynthesis, 261
terpene synthesis pathway, 262

W

Water relations

dehydration postponement, 28
PPFD, 28
solar radiation, 28

Whole-genome comparison

dotplot, papaya and peach, 209
Lander–Waterman theory, 207
papaya–peach syntenic dotplot, 208

Whole-genome shotgun sequences (WGS), 229

Wild papaya, 5, 6

Wound healing

active protein fraction, 393
chymopapains, 393–394
description, 393
green papaya skin, 393
papain, 393, 394
proteolytic enzymes, 394
skin grafting, 393

X

X chromosomes

BACs (*see* Bacterial artificial
chromosomes (BACs))
carpel and stamen abortions, 317

X chromosomes (*cont.*)

- collinear region and embryo abortion, 318
- gene content
 - coding sequences, 322
 - deleterious retrotransposons, 323
 - expansion and rearrangements, 318, 319
 - human Y chromosome, 322
 - Muller's ratchet, 322
 - paired genes, 322
 - palindromic sequences, 322
 - pseudogenes, 322
 - and trafficking, 323
 - Vasconcellea monoica*, 323
 - Y-specific genes, 323
- hermaphrodite flowers, 317
- Meullers ratchet, 318–319
- minimum tiling path, 338
- physical mapping, 334–335
- recombinational suppression and
 - sex-specific regions, 317
- reiterative chromosome walking of, 336
- sequence gaps, 333

- sex-specific SCAR markers, 324
- structure and expansion, 316, 317
- trafficking and gene content, 323–324
- and Y, 331
- Y and Y^h chromosome, 317
- Y^h specification, 336

Y

- Y chromosome degeneration
 - ancestral autosome, 322
 - carpel abortion and stamen development, 317
 - chimpanzee, 312
 - hermaphrodites and males, 310
 - Meullers ratchet, 319
 - mutant gene, 318
 - Selene, 322
 - and X, 309, 311

Z

- Zapote Nodriza, chichihualzapotl, 8