

Index

- Absorption and fate
 - of PCDD/Fs in humans, 126
- Accidental and occupational exposure
 - relative to background exposure, 106
- Accidental and uncontrolled releases of POPs, 10
- Air
 - contamination in Central and Eastern Europe, 240
- Air drying
 - potential influence on concentrations in archived samples, 200
- Air to milk transfer coefficients, 81
- Air-side mass transfer coefficient
 - influence on air-water exchange, 40
- Air-to-foliage transfer
 - equilibrium partitioning versus kinetically-limited gaseous deposition, 64
 - role of wet and dry deposition, 67
- Air-water exchange
 - theoretical treatments of, 37
- Animal products
 - contribution to human exposure, 79
- Arctic contamination
 - with organochlorines, 215
 - with PAH, 214
- Arctic Monitoring Assessment Programme, 213
- Atmosphere
 - as a source to polar regions, 213
- Atmospheric half-lives, 45
- Atmospheric reactivity, 45

- Bioaccumulation
 - definition of, 148
- Bioaccumulation factor
 - definition of, 148
 - influence of organism on, 151
- Bioavailability, 149
- Bioconcentration
 - definition of, 146
- Bioconcentration factor
 - definition of, 147
 - in terrestrial food animals, 87
- Biomagnification
 - definition of, 148
- Biomagnification factor
 - definition of, 148
- Biota-Sediment-Accumulation Factor (BSAF) 150
- Biotransfer Factors
 - in terrestrial food animals, 87
- Bioturbation of surface sediment
 - influence on temporal resolution, 172

Carry-Over Ratios

- in terrestrial food animals, 87
- of PCDD/Fs from diet to milk, 91

Clearance rates

- in terrestrial food animals, 93

Clausius-Clapeyron equation

- utility in distinguishing between long- and short-range transport, 31, 218

Cold condensation, 220

Diet-to-milk transfer coefficients

- of 1,2,3,6,7,8-HxCDD in cattle, 89

Dry deposition

- relevance to atmospheric behaviour, 43

Elimination in terrestrial food animals

- discussion of, 92

Emission inventories, 4

Environmental budgets, 15

Episodic transport of POPs to polar regions, 222

Factors influencing human exposure assessments, 117

Feeding and management practices

- influence on exposure of food animals, 83

Fish consumption

- significance as a human exposure pathway, 122

Food-chain bioaccumulation, 149

Food-chain multiplier

- definition of, 149

Fugacity models

- use in constructing source inventories and environmental budgets, 18
- to describe fate in food animals, 85
- of fate in cattle, 99
- of aquatic food chain transfer, 155

Gas-particle partitioning

- factors affecting, 32

Global fractionation, 219

- evidence for, 221

Half-lives

- atmospheric, 45
- of PCDD/Fs in lactating dairy cows, 94

Henry's Law Constant

- definition of, 38
- influence on air-water exchange, 39
- measurement of, 39
- temperature-dependence of, 39

INDEX**Human adipose tissue**

- Czech and Slovak - POPs concentrations in, 258
- Polish - POPs concentrations in, 256

Human blood serum

- Slovak - POPs concentrations in, 254, 255

Human exposure

- Canada, 109
- Central and Eastern Europe, 252
- Denmark, 109
- dependence on age, 125
- Finland, 110
- Germany, 110
- Italy, 111
- Japan, 111
- New Zealand, 112
- Norway, 113
- of indigenous Arctic peoples, 212
- Spain, 113
- Sweden, 113
- The Netherlands, 111
- United Kingdom, 114
- United States, 114

Human milk

- comparison of contamination in different countries, 121
- concentrations in and influence on infant exposure, 118
- contamination of in Central and Eastern Europe, 253
- contamination of in Inuit women, 212
- temporal trends in contamination of, 120

 K_{ow}

- as a predictor of bioaccumulation in terrestrial food animals, 85, 98
- definition of, 146
- relationship to aquatic BCFs, 152
- relationship to uptake by root crops, 57

 K_{OA}

- as a predictor of foliar uptake, 64
- as a predictor of gas-particle partitioning, 36

Long-range transport

- as a source of PCBs to the Arctic, 217

Marine environments

- contamination in Central and Eastern Europe, 243

Mass balance calculations

- comparison of emissions and deposition, 15

Measuring BCFs and BAFs

- empirical methods, 150
- semi-empirical methods, 152
- Meteorology
 - in the Arctic, 213
- Michalovce District, 246, 247, 254, 255
 - PCB concentrations in blood serum of inhabitants of, 255
- Natural sources
 - of PCDD/Fs - significance of relative to anthropogenic sources, 199
- Non-thermal sources
 - of PCDD/Fs, 9
- OCDD
 - as a marker of PCP-associated releases of PCDD/Fs, 10
- Octanol
 - unsuitability as a surrogate for plant lipid, 64
- Organochlorine pesticides
 - "old" versus "new" sources, 225
- Pharmacokinetic models
 - to describe fate in food animals, 83
- Physiologically based models
 - to describe fate in food animals, 85
- Physiologically Based Pharmacokinetic (PBPK) modeling
 - to describe fate and distribution in humans, 131
- Point sources
 - impact on human exposure, 123
- POPs Listed under the UN/ECE (LRTAP) and UNEP Initiatives, 2
- POPs contamination in polar regions
 - history of, 212
- Relationship between concentrations in body fat and milk fat in cattle, 97
- Root concentration factor, 57
- Scavenging, 47
- Soil
 - ingestion of as a source of exposure to food animals, 82
 - contamination of in Central and Eastern Europe, 246
- Sources
 - of OCPs in Central and Eastern Europe, 237
 - of PAH in Central and Eastern Europe, 236
 - of PCBs in Central and Eastern Europe, 239
 - of PCDD/Fs in Central and Eastern Europe, 239
- Source identification

INDEX

- using multi-media fate models, 14
 - via pattern recognition, 12
 - via use of enantiomeric ratios, 11
- Source inventories, 4
- Sources and pathways of animal exposure, 80
- Sources of elevated concentrations in animal feed, 82
- Structural influences
- on uptake in terrestrial food animals, 97
 - Sub-cooled liquid vapour pressure
 - as a predictor of gas-particle partitioning, 33
- Techniques for assessing temporal trends
- archived samples, 193
 - biota samples, 198
 - dated peat cores, 190
 - glacial ice cores, 192
 - overview, 171
 - sediment cores, 172
- Temperature controlled air-surface exchange, 217
- Temporal trends
- of PCBs and PCDD/Fs - overview, 198
 - of PCBs in archived air filter samples, 197
 - of PCBs in archived vegetation samples, 195
 - of PCBs in archived soil samples, 194
 - of PCBs in European sediment cores, 180
 - of PCBs in North American sediment cores, 177
 - of PCBs in sediment cores - rest of world, 183
 - of PCBs in sediment cores from the Great Lakes, 173
 - of PCBs in water samples, 197
 - of PCDD/Fs in archived vegetation samples, 195
 - of PCDD/Fs in archived sewage sludge samples, 196
 - of PCDD/Fs in archived soil samples, 194
 - of PCDD/Fs in European sediment cores, 187
 - of PCDD/Fs in North American sediment cores, 186
 - of PCDD/Fs in sediment cores - rest of world, 190
 - of PCDD/Fs in sediment cores from the Great Lakes, 184
- Theoretical models
- of uptake by aquatic organisms, 154
- Toxicokinetics
- of PCDD/Fs in humans, 126
- Transfer coefficients
- in terrestrial food animals, 87
- Transfer into above-ground vegetation
- air-to-foliage transfer, 62
 - soil-to-foliage transfer, 60
- Transfer into chicken eggs, 93
- Transfer into root crops, 56

Vegetation

- as a passive monitor of atmospheric contamination, 73
- as a source of human exposure to PAH, 55
- as a source of human exposure to PCBs, 54
- as a source of human exposure to PCDD/Fs, 54
- as an atmospheric removal pathway, 71

Volatilisation

- relationship with temperature, 31

Washout ratio, 47**Water-side mass transfer coefficient**

- influence on air-water exchange, 41

Wet precipitation

- influence on atmospheric behaviour, 46

Wildlife

- contamination of in Central and Eastern Europe, 248

UN/ECE Convention on Long Range Transboundary Air Pollution, 1**UNEP and UN/ECE Protocols**

- Criteria for adding POPs to, 2

Uptake by root crops

- theoretical treatments of, 56
- validity of equilibrium partitioning approach, 57

Uptake by terrestrial food animals

- relative significance of different pathways, 80

Yu-Cheng, 124**Yusho, 124**