

# INDEX

- $\beta$ -Agonists, 7, 237  
Alkaline protease, 398, 401, 409  
Ascorbate, 102–104, 196–197  
Ascorbic acid, 23, 118; *see also* Ascorbate
- Beef, 1, 84–86, 116, 119, 124, 128–129, 142–143, 205, 290–291, 293, 299, 301, 356, 361, 371  
Belly burst, 254–255  
Boar taint, 147  
Bovine plasma powder, 400, 410, 412–413
- Ca<sup>2+</sup> ATPase, 253, 256–258, 267, 273, 278–282, 311, 324, 369  
Calcium-activated tenderization, 7  
Calpain, 7, 229–230, 237–239, 241–245, 393, 398, 400–401, 409, 414  
    activity in postmortem muscle, 237–239, 243–244  
    autolysis, 238–239  
    effect on protein degradation in surimi, 401  
Calpastatin, 237, 239–245, 401, 409  
    activity in postmortem muscle, 243–244  
    beef tenderness, 244  
    deactivation, 244  
    inhibition of calpain, 241
- $\beta$ -Carotene, 20–21, 103, 200–201  
Cathepsins, 7, 255, 393, 414, 422  
    cathepsin C, 253, 256, 260–261, 409  
    cathepsins B, L, L-like, 393–404, 415, 422, 425  
    frozen storage stability, 399–400  
    gel-softening, 401–403  
    protein degradation, 400–401  
    cathepsins D and H, 394, 409, 422  
Chicken: *see* Poultry  
Collagen, 6–7, 9, 219–226, 378  
    content, 371  
    cross-linking, 220–226  
        3-hydroxylysylpyridinium, 221–222  
        ketosamine, 221–222  
Collagen (*cont.*)  
    cross-linking (*cont.*)  
        lysyl hydroxylase, 224  
        lysyl oxidase, 224  
        lysylpyridinoline, 222  
        putative mechanisms, 224  
    hydroxyproline, 220  
    meat tenderness, 223  
    posttranslational modification, 220  
    types, 224–225  
Collagenase, 255–256, 260  
Color, 2–4, 201, 212, 255, 366–367; *see also* Myoglobin  
Conjugated dienes, 271–272  
Conjugated linoleic acid (CLA), 61–70  
    in fish muscle tissue, 64, 67  
    in fish organs, 66, 68, 70  
Connective tissue, 6, 219–220, 360  
    endomysium, 219  
    epimysium, 219  
    perimysium, 219  
Controlled atmosphere packaging (CAP), 74  
Cystatin, 393
- Dark-firm-dry (DFD), 3, 311, 313, 358  
DNA hybridization, 289
- Emulsification, 9–10, 321, 331–332  
Emulsions, 10, 332, 353–362  
    emulsion capacity test, 355, 357  
    emulsion stability, 10, 199, 328, 353, 355  
        mechanism, 354  
    emulsion stability test, 355–357  
    optical sensors to assess meat emulsion stability, 358–362  
Enzyme-linked immunosorbent assay (ELISA), 287–303  
    detection of species adulteration, 295–296  
    for species identification, 288–292, 295  
    for end temperature determination, 292–295

- Fatty acids, 16, 63, 116, 151, 269  
 arachidonic, 17  
 linoleic, 17, 127  
 linolenic, 17, 200  
 4-methylnonanoic, 17, 151–152  
 4-methyloctanoic, 17, 151–152, 154, 156  
 n-3 fatty acids, 17, 19; *see also* Omega-3 fatty acids  
 oleic, 17–18, 106  
 Omega-3 fatty acids, 69  
 palmitic, 17  
 stearic, 17
- Fish, 1–5, 116, 211, 213, 216, 278, 371, 373, 376, 394–395, 421–426
- Flavors, 4–6, 115–130, 192–193, 197, 211–212, 214  
 cured meat, 192  
 grassy flavor, 5, 137, 216  
 headspace volatiles, 13, 148–149  
 meaty flavor, 116–120, 162, 165–171  
 natural meat flavor, 4  
 ovine flavor, 135–156  
 effect of castration, 147–156  
 effect of lipid oxidation, 139–140  
 3-methylindole, 19, 147–151, 153, 156  
 reduction, 140–143  
 volatiles, 148–151  
 pastoral flavors, 18  
 phenols and indoles, 19  
 process-generated, 5  
 skatole, 19  
 species flavor, 128  
 volatile components in chicken and beef broths, 117, 121–122  
 warmed-over flavor (WOF), 6, 139–140, 192
- Gelation, 7, 278–279, 321, 327–331, 337–339, 354, 370–383, 395–396, 421–426  
 mechanism, process, 328, 422  
 mixed gels, 380–381
- $\beta$ -Glucan, 348–349
- Glutathione peroxidase, 23
- Glycogen, 14–15, 56, 254, 366–367
- Guanosine monophosphate (GMP), 116, 159
- Hemoglobin, 2,3
- High pressure treatment, 253–267
- Hydrodyne, 7
- Hydrophobicity, 373, 381–382
- Hydroxy radical, 270
- Inosine monophosphate (IMP), 4, 116–118, 159–171
- Interfacial protein film, 354
- Lactate, 12, 104, 254; *see also* Lactic acid
- Lactic acid, 54, 230
- Lamb, 1, 4, 128–129, 135, 142–143, 205; *see also* Mutton
- Least-cost-formulation (LCF), 355, 357
- Lipid oxidation, 75, 123, 139–140, 192, 197, 199–200, 212, 269–270, 272, 277–278, 283–284
- Lipoxygenase, 254
- Lutein, 20
- Mackerel, 211, 254, 397, 400  
 consumer perceptions, 211  
 cooked aromas, 215–216  
 histamine formation, 215
- $\alpha_2$ -Macroglobulin, 264–267
- Maillard reactions, 116–117, 122, 127, 129–130, 161, 167, 192, 212
- Malonaldehyde (MDA), 39, 277–284
- Meat particle binding, 327
- Modori, 372, 394, 396–399, 408–409
- Monoclonal antibodies, 287–288, 292–298, 300, 303
- Monosodium glutamate (MSG), 116, 159
- Muscle polymorphism, 365–385  
 fiber types, 3, 368–369  
 factors influencing fiber types, 369  
 muscle types, 3, 223, 366–368, 378  
 myosin isozymes (isoforms), 323, 368–369, 382–383
- Mutton, 129, 135, 140, 142–143
- Myofibrillar fragmentation index, 231, 243
- Myofibrillar proteins, 7, 9, 277, 284, 354, 365, 370, 372–373, 375, 378, 380–383, 396, 421–422  
 actin, 9, 12, 311, 323, 354, 376  
 $\alpha$ -actinin, 7  
 C-protein, 323, 377  
 desmin, 7, 236–237  
 filamin, 235–236  
 myosin, 9, 12, 253, 277–284, 311, 323, 354, 373, 376–377, 380, 383–384, 396, 416, 422, 424–425  
 nebulin, 7, 232–234  
 titin, 7, 231–232  
 tropomyosin, 7  
 troponin-T, 7, 234–235
- Myofibrils, 230–237, 310, 319–333, 373  
 hydration, 8, 321–322
- M-line, 230–231, 323
- myofilament lattice, 310, 323
- $N_1$ -line, 231
- shrinkage, 310, 323
- swelling, 8, 310, 319, 323, 326
- Z-line, 230–233, 235–236, 324
- Myoglobin, 2, 3, 31, 75, 204, 366  
 bloomed color, 76  
 content, 367  
 deoxymyoglobin, 75, 83  
 metmyoglobin, 3, 31–37, 73, 81, 106, 200  
 nitrosylmyoglobin, 200, 204  
 oxymyoglobin, 3, 31–34, 73, 86, 98
- N*-nitrosamine, 196
- Nitrite/nitrate, 3, 139–140, 191–205
- Nitrite-free curing, 191, 196–201
- Ohmic heating, 421–424

- Packaging  
air-permeable packaging, 77  
case-ready packaging, 76–77  
modified atmosphere packaging (MAP), 73–88  
  active gas exchange, 82  
  passive gas exchange, 81  
MAP with gaseous environments, 80–81  
MAP with pasteurizing gases  
vacuum packaging, 79–80, 82  
vacuum skin packaging, 78, 82
- Pale-soft-exudative (PSE), 3, 8, 311–315, 358
- Phosphates, 101, 319–333  
antioxidative effect, 101–102  
hexametaphosphate, 319–332  
hydrolysis, 101–102  
influence on myofibrillar protein functionality,  
  327–332  
phosphate-induced changes in myofibrils, 322–327  
phosphate-induced hydration of meat, 321–323  
protein extraction, 322–327  
pyrophosphate, 197, 319–332  
tripolyphosphate, 101–102, 197, 201–202, 319–332
- Plasmolysis, 95
- Polyclonal antibodies, 288, 291, 300
- Pork, 1, 126–129, 199, 205, 290–291, 293, 297–298,  
  313, 356, 362, 368, 370–371, 374
- Poultry, 1, 116, 291, 365–366, 368, 371, 383  
chicken, 2, 8, 40, 119, 124, 127–129, 140, 205,  
  290–291, 293, 296, 299, 302–303, 322–326,  
  370, 373–374, 376–377, 380–383  
duck, 1, 293, 370  
goose, 2  
turkey, 2, 8, 40, 293, 358, 370–371
- Pressure treatment, 253–267
- Protein oxidation, 270, 274
- Pyrazines, 5, 115–117, 122–124, 126, 149, 159, 203
- Red-soft-exudative (RSE), 313–314
- Salmon, 45–56  
  high energy diet, 45–48
- Sarcoplasmic proteins, 309, 311–313
- Sarcoplasmic reticulum, 269–274
- Selenium, 3, 23
- Sheep, 141, 237, 290–291, 293, 367–368
- Singlet oxygen, 85
- Sodium chloride (NaCl), 8, 96–100, 103, 199,  
  321–332, 381–382, 399, 402  
antioxidative effect, 98–99, 199  
extraction of myofibrillar proteins, 322–327  
preserving effect, 197  
prooxidative effect, 96–98, 199
- Somatotropin, 7
- Strecker degradation, 116, 122, 164, 212, 216
- Superoxide anion, 85–86, 270
- Surimi, 9, 140, 279, 284, 370–371, 385, 394–403,  
  422–423  
  beef heart surimi, 407–417  
  effect of endogenous enzymes, 398–403  
  gelation, 395–396, 409–414  
  gel-weakening (softening), 401–402, 411–414
- Suwari, 395–396, 408–409
- Tenderization, 229–245  
  postmortem changes in muscle tissue, 230–231  
  proteolysis, 243–244  
  30 kDa polypeptide/component, 7, 244
- Texture softening, 254
- Thiamine, 117, 173  
  degradation, 117–118, 175–189  
  oxidation, 173–189
- Thiobarbituric acid-reactive substances (TBARS),  
  39–41, 96–102,  
  104–109, 139, 198–199, 254, 269, 271
- $\alpha$ -Tocopherol, 3, 22, 31, 34–42, 196; *see also* Vitamin  
  E  
  antioxidant activity, 34–41  
  desirable levels, 36–38
- Transglutaminase, 372, 400, 421, 425–426
- Umami, 115–116, 159
- Vitamin E, 31–42  
  antioxidant activity, 34–35  
  dietary supplement, 36–42
- Water-holding capacity (WHC), 95, 213, 217, 278,  
  309–316, 320–321, 327–327, 335–349, 354,  
  365, 396, 413  
  in low-fat meat batters, 344–349  
  influence of meat batter microstructure, 344–345  
  measurement, 339–343