

INDEX

- ACE inhibitors, *see also* Angiotensin converting enzyme
angiotensin (1–7) and, 233
cardiac overload and, 23, 60
myocardial fibrosis and, 61
myocardial infarction and, 23, 30
- Acholeplasma laidlawii, 238–239
- ACTH, 9, 105, 109, 112
- Adenylate cyclase
AT₁ receptors and, 11, 40, 123, 217
AT₂ receptors and, 148
- Adipose tissue, AT₁ receptors in, 39–47
- Adrenal glands
AT₁ receptors in
antagonists, 135, 136
antisense injections and, 89
with hypertension, 49–58
subtypes, 8, 9, 61
hypothalamic AT₁ receptors and, 110–111
- Adrenal glomerulosa cells, desensitization in, 65
- Adrenergic neurons, 94, 96
- Aggregation of receptors, 65–66
- Aldosterone, 20, 56, 60
- Amphibian receptors, 239–240, 242
- Angiogenesis
AT₂ receptors and, 172
avian receptors and, 240
- Angiotensin I, 45, 238
- Angiotensin II, *see also* Renin-angiotensin system
behavioural effects, 96, 98
in carbohydrate-induced hypertension, 49–58
cardiovascular effects, 59–60, 226–227, 237
drinking response, 86, 94–96, 97–98, 113–119
fluid balance effects, 11, 199
growth promoting effects, 219
hypothalamic-pituitary-adrenal effects, 105–106
as a neurotransmitter, 113
peptide analogues, 134–141
tissues expressed in, 1
- Angiotensin III
brain activity, 114
receptor affinities for, 45, 238, 253
- Angiotensin IV, 233, 242, 250, 253–263
- Angiotensin (1–7), 242–243
kidney binding, 231
medulla oblongata activity, 225–235
- Angiotensin (2–8), *see* Angiotensin III
- Angiotensin (3–7), 233
- Angiotensin (3–8), 233, 242, 250, 253–263
- Angiotensin (4–8), 260
- Angiotensin converting enzyme, *see also* ACE inhibitors
in cardiac hypertrophy, 23
substrates for, 199
- Angiotensin receptors, *see* AT₁ receptors; AT₂ receptors; Atypical receptors
- Angiotensinogen, 199; *See also* Renin-angiotensin system
in adipose tissue, 40
antisense oligonucleotide for, 82, 84, 85–86
in brain, 121, 127
in heart, 23, 60
hypertension and, 85–86, 90
regulation by angiotensin II, 127
- Antagonists
of AT₁ receptor, 131–143, 237–238
of AT₂ receptor, 237–238
- Antisense oligonucleotides
for angiotensinogen, 82, 84, 85–86
for AT₁ receptor, 82, 85–89
controls, 89
designing of, 79–82
effectiveness, 84–85
for inhibiting hypertension, 79, 85–90
mechanisms of action, 81, 83–84
specificity, 84
stability, 83
viral vectors, 89–90
- Aorta, AT₁ receptors in
for assay of angiotensin analogues, 136–137
with carbohydrate-induced hypertension, 49–58
- AP-1 binding sequences
of AT₁ gene, 46
of AT₂ gene, 151, 222

- Apoptosis, AT₂ receptors and
in ovary, 176, 180–181
in vascular smooth muscle, 219
- Arachidonic acid, 149
- Astrocytes, *see also* Glial cells
angiotensin II receptors in, 121–129
angiotensin (3–8) receptors in, 254–262
angiotensin(1–7)-stimulated prostaglandin release, 243
- AT₁ receptors
in adipose tissue, 39–47
amino acid sequence, 146
antagonists, 131–143, 237–238
antisense oligonucleotide for, 82, 85–89
of astrocytes, 121–129
brain distribution, 9, 93–94
with carbohydrate-induced hypertension, 49–58
catecholamine neurons and, 94, 99–100
desensitization, 63–66
gene regulation
after myocardial infarction, 23–32
cis-regulatory elements, 1–10, 46
human, 18–20
in spontaneously hypertensive rats, 71–78
with STAT proteins, 62–63
- genome organization, 14–18
of glial cells, 1–10, 108
in heart, 59–69
after myocardial infarction, 23–32
with hypertension, 49–58
- heterogeneity
human, 12, 14–16, 20
rat (*see* AT_{1a} receptors; AT_{1b} receptors)
- human, 11–21
in hypothalamus, 49–58, 88, 93, 106–108
internalization, 66–67
of intestinal epithelial cells, 241
in kidney, 33–37
in medulla oblongata, 94, 230–232
of neuroblastoma cells, 200
of ovarian granulosa cells, 178
signal transduction pathways, 11, 40, 59–60, 62–66, 67, 217
in astrocytes, 123, 125, 126
c-fos in, 76
- sodium intake and, 33–37
in subfornical organ, 113, 116
tissue distribution, 8–9, 15
- AT_{1a} receptors, 11–12
amino acid sequence, 146
in brain, 37, 88
desensitization, 63–66
gene regulation
after myocardial infarction, 24, 26, 28, 30–31
cis-regulatory elements, 1–10
in heart, 59–69
internalization, 66–67
in kidney, 33–37
signal transduction pathways, 62–66, 67
tissue distribution, 8
- AT_{1b} receptors, 11–12
in adrenal gland, 8, 9, 61
in brain, 37, 88
gene regulation
after myocardial infarction, 24, 26, 28, 30–31
promoter sequence, 9
- in heart, 60
in kidney, 33–37
in pituitary, 9
signal transduction pathways, 66, 67
tissue distribution, 8–9
- AT₂ receptors
amino acid sequence, 158
angiotensin III binding, 45
angiotensin (1–7) binding, 232, 233
antagonists, 237–238
of astrocytes, 123–125, 127–128
brain distribution, 93, 94
cloning of, 205–206, 218
human, 153–158
in development, 40, 61, 153, 171–172, 189
of brain, 127, 128, 153, 247
of kidney, 163–164, 191
in drinking response, 116–117
gene cloning, 145–146, 162–163, 168–169
gene regulation, 150–151, 222–223
after myocardial infarction, 24, 26–31
human, 163
in growth, 40, 61, 171–172
of vascular smooth muscle, 149, 172, 218–222, 237
- in heart, 60–61
hemodynamic effects, 149, 247
heterogeneity, 172
in murine neuroblastoma cells, 149, 200–208
in rat brain, 172, 247
in rat kidney and fetus, 189–197
homology with AT_{1a}, 23–24
human, 153–165
in hypothalamus, 148, 203
internalization not observed, 66
in medulla oblongata, 230–232
of neuroblastoma cells, 149, 167–173, 199–208
in neuronal function, 40
in ovaries, 183–187
granulosa cells, 175–182, 183–184
signal transduction pathways, 61, 148–149, 155, 209–215, 217, 221–222
structure, 147, 170
tissue distribution, 31, 149–150, 153, 161–162, 163, 175, 189, 217
- AT₃ receptors, 241
- AT₄ receptors, 242; *See also* Angiotensin (3–8)
- Atypical receptors, 134, 237–245
avian, 131, 240
for fragments of angiotensin II, 233, 242–243, 253–263
in gerbil brain, 247–252
mammalian, 240–242
of *Mycoplasma*, 238–239
of *Xenopus*, 239–240

- Avian receptors, 131, 240
- Baroreceptor reflexes, 225–227, 228, 232
drinking response and, 117
- Biphenylimidazole antagonists, 237
- Blood vessels
cerebral
angiotensin (3–8) and, 242
 AT_2 receptors in, 247, 248
endothelial cell receptors in, 121
repair of, 149, 172, 219–222, 237
- Bp3*, 218
- Brain
angiotensin (3–8) receptors, 242, 253–263
 AT_1 receptors
of astrocytes, 121–129
catecholamines and, 94, 99–100
distribution, 9, 93–94
gene regulation, 7
of glial cells, 1–10, 108
hypertension and, 71–78
in hypothalamus, 49–58, 88, 93, 106–112
in medulla oblongata, 230–232
of neuroblastoma cells, 200
sodium intake and, 37
in subfornical organ, 113, 116
- AT_2 receptors
of astrocytes, 123–125, 127–128
in cerebellum, 203, 251
in cerebral arteries, 248
in development, 127, 128, 153, 247
distribution, 93, 94, 202–203
heterogeneity, 149, 172, 200–208, 247
in hypothalamus, 148, 203
in medulla oblongata, 230–232
of neuroblastoma cells, 149, 167–173, 199–208
PP2A coupling to, 209–215
- atypical receptors of gerbil, 247–252
drinking response and, 113–119
renin-angiotensin system in, 121
catecholamines and, 93–103
glucocorticoids and, 111
hypertension and, 71, 75–76, 85–86, 121
pineal, 96
- Brainstem, PP2A activity in, 209–210; *See also* Cerebellum; Medulla oblongata
- Calcium, intracellular, *see also* Phosphoinositide/calcium pathway
desensitization of transients, 63, 64
Xenopus receptors and, 239
- Calcium channels, 11, 148, 155, 181
- Carbohydrate-induced hypertension, 49–58
- Cardiac angiotensin receptors, 59–69
- Cardiac growth, 59–60
- Cardiac hypertrophy, 60, 61
angiotensin IV and, 262
gene regulation in, 23–24, 30–31
with hypertension, 49, 53, 55–57
- Cardiac renin-angiotensin system, 23, 60
- Catecholamines
avian angiotensin receptors and, 240
in brain, 93–103, 107–108
in brown adipose tissue, 40
drinking response and, 94–96, 97–98, 117
plasma levels during stress, 110
- Cerebellum
angiotensin (3–8) receptors, 253–254
 AT_1 receptors, on astrocytes, 124–125, 127
 AT_2 receptors, 203, 251
- Cerebral cortex, AT_1 receptors in, 124, 126, 127
- CGP42112A, 189, 237
- Chicken receptors, 240
- Cis-regulatory elements
of human AT_{1a} receptor gene, 9
of mouse AT_2 receptor gene, 151
of rat AT_{1a} receptor gene, 1–10
- Clathrin-coated pits, 66
- Constitutive receptor activity, 132–134
- Corpus luteum, 185–186
- Corticotropin releasing hormone, 105, 106, 107, 108, 110, 111
- Cyclic AMP, *see* Adenylate cyclase
- Cyclic GMP
angiotensin (3–8) receptors and, 242
 AT_2 receptors and, 148, 238
atypical neuroblastoma receptors and, 240–241
- Cytokines, STAT proteins and, 62–63
- Desensitization of receptors, 63–66
- Development, *see* AT_2 receptors, in development
- Diacylglycerol, 62
- Dimerization of receptors, 65–66
- Dithiothreitol
 AT_1 vs. AT_2 binding and, 178, 238
 AT_2 subtypes binding and, 204
- DOCA-salt model, 76
- Dopamine, 94, 97–101, 107
- Drinking response
antisense inhibition of, 86
brain regions involved in, 113–115
catecholamines and, 94–96, 97–98
cholinergic mechanisms, 114
fluid intake and, 115–116
hypotension and, 117
hypovolemia and, 117–118
osmotic stimuli, 116
receptor types in, 116–117
sodium appetite and, 117–118
- DuP753, *see* Losartan
- Endocytosis of receptors, 66–67
- Estrogen, 186
- EXP3174, 65
- FAK¹²⁵, 62

- Fibroblasts
 AT_{1a} receptors in, 24, 30, 61–62
 AT₂ receptors in, 150, 222–223
 signal transduction in, 62
- Fluid balance, *see* Drinking response; Osmotic stimuli
- Focal adhesion proteins, 62
- Follicle stimulating hormone (FSH), 176, 178–181, 185
- c-fos, angiotensin induction of, 61, 62
 in spontaneously hypertensive rats, 71–78
- Fos immunoreactivity, 113, 114
- Fructose, hypertension and, 49–58
- G proteins
 AT₁ receptors and, 11, 61, 63, 64, 217, 238
 AT₂ receptors and, 148, 217, 238
 with PP2A, 209–210, 212–213
 with PTPase, 31, 61, 155
 with receptor subtypes, 204
 desensitization of receptors and, 63
- Gene regulation, *see* Transcription factors
- Gene transfer *in vivo*, 218–220
- Glia cells, *see also* Astrocytes
 angiotensin (3–8) receptors in, 254–262
 AT₁ receptors in, 1–10, 108
- Glial fibrillary acidic protein (GFAP), 121
- Glucocorticoids
 AT_{1a} gene expression and, 30
 hypothalamic AT₁ receptors and, 108, 110–111, 112
- Growth
 AT₁ receptors and, 164
 AT₂ receptors and, 40, 61, 164, 171–172
 cardiac, 59–60
 of vascular smooth muscle, 149, 172, 218–222
- Growth factors
 in cardiac hypertrophy, 61
 in ovarian apoptosis, 180–181
 STAT proteins and, 62–63
- Growth hormone, 107
- Guanylate cyclase, angiotensin (3–8) receptors and, 242
- Heart failure, 60
- Hemagglutinating Virus of Japan (HVJ), 219
- High mobility group 1 protein (HMG-1), 219
- Hippocampus
 angiotensin (3–8) receptors in, 253
 atypical gerbil receptors in, 248–251
- Hyperinsulinemia, hypertension and, 49–58
- Hypertension
 angiotensin (1–7) and, 233
 animal models, 72, 76
 antisense oligonucleotides for, 79, 85–90
 carbohydrate-induced, 49–58
 spontaneously hypertensive rats, 71–78, 115, 218
- Hypertrophic marker genes, 61
- Hypertrophy, *see* Cardiac hypertrophy
- Hypothalamic-pituitary-adrenal axis, 105–112
- Hypothalamic-thalamus-septum, 76
- Hypothalamus
 angiotensin (1–7) in, 227–228, 232–233
 antisense treatment and, 87–89
 astrocytes in, 124, 126, 127
 AT₁ receptors in, 88, 93, 106–108
 with hypertension, 49–58, 73
 AT₂ receptors in, 148, 203
 dopamine in, 97, 99–100
 in drinking response, 96, 113–118
 noradrenaline in, 96–97
 PP2A activity in, 209–210
 in stress response, 105–112
- Immediate-early genes, 61
- Inferior olive, 203, 206, 251
- Inositol triphosphate, *see* Phosphoinositide/cal-cium pathway
- Insulin
 AT₂ gene expression and, 151
 with carbohydrate-enriched diet, 49–58
- Interleukin-1 β , 151
- Internalization
 of AT₁ receptors, 66–67
 of AT₂ receptors, 66, 148
- Intestinal epithelium, atypical receptors in, 241
- Intrinsic receptor activity, 132–134
- Inverse agonist, 131–134
- IRF proteins, 222
- Kidney
 AT₁ receptors in, 33–37
 AT₂ receptors in, 163–164, 190–196
- Locus coeruleus, 203, 206
- Losartan (DuP753)
 binding locus, 134
 brain penetration, 117
 receptor internalization and, 66
 specificity, 116, 237
- Luteinizing hormone, 184
- Macrophage receptors, 251
- MAP kinase
 AT_{1a} receptors and, 62, 65
 AT₂ receptors and, 221–222
 PP2A and, 215
- mas* oncoprotein, 241–242
- Medulla oblongata
 angiotensin (1–7) activity in, 227–235, 243
 astrocytes in, 124–125, 126, 127
 AT₁ receptors in, 73, 94, 230–232
- c-myc, 223
- Mycoplasma
 binding sites on, 238–239
 contamination from, 239, 241
- Myocardial fibrosis, 61

- Myocardial infarction, 23–32, 60, 61
Myocytes, cardiac
 hypertrophy, 60, 61
 receptor subtypes in, 23–24, 30–31
 signal transduction in, 62
- Neointimal proliferation, 149, 172, 219–222, 237
Nerve growth factor, 150
Neuroblastoma cells
 AT₁ receptors in, 200
 AT₂ receptors in, 149, 167–173, 199–208
 atypical receptors in, 240–241
- Neurons
 angiotensin (3–8) receptors in, 254–262
 AT₁ receptors, in hypertension, 40
 AT₂ receptors in, 40
- Nodose ganglion, 225–227, 228
- Non-AT₁, non-AT₂ receptors, *see also* Atypical receptors
Norepinephrine
 angiotensin II and, 94, 96–97, 107
 in adipose tissue, 45–46
 angiotensin (1–7) and, 233
- Norepinephrine transporter, 71–78
- Obesity, genetic, 39, 41, 43–44, 45–46
- Oncogenes
 fos induction by angiotensin, 61, 62, 71–78
 in hypertrophy, 61
 MAP kinase and, 62
 mas protein receptor, 241–242
 myc in AT₂ gene regulation, 223
- Osmotic stimuli, 108–109, 110, 116
- Ovaries
 AT₂ receptors in, 183–187
 granulosa cells, 175–182, 183–184
 follicular atresia, 176, 178–179, 181, 184
- Ovulation, 186
- Oxytocin
 AT₁ receptors and, 107, 108, 111
 AT₂ receptors and, 203
- PAI-1, 71–72, 75–76
- Paraventricular nucleus
 angiotensin (1–7) activity, 228, 232–233
 AT₁ receptor distribution, 106–108
 AT₂ receptor distribution, 203
 in drinking response, 113–118
 in stress response, 105–106, 108–112
- Paxillin, 62
- PC12 cells
 AT_{1a} gene regulation in, 1–10
 AT₂ receptors in, 148, 150, 181, 210
 properties of, 9
- PD123177, 237
- PD123319
 AT₂ subtypes and, 168, 204
 specificity for AT₂, 116, 237
- Pertussis toxin, 209
- Phenobarbital, AT₁ density and, 39–40, 41, 44, 46
- Phosphatidic acid, 62
Phosphoinositide/calcium pathway
 angiotensin (3–8) receptors and, 242
 AT₁ receptors and, 11, 40, 62, 217
 in astrocytes, 123, 125, 126
 AT₂ receptors and, 148
 desensitization of calcium transients, 63, 64, 65
- Phospholipase A₂, 126–127
- Phospholipase C, 11, 62, 123, 125–126; *See also*
 Phosphoinositide/calcium pathway
- Phospholipase D, 62
- Pineal gland, renin in, 96
- Pit-1 transcription factor, 18
- Pituitary, 105, 106
- Potassium currents, 148, 209, 210
- Potassium depletion, 56
- POU transcription factors, 18, 223
- Pregnancy
 AT₂ down-regulation in, 150
 ovarian RAS during, 183
- Pressor response, catecholamines and, 94, 97, 106, 107–108
- Pressure-overload model, 60
- Prolactin, 98, 107
- Prostaglandins
 angiotensin (1–7) and, 232, 243
 AT₁ receptors and
 in adipose tissue, 45
 in astrocytes, 123, 125–128
 AT₂ receptors and, 155, 167, 232
 atypical neuroblastoma receptors and, 240
- Protein kinase A, 63–64
- Protein kinase C
 AT₁ coupling to, 59, 62
 in AT₁ desensitization, 63–64
 phosphorylation sites on AT₂, 147
- Protein phosphatase 2A (PP2A), 209–215
- Protein tyrosine phosphatase (PTPase), 31, 61, 155, 176
 in adrenal cortex, 181
 in COS-7 cells, 210
 in neuroblastoma cells, 167, 170–172
 in NG108–15 cells, 149
 in PC12W cells, 148, 181, 210
 T-type calcium channels and, 181
- Receptors, *see also* AT₁ receptors; AT₂ receptors; Atypical receptors
- Renin, angiotensin (3–8) and, 242
- Renin-angiotensin system, *see also* Angiotensinogen
 in adipose tissue, 40
 in brain, 121
 catecholamines and, 93–103
 glucocorticoids and, 111
 hypertension and, 71, 75–76, 85–86, 121
 pineal, 96
 drinking response and, 117
 in heart, 23, 60

- Renin-angiotensin system (*cont*)**
- hypertension and, 49–50, 55–57, 71, 75–76, 85–86, 121
 - kidney in, 199
 - in ovary, 176, 183–185
 - sodium intake and, 37
- Renin-transgenic rat**, 72, 76
- Saralasin**, 238
- [$\text{Sar}^1, \text{Ile}^8$]-angiotensin II, 238
- Second messengers**, *see* Signal transduction
- Sendai virus**, 219
- Serine kinases**, 63–64
- Serine/threonine phosphatases**
- AT₂ receptor and, 148, 149, 176
 - PP2A, 209–215
- Serotonin**
- angiotensin binding and, 233
 - drinking response and, 117
- SHC proteins**, 62
- Shear-stress inducible transcription factor**, 19
- Signal transduction**
- with angiotensin (3–8) receptors, 242
 - with AT₁ receptors, 11, 40, 59–60, 62–66, 67, 217
 - in astrocytes, 123, 125, 126
 - c-fos in, 76
 - with AT₂ receptors, 61, 148–149, 209–215, 217, 221–222
 - with AT₂ receptors, 61, 148–149, 209–215, 217, 221–222
- sis* inducing factor, 63
- Sodium intake**
- AT₁ receptors and, 33–37
 - drinking response and, 117–118
- Solitary tract**, nucleus of, 93, 94
- Somatostatin**, 107
- Spontaneously hypertensive rat**
- AT₁ expression, 71, 72, 75–76
 - brain Fos expression, 115
- Src kinases**, 63
- STAT pathway**, 59, 62–63, 65
- Stress responses**, 105, 108–110, 111, 112
- Stroke**, gerbil model for, 248
- Stroke-prone spontaneously hypertensive rat**
- AT₁ expression in, 71–78
 - Bp3* gene in, 218
- Subfornical organ**
- in DOCA-salt model, 76
 - in drinking response, 113–118
- Substance P**, 226, 227, 228, 232
- Supraoptic nucleus**
- AT₂ receptors in, 203
 - drinking response and, 113–118
- Sympathetic nervous system**
- angiotensin activation of, 60, 94–97, 109–110
 - angiotensin inhibition of, 226–227
 - in hypertension, 50, 55
- T-type calcium channels**, 148, 155, 181
- TCV-116, 50**
- Testosterone**, ovarian receptors and, 185
- Tetrahydroimidazopyridine antagonists**, 237
- Thirst**, *see* Drinking response
- Transcription factors**
- for AT₁ gene, 18–20, 46
 - for AT_{1a} gene, 1–10
 - for AT₂ gene, 151, 163, 222–223
 - MAP kinase and, 62
 - STAT proteins, 62–63
- Turkey receptors**, 240
- Tyrosine dephosphorylation**, *see* Protein tyrosine phosphatase
- Tyrosine hydroxylase**, 233
- Tyrosine phosphorylation**
- AT_{1a} activation of, 59, 62, 67
 - of AT_{1a} receptors, 63–65
- Uterus**, AT₁ receptors in, 8
- Vagal reflexes**, 225–226
- Vascular smooth muscle**, repair of, 149, 172, 218–222, 237
- Vasopressin**
- angiotensin (1–7) and, 242
 - AT₁ receptors and, 9
 - AT₂ receptors and, 203
 - catecholamines and, 94, 96, 97, 106, 107–108, 109, 111
- Viral-mediated gene transfer**, 218–220
- X chromosome**, AT₂ gene on, 218
- Xenopus* receptors, 239–240, 242