

# References

- Adler, D., Murdoch, D. and others (2017). **rgl**: 3D visualization using OpenGL. R package version 0.98.22. <https://r-forge.r-project.org/projects/rgl/>.
- Akaike, H. (1973). Maximum likelihood identification of Gaussian autoregressive moving average models. *Biometrika*, **60**, 255–265.
- Albert, J. (2007). *Bayesian Computation with R*. New York: Springer.
- Al Kadiri, M., Carroll, R.J. and Wand, M.P. (2010). Marginal longitudinal semiparametric regression via penalized splines. *Statistics and Probability Letters*, **80**, 1242–1252.
- Bachrach, L.K., Hastie, T., Wang, M.-C., Narasimhan, B. and Marcus, R. (1999). Bone mineral acquisition in healthy Asian, Hispanic, Black and Caucasian youth. A longitudinal study. *Journal of Clinical Endocrinology and Metabolism*, **84**, 4702–12.
- Baltagi, B.H. (2013). *Econometric Analysis of Panel Data, Fifth Edition*. Chichester, U.K.: John Wiley & Sons.
- Bande, M.F. and de la Fuente, M.O. (2016). **fda.usc**: Functional data analysis and utilities for statistical computing. R package version 1.3. <http://www.jstatsoft.org/v51/i04/>.
- Berry, S.M., Carroll, R.J. and Ruppert, D. (2002). Bayesian smoothing and regression splines for measurement error problems. *Journal of the American Statistical Association*, **97**, 160–169.
- Biecek, P. (2014). **PBI**misc: A set of datasets used in my classes or in the book, *Modele liniowe i mieszane w R, wraz z przykladami w analizie danych*. R package version 0.999. <http://cran.r-project.org>.
- Bishop, C.M. (2006). *Pattern Recognition and Machine Learning*. New York: Springer.
- Bivand, R.S., Pebesma, E.J. and Gómez-Rubio, V. (2008). *Applied Spatial Data Analysis with R*. New York: Springer.
- Boddy, L., Wilkins, M.F. and Morris, C.W. (2001). Pattern recognition in flow cytometry. *Cytometry*, **44**, 195–209.
- de Boor, C. (2001). *A Practical Guide to Splines, Revised Edition*. New York: Springer-Verlag.
- Braglia, L. (2016). **aplore3**: Datasets from Hosmer, Lemeshow and Sturdivant, ‘Applied Logistic Regression’ (Third Edition, 2013). R package version 0.9. <http://www.r-project.org>.
- Breiman, L. and Friedman, J.H. (1985). Estimating optimal transformations for multiple regression and correlation. *Journal of the American Statistical Association*, **80**, 580–598.
- Breslow, N.E. and Clayton, D.G. (1993). Approximate inference in generalized linear mixed models. *Journal of the American Statistical Association*, **88**, 9–25.
- Brinkman, N.D. (1981). Ethanol fuel-A single-cylinder engine study of efficiency and exhaust emissions. *SAE Transactions*, **90**, 1410–1424.
- Brooks, S.P. and Gelman, A. (1998). General methods for monitoring convergence of iterative simulations. *Journal of Computational and Graphical Statistics*, **7**, 434–455.

- Brumback, B.A. and Rice, J.A. (1998). Smoothing spline models for the analysis of nested and crossed samples of curves (with discussion). *Journal of the American Statistical Association*, **93**, 961–994.
- Cardot, H. and Sarda, P. (2011). Functional linear regression. In *The Oxford Handbook of Functional Data Analysis*, editors Ferraty, F. and Romain, Y., Oxford: Oxford University Press, pp. 21–46.
- Carlin, B.P. and Louis, T.A. (2009). *Bayesian Methods for Data Analysis, Third Edition*. Boca Raton, Florida: Chapman & Hall/CRC.
- Carpenter, B., Gelman, A., Hoffman, M.D., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M., Guo, J., Li, P. and Riddell, A. (2017). Stan: A probabilistic programming language. *Journal of Statistical Software*, **76**, Issue 1, 1–32.
- Carroll, R.J., Roeder, K. and Wasserman, L. (1999). Flexible parametric measurement error models, *Biometrics*, **55**, 44–54.
- Carroll, R.J. and Ruppert, D. (1988). *Transformation and Weighting in Regression*. New York: Chapman & Hall.
- Carroll, R.J., Ruppert, D., Stefanski, L.A. and Crainiceanu, C.M. (2006). *Measurement Error in Nonlinear Models, Second Edition*. Boca Raton, Florida: Chapman & Hall/CRC.
- Chang, C.-C. and Lin, C.-J. (2011). **LIBSVM** : a library for support vector machines. *Association for Computing Machinery Transactions on Intelligent Systems and Technology*, **2**, 27:1–27:27.
- Chouldechova, A., and Hastie, T. (2015). Generalized additive model selection. Unpublished manuscript. <https://arxiv.org/pdf/1506.03850>
- Chouldechova, A., Hastie, T., and Spinu, V. (2018). **gamsel**: Fit regularization path for generalized additive models. R package version 1.8.1 <http://www.r-project.org>.
- Cole, T.J. and Green, P.J. (1992). Smoothing reference centile curves: the LMS method and penalized likelihood. *Statistics in Medicine*, **11**, 1305–1319.
- Coull, B.A., Ruppert, D. and Wand, M.P. (2001). Simple incorporation of interactions into additive models. *Biometrics*, **57**, 539–545.
- Crainiceanu, C. and Ruppert, D. (2004). Likelihood ratio tests in linear mixed models with one variance component. *Journal of the Royal Statistical Society, Series B*, **66**, 165–185.
- Crainiceanu, C.M., Ruppert, D., Claeskens, G. and Wand, M.P. (2005). Exact likelihood ratio tests for penalized splines. *Biometrika*, **92**, 91–103.
- Crainiceanu, C., Ruppert, D. and Wand, M.P. (2005). Bayesian analysis for penalized spline regression using WinBUGS. *Journal of Statistical Software*, **14**, Issue 14, 1–24.
- Craven, P. and Wahba, G. (1979). Smoothing noisy data with spline functions: estimating the correct degree of smoothing by the method of generalized cross-validation. *Numerische Mathematik*, **31**, 377–403.
- Cressie, N.A.C. (2015). *Statistics for Spatial Data, Revised Edition*. New York: Wiley.
- Cressie, N. and Wikle, C. (2011). *Statistics for Spatio-Temporal Data*. Hoboken, New Jersey: John Wiley & Sons.
- Creswell, M. (1991). A multilevel bivariate model. In *Data Analysis with ML3*, editors Prosser, R., Rasbash, J. and Goldstein, H., London: Institute of Education, London, pp. 76–108.
- Cristianini, N. and Shawe-Taylor, J. (2000). *An Introduction to Support Vector Machines and Other Kernel-Based Learning Methods*. Cambridge, U.K.: Cambridge University Press.
- Croissant, Y. (2016). **Ecdat**: Data sets for econometrics. R package version 0.3. <http://www.r-project.org>.
- Daniels, M.J. and Hogan, J.W. (2008). *Missing Data in Longitudinal Studies*. Boca Raton, Florida: Chapman & Hall/CRC.
- Deb, P. and Trivedi, P.K. (1997). Demand for medical care by the elderly: A finite mixture approach. *Journal of Applied Econometrics*, **12**, 313–336.
- Diggle, P., Heagerty, P., Liang, K.-L. and Zeger, S. (2002). *Analysis of Longitudinal Data, Second Edition*. Oxford, U.K.: Oxford University Press.
- Diggle, P.J. and P.J. Ribeiro Jr. (2007). *Model-based Geostatistics*. New York: Springer.
- Di Narzo, A.F. and Stigler, M. (2016). **tsDyn**: Nonlinear time series models with regime switching. R package version 0.9. <http://github.com/MatthieuStigler/tsDyn/wiki>.

- Donnelly, C.A., Laird, N.M. and Ware, J.H. (1995). Prediction and creation of smooth curves for temporally correlated longitudinal data. *Journal of the American Statistical Association*, **90**, 984–989.
- Duan, N. (1983). Smearing estimate: a nonparametric retransformation method. *Journal of the American Statistical Association*, **78**, 605–610.
- Dunn, P.K. and Smyth, G.K. (1996). Randomized quantile residuals. *Journal of Computational and Graphical Statistics*, **5**, 236–244.
- Durbán, M., Harezlak, J., Wand, M.P. and Carroll, R.J. (2005). Simple fitting of subject-specific curves for longitudinal data. *Statistics in Medicine*, **24**, 1153–1167.
- Eilers, P.H.C. and Marx, B.D. (1996). Flexible smoothing with B-splines and penalties (with discussion). *Statistical Science*, **11**, 89–121.
- Fahrmeir, L. and Kneib, T. (2011). *Bayesian Smoothing and Regression for Longitudinal, Spatial, and Event History Data*. Oxford, U.K.: Oxford University Press.
- Fahrmeir, L. and Tutz, G. (1994). *Multivariate Statistical Modelling Based on Generalized Linear Models*. New York: Springer-Verlag.
- Faraway, J.J. (2006). *Extending the Linear Model with R*. Boca Raton, Florida: Chapman & Hall/CRC.
- Ferraty, F. and Romain, Y. (2011). *The Oxford Handbook of Functional Data Analysis*. Oxford: Oxford University Press.
- Ferraty, F. and Vieu, P. (2006). *Nonparametric Functional Data Analysis*. New York: Springer.
- Fitzmaurice, G., Davidian, M., Verbeke, G. and Molenberghs, G. (Editors) (2008). *Longitudinal Data Analysis*. Boca Raton, Florida: Chapman & Hall/CRC.
- Fitzmaurice, G.M., Laird, N.M. and Ware, J.H. (2004). *Applied Longitudinal Analysis*. Hoboken, New Jersey: John Wiley & Sons.
- Frees, E.W. (2004). *Longitudinal and Panel Data: Analysis and Applications in the Social Sciences*. Cambridge, U.K.: Cambridge University Press.
- Fuller, W.A. (1987). *Measurement Error Models*. New York: John Wiley & Sons.
- Gałecki, A. and Burzykowski, T. (2013). *Linear and Mixed-Effects Models Using R*. New York: Springer.
- Gelman, A. (2006). Prior distributions for variance parameters in hierarchical models (Comment on article by Browne and Draper). *Bayesian Analysis*, **3**, 515–534.
- Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., and Rubin, D. B. (2014). *Bayesian Data Analysis, 3rd Ed.*, Chapman & Hall Ltd (London; New York)
- Gelman, A. and Hill, J. (2007). *Data Analysis using Regression and Multilevel/Hierarchical Models*. New York: Cambridge University Press.
- Gelman, A. and Rubin, D.B. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, **7**, 457–511.
- Gelman, A., Sturtz, S., Ligges, U., Gorjanc, G. and Kerman, J. (2015). **R2WinBUGS**: Running WinBUGS and OpenBUGS from R/S-PLUS. R package version 2.1. <http://www.r-project.org>.
- Goldstein, H. (2010). *Multilevel Statistical Models, Fourth Edition*. Chichester, U.K.: John Wiley & Sons.
- Goldsmith, J., Bobb, J., Crainiceanu, C., Caffo, B. and Reich, D. (2011). Penalized functional regression. *Journal of Computational and Graphical Statistics*, **20**, 830–851.
- Goldsmith, J., Scheipl, F., Huang, L., Wrobel, J., Gellar, J., Harezlak, J., McLean, M.W., Swihart, B., Xiao, L., Crainiceanu, C., Reiss, P., Chen, (2016). **refund**: Regression with functional data. R package version 0.1. <http://www.r-project.org>.
- Green, P.J. and Silverman, B.W. (1994). *Nonparametric Regression and Generalized Linear Models*. London: Chapman and Hall.
- Gu, C. (2017). **gss**: General smoothing splines. R package version 2.1. <http://www.r-project.org>.
- Guo, W. (2002). Functional mixed effects models. *Biometrics*, **58**, 121–128.
- Guo, J., Gabry, J. and Goodrich, B. (2017). **rstan**: R interface to Stan. R package version 2.17.2. <http://mc-stan.org>.

- Gurrin, L.C., Scurrah, K.J. and Hazelton, M.L. (2005). Tutorial in biostatistics: spline smoothing with linear mixed models. *Statistics in Medicine*, **24**, 3361–3381.
- Härdle, W., Hall, P. and Marron, J.S. (1988). How far are automatically chosen regression smoothing parameters from their optimum? *Journal of the American Statistical Association*, **83**, 86–101.
- Hastie, T. (1996). Pseudosplines. *Journal of the Royal Statistical Society, Series B*, **58**, 379–396.
- Hastie, T. J. (1992) Generalized additive models. Chapter 7 of *Statistical Models in S* eds J. M. Chambers and T. J. Hastie, Wadsworth & Brooks/Cole.
- Hastie, T. (2017a). *gam*: Generalized additive models. R package version 1.14. <http://www.r-project.org>.
- Hastie, T. (2017b). *mda*: Mixture and flexible discriminant analysis. R package version 0.4. <http://www.r-project.org>.
- Hastie, T. (2016). *svmpath*: The support vector machine path algorithm. R package version 0.955 <http://www.jmlr.org/papers/volume5/hastie04a/hastie04a.pdf>.
- Hastie, T. and Efron, E. (2013). *lars*: Least angle regression, lasso and forward stagewise. R package version 1.2. <http://www.r-project.org>.
- Hastie, T.J. and Tibshirani, R.J. (1986). Generalized additive models. *Statistical Science*, **1**, 297–310.
- Hastie, T.J. and Tibshirani, R.J. (1990). *Generalized Additive Models*. Boca Raton, Florida: Chapman & Hall/CRC.
- Hastie, T. and Tibshirani, R. (1993). Varying-coefficient models. *Journal of the Royal Statistical Society, Series B*, **55**, 757–796.
- Hastie, T. and Tibshirani, R. (2000). Bayesian backfitting. *Statistical Science*, **15**, 196–223.
- Hastie, T., Tibshirani, R. and Friedman, J. (2009). *The Elements of Statistical Learning, Second Edition*. New York: Springer.
- Hastie, T., Tibshirani, R. and Wainwright, M. (2015). *Statistical Learning with Sparsity*. Boca Raton, Florida: CRC Press.
- Hastie, T. and Zhu, J. (2006). Comment on article by J.M. Moguerza and A. Muñoz. *Statistical Science*, **21**, 352–357.
- Hodges, J.S. (2014). *Richly Parameterized Linear Models*. Boca Raton, Florida: Chapman & Hall/CRC.
- Hoff, P.D. (2010). *A First Course in Bayesian Statistical Methods*. New York: Springer.
- Hoffman, M.D. and Gelman, A. (2014). The No-U-turn Sampler: adaptively setting path lengths in Hamiltonian Monte Carlo. *Journal of Machine Learning Research*, **15**, 1593–1623.
- Horváth, L. and Kokoszka, P. (2012). *Inference for Functional Data with Applications*. New York: Springer.
- Huang, A. and Wand, M.P. (2013). Simple marginally noninformative prior distributions for covariance matrices. *Bayesian Analysis*, **8**, 439–452.
- Huber, P.J. (1981). *Robust Statistics*. New York: Wiley.
- Huq, N.M. and Cleland, J. (1990). *Bangladesh Fertility Survey 1989 (Main Report)*. Dhaka, Bangladesh: National Institute of Population Research and Training.
- Hurvich, C. M., Simonoff, J. S. and Tsai, C. (1998). Smoothing parameter selection in nonparametric regression using an improved Akaike information criterion. *Journal of the Royal Statistical Society, Series B*, **60**, 271–293.
- Ivanescu, A., Staicu, A.-M., Scheipl, F. and Greven, S. (2015). Penalized function-on-function regression. *Computational Statistics*, **30**, 539–568.
- James, G. (2011). Sparseness and functional data analysis. In *The Oxford Handbook of Functional Data Analysis*, editors Ferraty, F. and Romain, Y., Oxford: Oxford University Press, pp. 298–326.
- Jones, M.C. and Faddy, M.J. (2003). A skew extension of the *t*-distribution, with applications. *Journal of the Royal Statistical Society, Series B*, **65**, 159–174.
- Kammann, E.E. and Wand, M.P. (2003). Geoadditive models. *Journal of the Royal Statistical Society, Series C*, **52**, 1–18.

- Karatzoglou, A., Smola, A. and Hornik, K. (2016). **kernlab**: Kernel-based machine learning lab. R package version 0.9. <http://www.r-project.org>.
- Kauermann, G., Krivobokova, T. and Fahrmeir, L. (2009). Some asymptotic results on generalized penalized spline smoothing. *Journal of the Royal Statistical Society, Series B*, **71**, 487–503.
- Kauermann, G. and Opsomer, J.D. (2011). Data-driven selection of the spline dimension in penalized spline regression. *Biometrika*, **98**, 225–230.
- Kayal, M., Vercelloni, J., Wand, M.P. and Adjeroud, M. (2015). Searching for the best bet in life-strategy: a quantitative approach to individual performance and population dynamics in reef-building corals. *Ecological Complexity*, **23**, 73–84.
- Kipnis, V., Subar, A.F., Midthune, D., Freedman, L.S., Ballard-Barbash, R., Troiano, R., Bingham, S., Schoeller, D.A., Schatzkin, A. and Carroll, R.J. (2003). The structure of dietary measurement error: results of the OPEN biomarker study. *American Journal of Epidemiology*, **158**, 14–21.
- Kleiber, C. and Kleiber, A. (2017). **AER**: Applied econometrics with R. R package version 1.2. <http://cran.r-project.org>.
- Kneib, T., Heinzl, F., Brezger, A., Sabanes, B. and Klein, N. (2014). **BayesX**: R utilities accompanying the software package BayesX. R package version 0.2. <http://www.bayesx.org>.
- Koenker, R. (2005). *Quantile Regression*. New York: Cambridge University Press.
- Koenker, R., Ng, P. and Portnoy, S. (1994). Quantile smoothing splines. *Biometrika*, **81**, 673–68.
- Koenker, R. (2017). **quantreg**: Quantile regression. R package version 5.34. <http://www.r-project.org>.
- Kooperberg, C. (2015). **polyspline**: Polynomial spline routines. R package version 1.1. <http://www.r-project.org>.
- Kou, S.C. and Efron, B. (2002). Smoothers and the  $C_p$ , generalized maximum likelihood, and extended exponential criteria. *Journal of the American Statistical Association*, **97**, 766–782.
- Krivobokova, T. (2013). Smoothing parameter selection in two frameworks for penalized splines. *Journal of the Royal Statistical Society, Series B*, **75**, 725–741.
- Landman, B.A., Huang, A.J., Gifford, A., Vikram, D.S., Lim, I.A.L, Farrell, J.A.D., Bogovic, J.A., Hua, J., Chen, M., Jarso, S., Smith, S.A., Joel, S., Mori, S., Pekar, J.J., Barker, P.B., Prince, J.L. and van Zijl, P.C.M. (2010). Multi-parametric neuroimaging reproducibility: A 3T resource study. *NeuroImage*, **54**, 2854–2866.
- Lange, K.L., Little, R.J.A. and Taylor, J.M.G. (1989). Robust statistical modeling using the  $t$ -distribution. *Journal of the American Statistical Association*, **84**, 881–896.
- Lauritzen, S.L. (1996). *Graphical Models*. Oxford, U.K.: Clarendon Press.
- Lee, M.-J. (1995). Semi-parametric estimation of simultaneous equations with limited dependent variables: a case study of female labour supply. *Journal of Applied Econometrics*, **10**, 187–200.
- Lee, P.M. (2012). *Bayesian Statistics*. Chichester, U.K.: John Wiley & Sons.
- Leisch, F. and Dimitriadou, E. (2010). **mlbench**: Machine learning benchmark problems. R package version 2.1. <http://www.r-project.org>.
- Li, Y. and Ruppert, D. (2008). On the asymptotics of penalized splines. *Biometrika*, **95**, 415–436.
- Ligges, U., Sturtz, S., Gelman, A., Gorjanc, G. and Jackson, C. (2017). **BRugs**: Interface to the OpenBUGS Markov chain Monte Carlo software. R package version 0.9. <http://www.r-project.org>.
- Lin, Y. and Zhang, H. (2006). Component selection and smoothing in multivariate nonparametric regression. *Annals of Statistics*, **34**, 2272–2297.
- Little, R.J. and Rubin, D.B. (2002). *Statistical Analysis with Missing Data, Second Edition*. Hoboken, New Jersey: John Wiley & Sons.
- Loader, C. (1999). *Local Regression and Likelihood*. New York: Springer.
- Loader, C. (2013). **locfit**: Local regression, likelihood and density estimation. R package version 1.5. <http://www.r-project.org>.
- Longford, N.T. (2005). *Missing Data and Small-Area Estimation*. New York: Springer.
- Lunn, D., Jackson, C., Best, N., Thomas, A. and Spiegelhalter, D. (2013). *The BUGS Book*. Boca Raton, Florida: CRC Press.

- McLean, M., Hooker, G., Staicu, A.-M., Scheipl, F. and Ruppert, D. (2013). Functional generalized additive models. *Journal of Computational and Graphical Statistics*, **23**, 249–269.
- Marley, J.K. and Wand, M.P. (2010). Non-standard semiparametric regression via **BRugs**. *Journal of Statistical Software*, **37**, Issue 5, 1–30.
- Maronna, R.A., Martin, D.R. and Yohai, V.J. (2006). *Robust Statistics: Theory and Methods*. London: Wiley.
- Marra, G. and Wood, S.N. (2011). Practical variable selection for generalized additive models. *Computational Statistics and Data Analysis*, **55**, 2372–2387.
- McCullagh, P. and Nelder, J.A. (1989). *Generalized Linear Models, Second edition*. London: Chapman and Hall.
- McCulloch, C.E., Searle, S.R. and Neuhaus, J.M. (2008). *Generalized, Linear, and Mixed Models, Second Edition*. New York: John Wiley & Sons.
- Meyer, D., Dimitriadou, E., Hornik, K., Weingessel, A., Leisch, F., Chang, C.-C., Lin, C.-C. (2017). **e1071**: Miscellaneous functions of the department of statistics, probability theory group (formerly: E1071), TU Wien. **R** package version 1.6. <http://www.r-project.org>.
- Minka, T., Winn, J., Guiver, J., Webster, S., Zaykov, Y., Yangel, B., Spengler, A. and Bronskill, J. (2014). **Infer.NET 2.6**, Microsoft Research Cambridge. <http://research.microsoft.com/infernet>
- Moguerza, J.M. and Muñoz, A. (2006). Support vector machines with applications (with discussion). *Statistical Science*, **21**, 322–362.
- Müller, H.G., Wu, Y. and Yao, F. (2013). Continuously additive models for nonlinear functional regression. *Biometrika*, **100**, 607–622.
- Müller, H.G. and Yao, F. (2008). Functional additive models. *Journal of the American Statistical Association*, **103**, 1534–1544.
- Munnell, A.H., Tootell, G.M.B., Browne, L.E. and McEneaney, J. (1996). Mortgage lending in Boston: Interpreting HDMA data. *American Economic Review*, 25–53.
- Nason, G.P. (2008). *Wavelet Methods in Statistics with R*. New York: Springer.
- Nason, G. (2016). **wavethresh**: Wavelets statistics and transforms. **R** package version 4.6.8. <http://www.r-project.org>.
- Nychka, D., Furrer, R., Paige, J. and Sain, S. (2017). **fields**: Tools for spatial data. **R** package version 9.0. <http://www.r-project.org>.
- O’Connell, M.A. and Wolfinger, R.D. (1997). Spatial regression models, response surfaces, and process optimization. *Journal of Computational and Graphical Statistics*, **6**, 224–241.
- Ormerod, J.T. and Wand, M.P. (2018). **LowRankQP**: Low-rank quadratic programming. **R** package version 1.0.3. <http://www.r-project.org>.
- Ormerod, J.T., Wand, M.P. and Koch, I. (2008). Penalised spline support vector classifiers: computational issues. *Computational Statistics*, **23**, 623–641.
- O’Sullivan, F. (1986). A statistical perspective on ill-posed inverse problems (with discussion). *Statistical Science*, **1**, 505–527.
- Parker, R.L. and Rice, J.A. (1985). Comment on article by B.W. Silverman. *Journal of the Royal Statistical Society, Series B*, **47**, 40–42.
- Pearce, N.D. and Wand, M.P. (2006). Penalized splines and reproducing kernel methods. *The American Statistician*, **60**, 233–240.
- Pearce, N.D. and Wand, M.P. (2009). Explicit connections between longitudinal data analysis and kernel machines. *Electronic Journal of Statistics*, **3**, 797–823.
- Pearl, J. (1988). *Probabilistic Reasoning in Intelligent Systems*. San Francisco: Morgan Kaufmann Publishers.
- Pinheiro, J.C. and Bates, D.M. (2000). *Mixed-Effects Models in S and S-PLUS*. New York: Springer-Verlag.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., **EISPACK** authors and **R** Core Team. (2017). **nlme**: Linear and nonlinear mixed effects models. **R** package version 3.1. <http://www.r-project.org>.
- Pope, C.A., Dockery, D.W., Spengler, J.D. and Raizenne, M.E. (1991). Respiratory health and PM<sub>10</sub> pollution: a daily time series analysis. *American Review of Respiratory Disease*, **144**, 668–674.

- Pratt, J.H., Jones, J.J., Miller, J.Z., Wagner, M.A. and Fineberg, N.S. (1989). Racial differences in aldosterone excretion and plasma aldosterone concentrations in children. *New England Journal of Medicine*, **321**, 1152–1157.
- Pya, N. (2017). **sca**m: Shape constrained additive models. R package version 1.2-2. <http://cran.r-project.org>.
- Pya, N. and Wood, S.N. (2016). A note on basis dimension selection in generalized additive modelling. Unpublished manuscript. <http://arxiv.org/abs/1602.06696>
- Ramsay, J.O., Hooker, G. and Graves, S. (2009). *Functional Data Analysis with R and MATLAB*. New York: Springer.
- Ramsay, J.O. and Silverman, B.W. (2006). *Functional Data Analysis, Second Edition*. New York: Springer.
- Ramsay, J.O., Wickham, H., Graves, S. and Hooker, G. (2017). **fda**: Functional data analysis. R package version 2.4.7. <http://www.functionaldata.org>.
- Rasmussen, C.E. and Williams, C.K.I. (2006). *Gaussian Processes for Machine Learning*. Cambridge, Massachusetts: MIT Press.
- Rao, J.N.K. and Molina, I. (2015). *Small Area Estimation, Second Edition*. Hoboken, New Jersey: John Wiley & Sons.
- Ravikumar, P., Lafferty, J., Liu, H. and Wasserman, L. (2009). Sparse additive models. *Journal of the Royal Statistical Society, Series B*, **71**, 1009–1030.
- Reiss, P.T. and Ogden, R.T. (2009). Smoothing parameter selection for a class of semiparametric linear models. *Journal of the Royal Statistical Society, Series B*, **71**, 505–523.
- Ribeiro, P.J. and Diggle, P.J. (2016). **geoR**: Analysis of geostatistical data. R package version 1.7. <http://www.leg.ufpr.br/geoR>.
- Richardson, S., Leblond, L., Jausset, I. and Green, P.J. (2002). Mixture models in measurement error problems with reference to epidemiological studies. *Journal of the Royal Statistical Society, Series A*, **165**, 549–566.
- Rigby, R.A. and Stasinopoulos, D.M. (2005). Generalized additive models for location, scale and shape. *Journal of the Royal Statistical Society, Series C*, **54**, 507–554.
- Ripley, B., Venables, B., Bates, D.M., Hornik, K., Gebhardt, A. and Firth, D. (2015). **MASS**: Support functions and datasets Venables and Ripley's 'Modern Applied Statistics with S'. R package version 7.3. <http://www.stats.ox.ac.uk/pub/MASS4/>.
- Robinson, G.K. (1991). That BLUP is a good thing: the estimation of random effects. *Statistical Science*, **6**, 15–51.
- Roeder, K., Carroll, R.J. and Lindsay, B.G. (1996). A semiparametric mixture approach to case-control studies with errors in covariables. **91**, 722–732.
- Ruppert, D. (2002). Selecting the number of knots for penalized splines. *Journal of Computational and Graphical Statistics*, **11**, 735–757.
- Ruppert, D. and Matteson, D.S. (2015). *Statistics and Data Analysis for Financial Engineering, Second Edition*. New York: Springer.
- Ruppert, D., Wand, M.P. and Carroll, R.J. (2003). *Semiparametric Regression*. Cambridge, U.K.: Cambridge University Press.
- Ruppert, D., Wand, M.P. and Carroll, R.J. (2009). Semiparametric regression during 2003–2007. *Electronic Journal of Statistics*, **3**, 1193–1256
- Salibian-Barrera, M., Cubranic, D. and Alimadad, A. (2014). **rgam**: Robust generalized additive model. R package version 0.6. <http://www.stat.ubc.ca/~matias/rgam/>.
- Sarkar, D. (2008). *Lattice*. New York: Springer.
- Sarkar, D. (2017). **lattice**: Trellis graphics for R. R package version 0.20. <http://lattice.r-forge.r-project.org/>.
- Scheipl, F. and Bolker, B. (2016). **RLRsim**: Exact (restricted) likelihood ratio tests for mixed and additive models. R package version 3.1. <https://github.com/fabian-s/RLRsim>.
- Schölkopf, B. and Smola, A.J. (2002). *Learning with Kernels*. Cambridge, Massachusetts: MIT Press.
- Sigrist, M.W. (Editor) (1994). *Air Monitoring by Spectroscopic Techniques*. New York: Wiley.
- Sommer, A. (1982). *Nutritional Blindness*. New York: Oxford University Press.

- Stan Development Team. (2017). *Stan Modeling Language User's Guide and Reference Manual, Stan Version 2.17.0*. <http://mc-stan.org>.
- Stasinopoulos, D.M. and Rigby, R.A. (2008). Generalized additive models for location scale and shape (GAMLSS) in *R. Journal of Statistical Software*, **23**, Issue 7, 1–46.
- Stasinopoulos, M. and Rigby, B. (2017). **gamlss**: Generalised additive models for location scale and shape. *R* package version 5.0. <http://www.gamlss.org/>.
- Svensson, L.E.O. (1994). Estimating and interpreting forward interest rates: Sweden 1992–1994. International Monetary Fund Working Paper WP/94/114. Washington D.C.: International Monetary Fund Publication Services.
- Taylor, J. and Tibshirani, R.J. (2015). Statistical learning and selective inference. *Proceedings of the National Academy of Sciences of the United States of America*, **112**, 7629–7634.
- Therneau, T.M. and Lumley, T. (2015). **survival**: Survival analysis. *R* package version 2.38. <http://www.r-project.org>.
- Tibshirani, R. (1996). Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society, Series B, Methodological*, **58**, 267–288.
- Tibshirani, R., Tibshirani, R., Taylor, J., Loftus, J. and Reid, S. (2017). **selectiveInference**: Tools for post-selection inference. *R* package version 1.2.4. <http://cran.r-project.org>.
- Umlauf, N., Adler, D., Kneib, T., Lang, S. and Zeileis, A. (2015). Structured additive regression models: an *R* interface to **BayesX**. *Journal of Statistical Software*, **63**, Issue 21, 1–47.
- Umlauf, N., Kneib, T., Lang, S. and Zeileis, A. (2016). **R2BayesX**: Estimate structured additive regression models with **BayesX**. *R* package version 1.1. <http://cran.r-project.org>.
- Verbeke, G. and Molenberghs, G. (2000). *Linear Mixed Models for Longitudinal Data*. New York: Springer-Verlag.
- Verbyla, A.P., Cullis, B.R., Kenward, M.G. and Welham, S.J. (1999). The analysis of designed experiments and longitudinal data by using smoothing splines (with discussion). *Journal of the Royal Statistical Society, Series C*, **48**, 269–312.
- Verdinelli, I. and Wasserman, L. (1991). Bayesian analysis of outlier problems using the Gibbs sampler. *Statistics and Computing*, **1**, 105–117.
- Wahba, G. (1990). *Spline Models for Observational Data*. Philadelphia: SIAM.
- Wand, M.P. (2009). Semiparametric regression and graphical models. *Australian and New Zealand Journal of Statistics*, **51**, 9–41.
- Wand, M.P. and Ripley, B.D. (2015). **KernSmooth**: Functions for kernel smoothing supporting Wand and Jones (1995). *R* package version 2.23. <http://cran.r-project.org>
- Wand, M.P. and Jones, M.C. (1995). *Kernel Smoothing*. London: Chapman and Hall.
- Wand, M.P. and Ormerod, J.T. (2008). On semiparametric regression with O'Sullivan penalized splines. *Australian and New Zealand Journal of Statistics*, **50**, 179–198.
- Wand, M.P. and Ormerod, J.T. (2011). Penalized wavelets: embedding wavelets into semiparametric regression. *Electronic Journal of Statistics*, **5**, 1654–1717.
- Wang, Y. (1998). Mixed effects smoothing spline analysis of variance. *Journal of the Royal Statistical Society, Series B*, **60**, 159–174.
- Wickham, H. and Chang, W. (2016). **ggplot2**: Create elegant data visualisations using the grammar of graphics. *R* package version 2.2.1 <https://ggplot2.tidyverse.org>.
- Wong, R.K.W., Yao, F. and Lee, T.C.M. (2013). **robustgam**: Robust estimation for generalized additive models. *R* package version 0.1.7. <http://www.r-project.org>.
- Wood, S.N. (2003). Thin plate regression splines. *Journal of the Royal Statistical Society, Series B*, **65**, 95–114.
- Wood, S.N. (2006a). *Generalized Additive Models*. Boca Raton, Florida: Chapman & Hall/CRC.
- Wood, S.N. (2006b). Low rank scale invariant tensor product smooths for generalized additive mixed models. *Biometrics*, **62**, 1025–1036.
- Wood, S.N. (2017). **mgcv**: Mixed GAM computation vehicle with GCV/AIC/REML smoothness estimation. *R* package version 1.8. <http://cran.r-project.org>.
- Wood, S.N., Li, Z., Shaddick, G. and Augustin, N.H. (2017). Generalized additive models for gigadata: modelling the U.K. black smoke network daily data. *Journal of the American Statistical Association*, **112**, 1199–1210.



- Wu, H. and Zhang, J.-T. (2006). *Nonparametric Regression Methods for Longitudinal Data Analysis*. Hoboken, New Jersey: John Wiley & Sons.
- Xiao, L., Li, Y. and Ruppert, D. (2013). Fast bivariate P-splines: the sandwich smoother. *Journal of the Royal Statistical Society, Series B*, **75**, 577–599.
- Xiao, L., Ruppert, D., Zipunnikov, V. and Crainiceanu, C. (2016). Fast covariance estimation for high-dimensional function data. *Statistics and Computing*, **26**, 409–421.
- Yee, T.W. (2004). Quantile regression via vector generalized additive models. *Statistics in Medicine*, **23**, 2295–2315.
- Yee, T.W. (2010). The **VGAM** package for categorical data analysis. *Journal of Statistical Software*, **32**, Issue 10, 1–34.
- Yee, T.W. (2017). **VGAM**: Vector generalized linear and additive models. **R** package version 1.0. <http://www.stat.auckland.ac.nz/~yee/VGAM>.
- Yeo, I.-K. and Johnson, R.A. (2000). A new family of power transformations to improve normality or symmetry. *Biometrika*, **87**, 954–959.
- Zeger, S. and Diggle, P.J. (1994). Semiparametric models for longitudinal data with application to CD4 cell numbers in HIV seroconverters. *Biometrics*, **50**, 689–699.
- Zeger, S., Liang, K.-Y. and Albert, P.S. (1988). Models for longitudinal data: a generalized estimating equation approach. *Biometrics*, **44**, 1049–1060.
- Zhang, C., Shin, S.J., Wang, J., Wu, Y., Zhang, H.H. and Liu, Y. (2013). **probsvm**: Class probability estimation for support vector machines. **R** package version 1.00. <http://www.r-project.org>.
- Zhang, D., Lin, X., Raz, J. and Sowers, M. (1998). Semiparametric stochastic mixed models for longitudinal data. *Journal of the American Statistical Association*, **93**, 710–719.
- Zhang, H.H. and Lin, C.-Y. (2013). **cozzo**: Fit regularized nonparametric regression models using COSSO. **R** package version 2.1. <http://www.r-project.org>.
- Zhao, T., Li, X., Liu, H. and Roeder, K. (2014) **SAM**: Sparse additive modeling. **R** package version 1.0.5. <http://www.r-project.org>.
- Zhao, Y., Staudenmayer, J., Coull, B.A. and Wand, M.P. (2006). General design Bayesian generalized linear mixed models. *Statistical Science*, **21**, 35–51.

# Index

## A

Additive mixed model, 132, 133, 135, 137, 139, 142, 163, 167, 168, 174  
  generalized, 164  
Additive model, 71, 230, 250, 251, 253, 273, 311  
  functional with principal component scores, 251  
  heteroscedastic, 302, 305  
  quantile, 240  
  robust, 231  
AIC, *see* Akaike's information criterion  
Akaike's information criterion, 30, 31, 37, 254, 311  
  corrected, 31

## B

Basis, 16, 17, 19, 20  
Bayes estimate, 46, 47, 49, 58, 123, 139, 171, 175, 284, 288, 292, 295, 315, 317, 318  
Bayesian analysis, 137  
Bayesian inference engine, 225, 276, 279, 295, 297, 298, 306, 308  
Bernoulli distribution, 75  
Best linear unbiased predictor, 31, 33, 52, 157  
Box–Cox transformation, 237, 238  
Brain imaging, 213, 243  
Brooks–Gelman–Rubin convergence  
  diagnostics, 50  
BUGS, 41, 59, 225, 279, 280, 295, 297, 308

## C

Canonical correlation analysis, 206  
Capital Asset Pricing Model, 199

Classification, 124, 127, 265, 267–274, 276, 311–314  
Component selection and smoothing operator, 99–101, 103, 120  
Comprehensive R Archive Network, 1  
Confidence interval, 36, 46, 121, 136, 166  
  pointwise, 2–4, 35, 36, 65, 84, 87, 92, 94, 97, 111, 115, 117, 119, 134, 148, 167, 172, 175, 190, 191, 200–203, 249  
Confusion matrix, 125, 126, 272, 275, 311, 313  
Contrast function, 8, 58, 59, 69, 145–149, 171–174, 289, 291, 299, 302  
Coronary heart disease, 293–295  
Corpus callosum, 243  
Covariance function, 205, 207, 211, 212  
  estimation, 204, 207, 208, 210, 212, 223, 224, 249  
Credible interval, *see* credible set  
Credible set, 46, 47, 50, 139, 161, 175, 233, 288, 292, 306  
  pointwise, 6, 47, 58, 59, 65, 69, 123, 139, 148, 159, 161, 168, 171–174, 284, 288, 291, 295, 302, 306, 317, 318  
Credit rating, 76, 93, 94, 119  
Curse of dimensionality, 178

## D

Dataset  
  Bangladesh contraception, 174  
  Boston mortgage applications, 4, 75, 76, 90  
  California schools, 82, 83, 111  
  car auctions, 269, 273, 276, 313  
  coral ecology, 295, 298, 300–302

Dataset (*cont.*)

- coronary heart disease, 294
  - diffusion tensor imaging, 243, 244
  - ethanol exhaust fumes, 222
  - gasoline spectra, 204–206, 210, 248, 249
  - high-flux hemodialyzer, 172
  - Indiana adolescent growth, 6, 140, 142, 145, 148–150, 172–174
  - Indonesian children respiratory health, 163, 164, 166–168
  - intensive care unit, 122
  - lake acidity, 220
  - light detection and ranging, 316
  - meat content, 246, 247, 251–254
  - mental health clinical trial, 289
  - Michigan income dynamics, 11, 226, 227, 229–232, 236–238, 314
  - nutritional epidemiology, 153, 155
  - ozone in Los Angeles, 302–306
  - ozone in Midwest USA, 179, 181, 185
  - physician office visits, 78, 85, 86
  - Pima Indians diabetes, 286, 311
  - plankton flow cytometry, 311–313
  - protein in cows' milk, 173
  - ragweed pollen, 128, 317
  - school results, 173
  - spam e-mail, 124
  - spinal bone mineral density, 132, 134, 136, 137, 139, 171, 210–212
  - stock returns, 198, 199, 203, 204
  - Sydney real estate, 9, 188, 190, 191, 193, 197
  - Utah peak expiratory flow, 174
  - Warsaw apartments, 2, 4, 16, 38, 40, 42, 47, 53, 54, 56, 113, 267, 309
  - yield curves, 256, 258, 261, 264
- Density estimation, 124
- bivariate, 221
- Deviance, 28, 73, 80, 97
- scaled, 73
- Diabetes, 286, 288, 311
- Diagnostics, 23–26, 50, 65, 98, 120, 122, 123, 128, 175, 181–183, 185, 188, 193, 194, 215, 221, 291
- Diffusion tensor imaging, 243
- Directed acyclic graph, 277–281, 287, 300, 306
- E**
- EDF, *see* effective degrees of freedom
- Effective degrees of freedom, 22, 28–30, 37, 40, 47, 49, 93, 96, 110, 136, 147, 148, 165, 183, 231, 240, 267, 268, 317
- Effective Monte Carlo sample size, 49

Expectation propagation, 279

Exponential family, 72

**F**

- FACE, *see* fast covariance estimator
- Factor-by-curve interaction model, *see* interaction
- Fast covariance estimator, 207–209, 214, 248
- Finite mixture model, 292
- F*-test, 37–40, 56, 66, 85, 199, 200, 202
- F*-statistic, 39
- Functional data, 204–206, 215, 241, 250, 251, 254, 256–258, 260, 308
- Functional magnetic resonance imaging, 213
- Function-on-function regression, 254–256, 258, 262–265

**G**

- GAM, *see* generalized additive model
- Gaussian processes, 177
- GCV, *see* generalized cross-validation
- Generalized additive model, 71, 124, 125, 251
- Generalized cross-validation, 21, 22, 30, 31, 34, 39, 52, 65–68
- Generalized linear model, 71, 125, 126
- Geoadditive model, 187, 188, 197, 220, 222
- Geostatistics, 177, 215
- GLM, *see* generalized linear model
- GNU Linear Programming Kit**, 99
- Gram matrix, 266, 267, 271, 274
- Graphical model, 225, 276, 277, 279–281, 295, 298, 306, 308
- Group-specific curves, 140, 142, 145, 148, 150, 172–174, 297, 298, 300, 302

**H**

- Half-Cauchy prior on a standard deviation parameter, 41, 43, 137, 157, 167, 232, 278, 283, 287, 290, 294, 299, 306, 315, 318
- Hamiltonian Monte Carlo, 41
- Heat map, 214, 216, 251
- Heteroscedasticity, 40, 52, 70, 185, 194, 300, 302–304, 306, 316, 317
- Hinge loss, 268, 271
- Hyperparameter, 41, 42, 69, 138, 232, 283, 290, 295, 315, 318
- Hypothesis test, 23, 36, 38, 39, 56, 66, 67, 85, 86, 92, 93, 118, 122, 136, 183, 199, 200, 202, 204, 244

**I**

- Infer.NET, 279
- Interaction, 55–57, 115, 160, 177, 178
  - factor-by-curve, 58, 59, 68, 113, 115, 117, 119, 127, 128, 134, 146, 161, 318
  - factor-by-curve additive/interaction, 317

**J**

- Jones–Faddy skewed  $t$  distribution, 236

**K**

- Kaggle, 269–272
- Karush–Kuhn–Tucker conditions, 274
- Kernel, 265
  - radial basis function, 267, 271–273, 275, 312–314
- Kernel machine, 177, 265–268, 271
  - least-squares, 266
- $k$ -fold cross-validation, 100, 101, 103, 271, 274
- $k$ -index diagnostic, 24, 122, 182, 183, 185, 197
- Knot, 16–20, 33, 34, 62, 64, 126, 209, 213, 248, 283, 284, 287, 316, 317
- Kriging, 177

**L**

- LASSO, *see* least absolute shrinkage and selection operator, 99
- Least absolute shrinkage and selection operator, 99
- LIBSVM, 270
- Likelihood ratio test, 36
- Linear mixed model, 31, 33, 164
- Link, 72, 73, 75, 82, 87, 92, 250, 299
  - canonical, 72, 73
  - comparison of logit and probit, 73
  - logit, 72
  - probit, 73
- Logistic regression, 75, 91, 125
  - nonparametric, 286, 288
- Logit function, 72
- log-odds, 72, 91
- Longitudinal data, 6, 52, 131, 132, 149, 153, 161, 168, 171–174, 266
- Loss function, 265–268, 271

**M**

- Marginal model, 150, 151, 156
  - additive, 160
- Marginal nonparametric regression, *see* nonparametric regression

- Markov chain Monte Carlo, 41, 42, 44–50, 52, 59, 65, 167, 232, 284–288, 291–293, 295, 296, 302, 304, 306, 308, 315
- MCMC, *see* Markov chain Monte Carlo
- Mean field variational Bayes, 279
- Measurement error, 276
  - Berkson, 281
  - classical, 281, 282, 284, 285, 293
- M-estimation, 227, 228
- Misclassification rate, 125, 126, 271, 272, 275, 311, 313, 314
- Missing data, 276, 277, 280, 286, 295, 300
  - at random, 281
  - completely at random, 281, 306, 317
  - not at random, 281, 286
  - at random, 281
- Mixed model, *see* linear mixed model
- Mixed model-based penalized splines, *see* penalized splines
- Model
  - saturated, 73
- Model selection
  - penalty-based, 99
- Monte Carlo standard error, 49
- Moral graph, 279
- Multilevel model, 131, 169
- Multiple sclerosis, 243

**N**

- Near infrared, 246, 248
- Negative Binomial response, 78, 87, 317, 318
- Nonparametric regression, 2, 15, 21, 22, 29, 31, 39, 41, 43, 45, 47, 50, 53, 56, 61, 64–67, 226, 273, 277, 281, 288
  - bivariate, 178, 179, 198, 199, 202, 203, 223
  - heteroscedastic, 69, 316
  - marginal, 151, 156, 157
  - measurement error, 282, 284
  - partially observed data, 280
  - skewed  $t$  response, 236
  - $t$  response, 232
- No U-turn sampler, 41

**O**

- Odds ratio, 75–77, 119, 175, 299, 302
- O’Sullivan spline, *see* spline
- Overdispersion, 74, 78, 80

**P**

- Panel data, 131, 169
- Penalized least-squares, 16

Penalized quasi-likelihood, 164, 298  
 Penalized spline, 15–23, 25–32, 34–36, 39–41, 48, 49, 51, 58, 59  
   Bayesian, 40, 47, 52  
   bivariate, 177, 180, 181, 183, 185, 188, 193, 197, 212, 213, 215, 218–221, 223, 224  
   mixed model-based, 31–34, 36–40, 47, 52, 65, 132, 133, 135, 137, 139, 142, 156, 163, 164, 167, 168, 174, 232, 273, 290, 305, 315, 318  
   tensor product, 181, 185, 213, 220, 224  
   thin plate, 180, 181, 183, 185, 188, 193, 197, 213, 220, 224  
   trivariate, 178  
 Poisson regression, 78, 80  
 PQL, *see* penalized quasi-likelihood  
 Principal components analysis, 205, 206, 244, 248, 251  
 Principal component scores, 251–254, 308, 311  
*p*-value, 24, 38–40, 57, 66, 80, 81, 85, 86, 94, 117, 137, 183, 185

## Q

Quadratic programming problem, 271, 274  
 Quantile-quantile plot, 26, 89, 185, 194–196  
 Quantile regression, 11, 196, 225, 237, 240, 309  
   LMS approach, 237, 238, 240, 241, 310  
 Quantile semiparametric regression, 237  
 Quantile smoothing spline, 228, 234, 238, 240, 241  
 Quasi-likelihood, 74, 75, 128  
 Quasi-Poisson model, 75, 80

## R

Random intercepts model, 155  
 Randomized quantile residual, 89  
 REML, *see* restricted maximum likelihood  
 Reproducing kernel Hilbert space, 265  
 Residual, 26, 89  
   deviance, 28, 73  
 Residual maximum likelihood, 66  
 Resistant regression, 225, 227, 229, 310  
 Resistant scatterplot smoothing, 226  
 Restricted likelihood ratio test, 37  
 Restricted maximum likelihood, 31–34, 39, 52, 65, 66, 68, 157  
 Return  
   excess, 198  
   on a stock, 198

## R function

`af()`, 251  
`anova()`, 57  
`apply()`, 208  
`BGRinterval()`, 50  
`bifd()`, 259  
`bkde()`, 46  
`bruto()`, 95  
`cosso()`, 101  
`create.bspline.basis()`, 246  
`createBoundary()`, 192  
`cv.gamsel()`, 105  
`dpik()`, 46  
`eigen()`, 208  
`eval.bifd()`, 260  
`eval.fd()`, 246  
`exactRLRT()`, 38, 40  
`expand.grid()`, 208  
`fbps()`, 213  
`fdPar()`, 259  
`ff()`, 263, 265  
`ffpc()`, 263, 265  
`fpca.face()`, 207–209, 214  
`gam()`, 21, 23, 25, 26, 28, 33, 39, 77, 78, 80, 82–84, 95, 96, 98, 113, 120, 123, 177, 179, 180, 190, 191, 198, 201, 204, 208, 209, 211–215, 218, 220–222, 249  
`gam.check()`, 23, 24, 26, 122, 181, 185, 188, 193  
`gamlss()`, 78, 87  
`gamm()`, 40, 134, 164, 171, 174, 177, 204  
`gamsel()`, 126  
`glm()`, 75, 77  
`glmmPQL()`, 164, 298  
`hare()`, 309  
`image()`, 217  
`image.plot()`, 185, 217, 260  
`ksline()`, 177  
`lf()`, 248, 251  
`linmod()`, 255, 260, 263, 264  
`lme()`, 32, 33, 142, 143, 148, 157, 172, 298  
`lms.bcn()`, 239, 240, 309  
`lms.yjn()`, 241  
`lms.yjn2()`, 241  
`locfit()`, 35  
`loess()`, 226, 228  
`monitor()`, 47  
`pairs()`, 188  
`pb()`, 89  
`persp()`, 187, 217  
`pfpr()`, 255, 263, 264  
`pfr()`, 244, 247, 248, 251  
`plogis()`, 91

- `plot.cosso()`, 104
- `plot.gam()`, 84, 91, 251
- `plotvgam()`, 231
- `pointsInPoly()`, 192
- `pol spline()`, 309
- `polymars()`, 95, 98, 126, 127
- `pspline()`, 309
- `qqmath()`, 196
- `qss()`, 228
- `range()`, 212
- `rgl.surface()`, 187
- `rqss()`, 228, 238, 240, 241, 309
- `s()`, 22
- `sd()`, 196
- `skewness()`, 195, 196
- `smooth.basis()`, 260
- `smooth.basisPar()`, 246
- `smooth.spline()`, 19–22, 29, 317
- `stan()`, 42, 44, 50, 232
- `step.Gam()`, 95–98, 121–123, 125, 222
- `stl()`, 309
- `studentt3()`, 230, 311
- `summMCMC()`, 47, 50
- `svm()`, 270–272, 311
- `tapply()`, 196
- `terrain.colors()`, 180, 181, 185, 186, 192, 202, 212, 216–218, 221
- `topo.colors()`, 202
- `Tps()`, 177
- `transform()`, 229
- `tune.cosso()`, 100, 102, 103
- `tune.svm()`, 271, 272, 311
- `US()`, 180, 185
- `vgam()`, 110, 229, 231, 239, 309–311
- `vis.gam()`, 191
- `xLMS()`, 239
- `xyplot()`, 132, 154
- `ZDaub()`, 309
- `ZOSull()`, 18, 309
- R function's argument
  - by (`gam()`), 198
  - family (`gam()`), 75, 80, 120, 128
  - family (`vgam()`), 230, 240
  - k (`s()`), 22
  - panel (`xyplot()`), 133
  - presmooth (`lf()`), 248
  - pve (`fpca.face()`), 207
  - Qtransform (`pfr()`), 251
  - s (`gam()`), 83
  - scale (`gam()`), 80
  - scheme (`plot.gam()`), 251
  - scope (`gam()`), 96
  - select (`gam()`), 104, 201
  - select (`plot.gam()`), 190
  - seWithMean (`plot.gam()`), 201
  - sp (`gam()`), 84
  - span (`loess()`), 226
- Robust regression, 225–229, 231, 310
- Robust scatterplot smoothing, 226
- Robust semiparametric regression, 229
- R package
  - AER, 120
  - aplore3, 122
  - BayesX, 59
  - BRugs, 41, 280, 295, 308
  - cosso, 99–101
  - e1071, 195, 270, 272, 311
  - Ecdat, 4, 64, 78, 82, 126, 226, 230
  - fda, 215, 246, 255, 258, 263, 308
  - fda.usc, 246
  - fields, 177, 179, 180, 185, 215, 217, 223, 260
  - gam, 71, 82, 95, 96, 110, 121–123, 125, 222
  - gamlss, 71, 78, 87
  - gamsel, 99–101, 105, 125, 126
  - geoR, 177
  - ggplot2, 133
  - gss, 122, 220
  - HRW, 15, 16, 18, 47, 50, 54, 57, 59, 84, 92, 97, 111, 117, 118, 132, 140, 148, 153, 157, 163, 172–174, 179, 185, 186, 192, 198, 203, 211, 215, 217–219, 222, 230, 233, 236, 240, 247, 251, 256, 269, 272, 274, 276, 281, 287–289, 294, 295, 298, 306, 309–311, 313, 316, 317
  - kernlab, 124, 127
  - KernSmooth, 46
  - lars, 309
  - lattice, 132, 133, 144, 196, 224
  - lme4, 59
  - locfit, 35
  - LowRankQP, 273, 314
  - MASS, 164, 298
  - mda, 95
  - mgcv, 15, 21, 23, 26, 28, 33, 37, 40, 52, 71, 77, 78, 82, 83, 96, 104, 110, 123, 124, 131, 134, 137, 164, 168, 171, 174, 177, 179, 180, 215, 218, 220, 222
  - mlbench, 286, 302, 311
  - nlme, 15, 32, 59, 131, 142, 168, 172, 173, 223
  - PBImisc, 2
  - pol spline, 71, 95, 98, 126
  - probsvm, 270
  - quantreg, 228, 237, 238, 240, 241, 309
  - refund, 204, 207, 213, 223, 243, 244, 255, 263
  - rgam, 229

R package (*cont.*)

[rgl](#), 187, 219  
[Rglpk](#), 99  
[RLRsim](#), 38, 40  
[robustgam](#), 229  
[R2WinBUGS](#), 41, 280  
[rstan](#), 15, 41, 42, 50, 52, 58, 123, 131,  
 137, 138, 148, 150, 157, 164, 167,  
 171–173, 175, 232, 235, 280, 287, 289,  
 291, 297, 301, 308, 310, 315, 316, 318  
[SAM](#), 99  
[scam](#), 120  
[survival](#), 308  
[svmpath](#), 270  
[tsDyn](#), 309  
[VGAM](#), 71, 110, 229, 237, 239, 241, 309–311  
[wavethresh](#), 309

## R script

[absorbBivarFigs.R](#), 251  
[absorbFuncsDrvs.R](#), 247  
[absorbScalOnFuncAna.R](#), 247  
[BCRana.R](#), 291, 292  
[BostMortFacByCurv.R](#), 118, 119  
[BostMortGAMfit.R](#), 82, 91  
[carAucPenSplSVM.R](#), 274, 314  
[carAucPenSplSVMtune.R](#), 274  
[carAucRadialSVM.R](#), 272  
[carAucRadialSVMtune.R](#), 272  
[CaSchoolGAMfit.R](#), 84  
[CaSchoolVGAMfit.R](#), 111  
[coralAna.R](#), 301, 302  
[femSBMDBayes.R](#), 138  
[IndonRespirBayes.R](#), 167  
[maleGrowthIndianaBayes.R](#), 148,  
 172–174  
[maleGrowthIndiananlme.R](#), 145, 148  
[margAddMod.R](#), 160  
[MichIncLMS.R](#), 240, 310  
[MichIncMCMCskewt.R](#), 236  
[MichIncMCMCt.R](#), 233  
[MichIncMultQSS.R](#), 238, 310  
[MichInctAddMod.R](#), 230, 310  
[npReg.R](#), 281  
[npRegBerksonMeaErr.R](#), 281  
[npRegClassicMeaErr.R](#), 281–283, 285  
[npRegMAR.R](#), 281  
[npRegMCAR.R](#), 281  
[npRegMNAR.R](#), 281  
[ozoneAna.R](#), 306  
[OFFPGAMfit.R](#), 82, 86  
[ozoneDisplayConvHull.R](#), 186  
[ozone3Dspin.R](#), 219  
[ozoneDisplayConvHull.R](#), 186  
[PIDana.R](#), 287

[proteinMargAddInt.R](#), 161  
[proteinMargNPREgn.R](#), 157  
[returnsDrvFig.R](#), 203  
[RLJGana.R](#), 295  
[SydneyDisplaySophis.R](#), 192  
[WarsawAptsBayes.R](#), 42, 47, 50, 65, 123  
[WarsawAptsGAMfit.R](#), 117  
[WarsawAptsSimpFacByCurv.R](#), 58, 59  
[WarsawAptsSimpSemi.R](#), 54  
[WarsawAptsSimpSemiInt.R](#), 56, 57

## S

Sandwich smoother, 213, 214, 224  
 Scalar-on-function regression, 241, 250, 251,  
 254, 255, 257  
 Scale parameter  
   of the quasi-likelihood function, 74  
 Scatterplot matrix, 188  
 Scree plot, 252, 253  
 Serial correlation, 65, 139, 174  
 Shape constraint, 120  
 Simple semiparametric regression, 53  
 Skewness, 26, 89, 194, 195  
 Smearing estimate, 221  
 Smoothing parameter, 16, 21, 23, 28, 30, 31,  
 65  
   selection, 22, 31, 32, 39, 65, 66  
 Smoothing spline, 19, 20, 22, 23  
   low-rank, 26  
 Soap film smooths, 179  
 Somatic growth, 140  
 Sparse additive models, 120  
 Spline  
   B-spline, 17, 19, 61–64, 246, 251, 258  
   O’Sullivan, 17–20, 23, 26, 69, 81, 138, 142,  
   164, 172, 173, 266, 267, 283, 290, 305,  
   309, 315  
   tensor product, 178–181, 183–185, 202,  
   207–209, 213, 220, 224, 251  
   thin plate, 65, 178–181, 183, 185, 188, 193,  
   197, 213, 220, 224, 249  
[Stan](#), 41–43, 49, 59, 138, 157, 167, 171–173,  
 225, 232, 235, 236, 279–281, 283, 284,  
 287, 291, 292, 295, 297, 298, 301, 302,  
 306, 308, 310, 315–318  
   compared to [BUGS](#), 308  
 Standard & Poor’s 500 (S&P 500) index, 198,  
 200, 202, 203  
 Stepwise model selection, 95, 96  
 Support vector machine, 265, 268, 270–273,  
 311, 312, 314  
   low-rank, 272  
   penalized spline, 273–275  
 Survival analysis, 308

**T**

*t* distribution, 229–231, 233, 234, 310  
Tecator Infratec Food and Feed Analyzer, 246  
Tensor product spline, *see* spline  
Thinning, 44, 287, 291  
Thin plate spline, *see* spline  
Time series analysis, 69, 309  
Trace plot, 46, 47  
Treasury bill, 198

**V**

Variability band, 3, 35, 36, 65, 117, 148, 190, 244, 249  
Variance-stabilizing transformation, 185

Varying-coefficient model, 197–200, 202–204, 222

Vector response, 110

**W**

Warm-up, 44, 46, 47  
Wavelet regression, 309  
White matter tract, 243

**Y**

Yeo–Johnson transformation, 241  
Yield curve, 256–258, 260