
Index

A

- Aflatoxin, *see also specific aflatoxins*,
analytical variance, 19, 20
capillary electrophoresis,
 data analysis, 56
 electrophoresis, 56
 extraction, 55, 56
 fluorescence detection and
 sensitivity, 53
 materials and instrumentation,
 53–57
 micellar electrokinetic capillary
 chromatography, 52, 53
 rapidity, 53
 standards, 56
distribution in samples, 17, 18
fluorescence, 52
intra-lot variability of testing, 12, 13
sample preparation variability, 18
sample size effects on assay
 variability, 20–22
standards, preparation,
 materials, 33
 overview, 31, 33
 purity analysis,
 liquid chromatography, 34
 thin-layer chromatography,
 34, 35
 solution preparation, 33–35
 stability, 35
 vial preparation of dry film
 standards, 34–36
ultraviolet absorption, 51
- Aflatoxin B₁, screening, 51
- Aflatoxin M₁,
 extraction considerations, 61
 milk analysis with immunoaffinity
 chromatography cleanup and
 high-performance liquid
 chromatography,
 apparatus, 64, 67
 data analysis, 66
 fluorescence detection and
 sensitivity, 63
 high-performance liquid
 chromatography, 65, 66, 68
 immunoaffinity chromatography,
 cleanup, 65, 68
 principle, 61
 performance, 63
 reagents, 64, 66, 67
 sample preparation,
 dry milk, 65, 68
 liquid milk, 64, 68
 validation, 61, 63
 origins in milk, 59
 seasonal trends in milk
 contamination, 59, 60
 structure, 59, 60
 tolerance levels, 60, 61
 toxicity, 59
 trends in milk contamination, 60
- Alternaria* toxins,
 crop distribution, 226
 liquid chromatography analysis of
 alternariol and methyl ester in
 apple juice,
 chromatography conditions, 229

- cleanup,
 - aminopropyl column, 228
 - cloudy samples, 228, 229
 - reverse-phase solid-phase extraction, 227, 228
- materials, 227, 231
- overview, 226, 227
- performance, 229, 230
- toxicity, 225, 226
- types and structures, 225, 226
- Analytical variance,
 - calculation, 19, 20
 - technique dependence, 19, 20
- Antibody preparation, *see* Citrinin; Ochratoxin A; Zearalenone
- AOACI, Official Method validation, 6, 63
- Apple juice, *see* *Alternaria* toxins; Patulin
- Atrones, *see* *Stachybotrys atra* toxins
- B**
- Blood analysis, *see* Ochratoxin A
- C**
- Calibrating standards, *see* Standards
- Capillary electrophoresis (CE),
 - aflatoxins,
 - data analysis, 56
 - electrophoresis, 56
 - extraction, 55, 56
 - fluorescence detection and sensitivity, 53
 - materials and instrumentation, 53–57
 - micellar electrokinetic capillary chromatography, 52, 53
 - rapidity, 53
 - standards, 56
 - cyclopiazonic acid, 73
 - formats, 52, 53
 - principle, 52
- CE, *see* Capillary electrophoresis
- Charcoal-aluminum-celite column,
 - preparation, 119, 120
 - trichothecene cleanup, 120, 121
- Citrinin,
 - calibration by ultraviolet spectroscopy, 32
 - crop and geographic distribution, 195, 196
 - enzyme immunoassays,
 - antigen conjugate preparations, 197, 199
 - direct assay, 198, 200
 - immunization of rabbits, 197–200
 - indirect assay, 198–200
 - materials, 196, 197, 199
 - sample preparation, 198
 - species in production, 195
 - structure, 199, 200
 - toxicity, 195
- CPA, *see* Cyclopiazonic acid
- Cross-cut sampler,
 - automated sampling, 15, 16
 - installation, 16
 - sample size, 17
 - time between cuts, 17
- Cyclopiazonic acid (CPA),
 - analysis with immunoaffinity
 - chromatography cleanup and high-performance liquid chromatography,
 - drought stress growth effects, 77, 78
 - high-performance liquid chromatography, 75
 - immunoaffinity
 - chromatography, 75, 76
 - materials, 74, 75
 - performance, 76
 - sample preparation and extraction
 - from peanuts, 75

calibration by ultraviolet spectroscopy, 32
 capillary electrophoresis, 73
 disease roles, 72
 enzyme-linked immunosorbent assay, 73, 74
 food distribution, 71
 high-performance liquid chromatography separation systems, 72, 73
 species in production, 71
 spectrophotometric assays, 73
 structure, 72
 thin-layer chromatography, 72, 73

D

DAS, *see* Diacetoxyscirpenol
 Deoxynivalenol (DON), *see also* Trichothecenes,
 advisory levels in feeds, 97, 98
 calibration by ultraviolet spectroscopy, 32
 fluorescence assay,
 accuracy and precision, 105–107
 cleanup column, 100, 101, 111
 cross-reactivity analysis, 104, 105
 derivatization for fluorescence, 102
 fluorometry,
 calibration, 102–104
 operating procedure, 103, 104, 111
 linearity of assay, 104
 materials, 99
 overview, 98, 99
 sample preparation and extraction, 100, 110, 111
 validation by high-performance liquid chromatography,
 external standard calibration, 110
 materials, 99, 100

purification of samples, 108, 109, 112
 sample preparation and extraction, 107
 spike preparation, 108, 111, 112
 growth conditions and grain contamination, 98
 monoacetate derivatives, 115, 116
 reverse-phase high-performance liquid chromatography, 122, 126
 species in production, 97, 115
 structure, 97, 98
 toxicity, 97, 117

Diacetoxyscirpenol (DAS),
 analysis in feeds, *see* Trichothecenes
 species in production, 115
 toxicity, 116, 117

DON, *see* Deoxynivalenol

E

ELISA, *see* Enzyme-linked immunosorbent assay
 Enzyme-linked immunosorbent assay (ELISA),
 citrinin,
 antigen conjugate preparations, 197, 199
 direct assay, 198, 200
 immunization of rabbits, 197–200
 indirect assay, 198–200
 materials, 196, 197, 199
 sample preparation, 198
 ochratoxin A,
 antibody preparation,
 horseradish peroxidase conjugate preparation as antigen, 85, 90
 immunization of rabbits, 85, 90, 91
 materials, 83, 89

serum albumin conjugate
 preparation as antigen,
 84, 85, 90
 titer determination, 83,
 85–87, 91
 competitive direct enzyme
 immunoassay,
 cereal sample preparation,
 87, 91
 incubation conditions, 88,
 89, 91
 materials, 84, 90
 overview, 82
 porcine blood serum sample
 preparation, 87, 88, 92
 overview of mycotoxin analysis,
 73, 74
 zearalenone assays,
 detection limits, 168, 169
 direct competitive assay,
 overview, 160, 16
 indirect competitive assay,
 overview, 160–163
 materials, 164
 monoclonal antibody assays,
 165, 166
 polyclonal antibody assays,
 165, 166
 recombinant antibody,
 competitive direct assay
 application, 166
 competitive indirect assay
 application, 162, 163,
 167, 168
 cross-reactivity compared
 with monoclonal
 antibody, 162, 163
 generation, 162
 sample preparation, 165
 Ergot alkaloids,
 crop and geographic distribution, 215

disease roles, 215, 217
 economic significance, 217
 liquid chromatography analysis,
 chromatography conditions, 221
 data analysis, 221, 223
 extraction and cleanup, 220,
 221, 223
 materials, 219–221
 overview, 218, 219
 standards, 220, 221, 223
 species in production, 215
 structures, 215, 216
 toxicity, 216, 217

F

F-2 toxin, *see* Zearalenone
 Florisil column,
 preparation, 120
 trichothecene cleanup, 121, 126
 Fumonisin,
 classification, 147
 disease roles, 26, 147, 148
 economic impact, 148
 extraction considerations, 149
 high-performance liquid
 chromatography, corn analysis,
 chromatography conditions,
 152, 154
 cleanup, 151, 154
 data analysis, 155, 156
 extraction, 151, 154
 materials, 150, 151
 overview, 149, 150
 precolum fluorescence
 derivatization, 152
 standards and calibration, 151,
 152, 154
 high-performance liquid
 chromatography/electrospray
 ionization mass spectrometry
 analysis,

- data acquisition, 43
 - instrumentation, 42, 44
 - materials, 42–44
 - overview, 41, 42
 - purity calculations, 43, 45, 46
 - sample preparation, 42
 - history of study, 25
 - isolation from solid phase fungal culture,
 - ammonium salt conversion, 28
 - extraction, 27, 28
 - Fusarium proliferatum* culture, 26, 28
 - high-performance liquid chromatography,
 - analytical equipment, 26, 27
 - preparatory equipment, 26
 - running conditions, 27
 - reagents, 27
 - molar response factors in detection, 41
 - species in production, 147
 - structures, 149
 - tolerance levels, 148, 149
 - toxicity mechanisms, 148
- G**
- Gas chromatography (GC),
 - Stachybotrys atra* toxins, gas chromatography/mass spectrometry,
 - data acquisition and analysis, 189, 192
 - derivatization, 182, 183, 189
 - hydrolysis of crude extract, 188, 189, 191, 192
 - materials, 184
 - trichothecenes,
 - electron-capture detection, 124, 125
 - heptafluorobutyrylation
 - derivatization, 123, 126
 - mass spectrometry detection, 124
 - trimethylsilylation derivatization, 123, 126
 - GC, *see* Gas chromatography
- H**
- High-performance liquid chromatography (HPLC),
 - aflatoxins, *see* Aflatoxin M₁
 - analytical variance, 19
 - cyclopiazonic acid, 72, 73, 75, 76
 - deoxynivalenol fluorescence assay
 - validation,
 - external standard calibration, 110
 - materials, 99, 100
 - purification of samples, 108, 109, 112
 - sample preparation and extraction, 107
 - spike preparation, 108, 111, 112
 - fumonisin isolation from solid phase fungal culture,
 - analytical equipment, 26, 27
 - preparatory equipment, 26
 - running conditions, 27
 - moniliformin,
 - apparatus, 135
 - buffers, 136
 - chromatography conditions, 137
 - cleanup, 137–139
 - overview of analytical approaches, 134, 135
 - performance, 142
 - reagents, 136
 - sample preparation, 136–138
 - spiking and recovery, 137, 139–142
 - stability of toxin, 138
 - standards, 136
 - resolution of mycotoxins, 37
 - Stachybotrys atra* toxins,
 - calibration curve, 187, 188, 191

chromatography conditions, 187
 culture preparation, 184, 189, 190
 extraction,
 aqueous, 184, 185, 190
 organic, 184, 190
 materials, 183, 189
 overview, 182
 polyethyleneimine silica cleanup,
 181, 182, 185, 189
 sample preparation, 187, 190, 191
 stationary phases, 176
 toxin isolation,
 aqueous extract, 186, 190
 organic extract, 185
 ultraviolet-visible diode array
 detection, 175, 176
 trichothecenes, 122, 126
 High-performance liquid
 chromatography/electrospray
 ionization mass spectrometry
 (HPLC/ES/MS),
 advantages over ultraviolet
 detection, 39, 40
 detection limits, 38
 fumonsin analysis,
 data acquisition, 43
 instrumentation, 42, 44
 materials, 42–44
 overview, 41, 42
 purity calculations, 43, 45, 46
 sample preparation, 42
 ionizability of specific mycotoxins,
 38, 39
 molar response, 40, 41
 overview, 37, 38
 rationale for mycotoxin analysis, 37
 HORRAT, *see* Horwitz Horn
 Horwitz equation, 4, 5
 Horwitz Horn, precision estimate by
 analyte concentration, 4
 Horwitz ratio (HORRAT),

calculation, 5
 method evaluation, 6
 table of evaluated methods, 7, 8
 HPLC, *see* High-performance liquid
 chromatography
 HPLC/ES/MS, *see* High-performance
 liquid chromatography/
 electrospray ionization mass
 spectrometry HT-2 toxin,
 analysis in feeds, *see* Trichothecenes,
 species in production, 115

I

Immunization, *see* Citrinin;
 Ochratoxin A
 Immunoaffinity chromatography
 cleanup, *see* Aflatoxin M₁;
 Cyclopiazonic acid
 Immunoassay,
 analytical variance, 19, 20
 enzyme-linked immunosorbent
 assay, *see* Enzyme-linked
 immunosorbent assay

L

LC, *see* Liquid chromatography
 Liquid chromatography (LC), *see also*
 High-performance liquid
 chromatography,
 aflatoxin standards, 34
 alternariol and methyl ester
 analysis in apple juice,
 chromatography conditions, 229
 cleanup,
 aminopropyl column, 228
 cloudy samples, 228, 229
 reverse-phase solid-phase
 extraction, 227, 228
 materials, 227, 231
 overview, 226, 227
 performance, 229, 230
 ergot alkaloids,

- chromatography conditions, 221
 - data analysis, 221, 223
 - extraction and cleanup, 220, 221, 223
 - materials, 219–221
 - overview, 218, 219
 - standards, 220, 221, 223
 - patulin assay in apple juices, calculations, 209, 210
 - chromatography and ultraviolet detection, 209, 211, 212
 - materials, 207, 208, 210
 - overview, 207
 - sensitivity, 207, 210,
 - solid-phase extraction, 208–211
- M**
- Mass spectrometry,
 - analyte identification, 9
 - atmospheric pressure chemical ionization, 38
 - electrospray ionization, *see* High-performance liquid chromatography/electrospray ionization mass spectrometry
 - fast atom bombardment, 41
 - gas chromatography/mass spectrometry, *see* Gas chromatography
 - Milk, *see* Aflatoxin M₁
 - Moniliformin,
 - calibration by ultraviolet spectroscopy, 32
 - crop and geographic distribution, 131, 132
 - high-performance liquid chromatography, apparatus, 135
 - buffers, 136
 - chromatography conditions, 137
 - cleanup, 137–139
 - overview of analytical approaches, 134, 135
 - performance, 142
 - reagents, 136
 - sample preparation, 136–138
 - spiking and recovery, 137, 139–142
 - stability of toxin, 138
 - standards, 136
 - species in production, 131
 - structure, 131, 132
 - thin-layer chromatography, 133, 134
 - toxicity, 132, 133
- N**
- NIV, *see* Nivalenol
 - Nivalenol (NIV),
 - analysis in feeds, *see* Trichothecenes
 - sources, 115, 116
- O**
- Ochratoxin A,
 - calibration by ultraviolet spectroscopy, 32
 - disease roles, 81
 - enzyme immunoassay, antibody preparation, horseradish peroxidase conjugate preparation as antigen, 85, 90
 - immunization of rabbits, 85, 90, 91
 - materials, 83, 89
 - serum albumin conjugate preparation as antigen, 84, 85, 90
 - titer determination, 83, 85–87, 91
 - competitive direct enzyme immunoassay, cereal sample preparation, 87, 91

- incubation conditions, 88, 89, 91
 - materials, 84, 90
 - overview, 82
 - porcine blood serum sample preparation, 87, 88, 92
 - plasma concentration versus intake, 81, 82
 - structure, 81, 82
 - tolerable daily intake, 82
- P**
- Patulin,
- calibration by ultraviolet spectroscopy, 32
 - liquid chromatography assay of apple juices,
 - calculations, 209, 210
 - chromatography and ultraviolet detection, 209, 211, 212
 - materials, 207, 208, 210
 - overview, 207
 - sensitivity, 207, 210
 - solid-phase extraction, 208–211
 - overview of apple juice analysis, 206, 207
 - properties, 205
 - species in production, 205
 - structure, 205, 206
 - tolerable daily intake, 206
 - toxicity, 205, 206
- Precision, mycotoxin assays, 4
- R**
- Reproducibility, mycotoxin assays, 4
- Roridins, *see Stachybotrys atra* toxins
- S**
- Sample preparation, variability, 18
- Sampling,
 - contamination distribution, 17, 18
 - selection methods,
 - cross-cut samplers, 15–17
 - dynamic lots, 15–17
 - random sampling, 14
 - static lots, 14, 15
 - size effects on assay variability, 20–22
 - USDA recommendations, 3
- Silica gel column,
 - preparation, 120
 - trichothecene cleanup, 121, 122
- Stachybotrys atra* toxins,
 - absorption and extinction coefficients, 176, 178, 179
 - classification, 176, 177, 180
 - extraction considerations, 180–182
 - gas chromatography/mass spectrometry,
 - data acquisition and analysis, 189, 192
 - derivatization, 182, 183, 189
 - hydrolysis of crude extract, 188, 189, 191, 192
 - materials, 184
 - high-performance liquid chromatography,
 - calibration curve, 187, 188, 191
 - chromatography conditions, 187
 - culture preparation, 184, 189, 190
 - extraction,
 - aqueous, 184, 185, 190
 - organic, 184, 190
 - materials, 183, 189
 - overview, 182
 - polyethyleneimine silica cleanup, 181, 182, 185, 189
 - sample preparation, 187, 190, 191
 - stationary phases, 176
 - toxin isolation,
 - aqueous extract, 186, 190
 - organic extract, 185

- ultraviolet-visible diode array
 - detection, 175, 176
- history of study, 173
- structures, 174, 175
- toxicity, 173, 174
- Standards,
 - aflatoxin standard preparation,
 - materials, 33
 - overview, 31, 33
 - purity analysis,
 - liquid chromatography, 34
 - thin-layer chromatography, 34, 35
 - solution preparation, 33–35
 - stability, 35
 - vial preparation of dry film
 - standards, 34–36
 - ultraviolet spectroscopy
 - calibration, 31, 32
- Sterigmatocystin, calibration by
 - ultraviolet spectroscopy, 32
- Stratoxins, *see Stachybotrys atra* toxins
- T**
- T-2 toxin,
 - analysis in feeds, *see* Trichothecenes,
 - species in production, 115
 - toxicity, 116, 117
- Thin-layer chromatography (TLC),
 - aflatoxin standards, 34, 35
 - analytical variance, 19
 - cyclopiazonic acid, 72, 73
 - moniliformin, 133, 134
 - trichothecenes, 122, 123, 126
- TLC, *see* Thin-layer chromatography
- Trichoverrols, *see Stachybotrys atra* toxins
- Trichothecenes, *see also* Deoxynivalenol; Diacetoxyscirpenol; Nivalenol; HT-2 toxin; *Stachybotrys atra* toxins; T-2 toxin,
 - advisory levels, 116
 - analysis in feeds,
 - apparatus, 119
 - charcoal-aluminum-celite column,
 - cleanup, 120, 121
 - preparation, 119, 120
 - Florisil column,
 - cleanup, 121, 126
 - preparation, 120
 - gas chromatography,
 - electron-capture detection,
 - 124, 125
 - heptafluorobutyrylation
 - derivatization, 123, 126
 - mass spectrometry
 - detection, 124
 - trimethylsilylation
 - derivatization, 123, 126
 - high-performance liquid
 - chromatography, 122, 126
 - overview, 117
 - reagents, 117, 119, 126
 - silica gel column,
 - cleanup, 121, 122
 - preparation, 120
 - solid-phase extraction column
 - cleanup, 120, 126
 - thin-layer chromatography, 122,
 - 123, 126
 - classification, 115
 - species in production, 115
 - toxicity, 116, 117
- U**
- USDA recommendations, sample size
 - for mycotoxin testing, 3
- V**
- Validation, mycotoxin assays, 6, 9
- Variability, *see also* Horwitz ratio,
 - analytical variability, 19, 20
 - intra-lot variation, 12, 13
 - reduction techniques, 20–22

- sample preparation variability, 18
- sampling variability, 13–18
- total variability components, 13, 22
- Verrucarins, *see Stachybotrys atra* toxins

Z

Zearalenone,

- calibration by ultraviolet spectroscopy, 32
- crop and geographic distribution, 159
- enzyme-linked immunosorbent assay,
 - detection limits, 168, 169
 - direct competitive assay,
 - overview, 160, 161
 - indirect competitive assay,
 - overview, 160–163
 - materials, 164

- monoclonal antibody assays,
 - 165, 166
- polyclonal antibody assays,
 - 165, 166
- recombinant antibody,
 - competitive direct assay
 - application, 166
 - competitive indirect assay
 - application, 162, 163, 167, 168
 - cross-reactivity compared with monoclonal antibody, 162, 163
 - generation, 162
 - sample preparation, 165
- species in production, 159
- tolerable levels, 160
- toxicity, 159, 160