

Index

2-D DIGE, *see* Two-Dimensional Difference Gel Electrophoresis

A

aCGH, *see* Comparative Genomic Hybridisation, array
Allele-specific primer extension-based genotyping, 54
Antibody microarrays, 199–200

B

Bac-to-Bac system, 270–271
Baculovirus expression systems, 269–271
Basic Local Alignment Search Tool, 43–44, 262, 379, 390, 392–394, 398, 409, 417
BeadArrays, 53–55, 159, 169
BEVS, *see* Baculovirus expression systems
Bioprocessor, 192, 195
BLAST, *see* Basic Local Alignment Search Tool
Bovine serum albumin, 113, 122, 201, 226, 343, 407, 409, 413
BSA, *see* Bovine Serum Albumin

C

cDNA
 first strand synthesis, 151, 154, 411
 second strand synthesis, 154, 156
 microarrays, 181, 211
Cell
 detection on antibody microarray, 203–206
 mammalian, 179, 181, 316, 319, 327–328, 340, 343, 347–348, 404, 406, 410
 COS-7, 354–355, 361, 363
 preparation of suspensions, 202–203

CEPH, *see* Centre d'Etude du Polymorphisme Humain

CGH, *see* Comparative Genomic Hybridisation

Chemiluminescence, 331, 334–336, 356, 363, 414

ChIP-chip, *see* Chromatin

 Immunoprecipitation on a Chip

Chromatin Immunoprecipitation on a Chip, 131, 143

Clinical biospecimens, 88

Co-immunoprecipitation, 291–292, 294, 298, 305, 327, 336

of in vivo cross-linked DNA and proteins, 131–132, 135–136

Comparative Genomic Hybridization, 87
 array, 87, 91–95

Comparative genomics
 prediction of protein function, 387–389

Coomassie blue staining of protein gels, 300–301

CpG islands, 110, 113, 116, 123, 126, 128
 array, 123

 methylation-sensitive cleavage, 112,

 115, 126

 specific adapter, 112, 114, 118

CTCE, *see* Cycling temperature capillary electrophoresis

Cycling temperature capillary electrophoresis, 23

CyDyes, 113, 120, 126, 143, 216, 222–223

D

DCE, *see* Denaturant capillary electrophoresis

Denaturant capillary electrophoresis, 19, 23

DNA

 bacmid, 278–280, 286

DNA (*Cont.*)

Centre d'Etude du Polymorphisme Humain, 5, 12

genomic

- amplification, 1, 6, 10, 143
- copy number determination, 87, 101–102, 106–107, 351
- degraded, 2, 14, 102, 110, 115
- isolation
 - from blood, 1, 2, 6, 77, 81
 - from buccal swabs, 6, 77, 82
 - from tissue, 2, 6, 10, 88, 90

labeling

- aminoallyl and N-hydroxysuccinimide functionalised dyes, 118–120
- random prime, 91

junk, 12

- melting probability calculation, 20, 27
- purification, 9, 272, 330, 333, 351, 355, 361
- quantification

PicoGreen assay, 10, 12, 56

sequence alignment

- multiple, 391
- pairwise, using BLAST, 43–44, 262, 379, 390, 392–394, 398, 409, 417
- pairwise, using Blat, 64, 67, 69

sequence variation

- identification
 - by denaturant capillary electrophoresis, 19, 23
 - by sequence analysis, 35–36, 42, 48
- whole genome amplified, 2, 59, 69

DNA methylation

- analysis, 109–110
- profiling, 109, 113, 115
- patterns, 109, 123–124, 127

DNA-Protein interactions, 131–132

- discovery by chromatin
 - Immunoprecipitation on a Chip, 131–132, 135–136

E

- Electropherogram, 23, 35, 42–44, 46, 148
- Epitope, 132, 141–142, 303, 352, 354, 356–358, 362–365
- Expression vector, 267, 276–277, 329, 336, 341, 353–355, 357–358, 360, 365
 - mammalian, 329, 353, 357, 359

F

- FFPE tissue, *see* Formalin-Fixed, Paraffin-Embedded tissue

Fluorescent dyes, 42, 48, 110, 120, 211–213, 215, 224

- CyDyes, 113, 120, 126, 143, 216, 222–223
 - dye-swap, 95–96, 98

Formalin-Fixed, Paraffin-Embedded tissue

- expression profiling, 159–161, 166, 175
- genomic DNA copy number determination, 101–102, 106–107
- genomic DNA isolation, 103
- RNA isolation, 165

G

GC-clamp, 20–21, 25, 27, 29, 31,

Gene

- expression data
 - integration with protein-protein interaction data, 369–370
- expression profiling, 147, 149, 159–162, 174
- function
 - comparative genomics-based prediction, 387–389
 - prediction using multiple sources of data, 369–370, 380–382
 - sequence homology-based prediction, 387–388
- silencing, 403–405

Genetic mapping

- candidate gene, 75–76, 78
- case-control association study, 76, 83
- linkage analysis, 76, 83
- linkage disequilibrium, 80, 84

Genotyping

- BeadArray-based, 53–55, 57–64
- PCR-based, 15

H

High Performance Liquid Chromatography

- two-dimensional, coupled with mass spectrometry for identification of peptides, 257–258, 261–262, 265–266

HPLC, *see* High Performance Liquid Chromatography

I

- ICAT, *see* Isotope-coded affinity tag-based quantitative protein profiling

IEF, *see* Isoelectric focusing

Immunoblotting, 292, 295, 297, 299, 300, 306, 346

In vitro transcription, 150, 155, 161

Isoelectric focusing, 213, 216–217, 265

Isotope-coded affinity tag-based
quantitative protein profiling,
225–227, 229, 231–233,
238–239

L

Label-free quantitative protein profiling,
241–243, 255–256

Ligation-mediated PCR, 132

LM-PCR, *see* Ligation-mediated PCR

M

Mammalian Two-Hybrid System, 327, 329,
331, 334, 336

Mass spectrometry

compatible Coomassie blue staining,
300–301

compatible silver staining, 300

coupled with liquid chromatography for
detection of differentially
expressed

peptides, 225–227, 229, 233–234, 241,
247–253

tandem, coupled with liquid
chromatography for
identification of peptides,
225–227, 229, 234–237, 242,
253–254, 257–258, 266,
302–303

time-of-flight, in combination with
SELDI, 191–192

Microarrays

cDNA, 181, 211

oligomer or oligonucleotide, 36, 87, 113,
155

microRNAs, 179

expression profiling, 179–186

polyA polymerase 3'-end labelling, 184

Universal Labeling System
labelling, 184

Microsatellites

genotyping, 83

miRNAs, *see* microRNAs

MLPA, *see* Multiplex Ligation-Dependent
Probe Amplification

mRNA

isoforms, 171

MS, *see* Mass spectrometry

MudPIT, *see* Multi-dimensional protein
identification technique

Multi-dimensional protein identification
technique, 264–265, 309, 312,
323–324

Multiplex Ligation-Dependent Probe
Amplification, 101–102,
103–107

Mutation

detection by sequence analysis, 36, 42, 49

N

Non-isotopic quantitative protein profiling,
241–243, 255–256

O

Oligomer or oligonucleotide

microarrays, 36, 87, 113, 155

Open Reading Frame, 353, 355–357, 370

ORF, *see* Open Reading Frame

Orthologues, 390, 391, 395, 398

Ozone

destruct unit, 88

interceptor, 90

level, 98

P

PCA, *see* Principal Component Analysis

PCR, *see* Polymerase Chain Reaction

Peptide

fractionation of complex mixtures
by two-dimensional high
performance

liquid chromatography, 257–258, 261

identification by mass fingerprinting, 191,
196, 221–222

identification by tandem
mass spectrometry, 257–258,
262

Phylogenetic profiling, 395–396

Polymerase Chain Reaction

ligation-mediated, 132

optimisation, 22, 27

primer design, 21, 22, 27, 80–81

quantitative, 134, 136, 142–143,
159, 161, 171–173, 175,
411–412

semi-quantitative, 101–102

sex, 12, 14, 16

thermostable DNA polymerases, 27

Positional gene cloning, 75–76

- Primers
 vector
 T7, 37
 SP6, 37
 sequencing
 T7 universal, 37
- Principal component analysis, 206–207
- Protein
 affinity tag purification, 309–310, 315–320
 blotting, 142, 295, 299–300, 318,
 320–321, 324, 356, 363, 407,
 413–414
 Chip, 191–192
 disease biomarker identification, 191,
 199–200
 expression, 179, 191–192, 211–212, 215,
 238, 242, 255, 354, 361
 in baculovirus, 269–271
 transient, in COS–7 cells, 354
 function
 comparative genomics-based
 prediction, 387–389
 prediction using multiple sources
 of data, 369–370, 380–381
 sequence homology-based prediction,
 387–388
 labelling, 213, 215, 237, 255
 microarray, 199–200
 precipitation, 230, 238, 245, 255, 317
 profiling
 microarray based, 199–200
 quantitative, ICAT and liquid
 chromatography-mass
 spectrometry
 based, 225–227, 229, 231–233, 238–239
 quantitative, liquid chromatography-
 mass spectrometry based,
 241–243, 247–256
 quantitative, two-dimensional
 difference gel electrophoresis
 based, 211–212, 215–216,
 219–221
 SELDI-TOF mass spectrometry based,
 191–192
 purification, 196, 274, 309
 serum, profiling, 191–192
 subcellular localisation, 353–354
 sequence, 43, 48, 79, 238, 255, 260, 262,
 264, 266, 387, 389–391, 399
 tagged, fusion, 353–354
 Protein fragment complementation, 339–340
 Protein-DNA interactions, 131–132
 discovery by chromatin
 Immunoprecipitation on a Chip,
 131–132, 135–136
- Protein-protein complexes
 isolation by co-immunoprecipitation,
 291–293, 297–299, 303
 isolation by tandem affinity purification
 under native conditions,
 309–310, 315–317
 isolation by tandem affinity purification,
 coupled with *in vivo*
 cross-linking,
 under denaturing conditions, 309–312,
 318–320
- Protein-protein interactions
 detection *in vivo* by mammalian two-
 hybrid system, 327–329
 identification by co-immunoprecipitation
 and immunoblotting, 291–293,
 297–298
 identification by co-immunoprecipitation
 and tandem mass spectrometry,
 291–293, 297
 identification by tandem affinity
 purification and tandem mass
 spectrometry, 309–310,
 321–323
 integration of data with gene expression
 data, 369–370
 protein fragment complementation for
 real-time quantitative detection
 in mammalian cell lysates, 344–345
 in live mammalian cells, 345–347
 in mice, 348–350
- Q**
 qPCR, *see* Polymerase Chain Reaction,
 quantitative
- R**
 Reverse transcription, 151, 181
 RISC, *see* RNA-induced silencing complex
 RNA
 integrity, 148, 165
 interference, 179, 403, 409
 isolation, 153–154, 165
 profiling, 147, 149, 159–162, 174
 terminal continuation amplification, 147,
 149–151
 RNAi, *see* RNA, interference
 RNA-induced silencing complex, 177, 404,
- S**
 SDS-PAGE, *see* SDS-Polyacrylamide gel
 electrophoresis

- SDS-Polyacrylamide Gel Electrophoresis, 217–218, 299, 413–414
- SELDI, *see* surface-enhanced laser desorption-ionization
- Sequence
- alignment
 - multiple, 391
 - pairwise, using BLAST, 43–44, 390
 - pairwise, using Blat, 64, 65, 68
 - genome, 35, 53, 64, 66, 69, 76–77, 389
- Sequencing
- artefacts, 47–48
 - primers
 - T7 universal, 37
 - template, 37–38
- Silver staining of protein gels, 300
- Single nucleotide polymorphism, 35–36
- detection
 - by denaturant capillary electrophoresis, 19, 23
 - by sequence analysis, 36, 42, 49
 - genotyping, 54–55
- siRNA, *see* small interfering RNA
- Small interfering RNA, 403–405
- delivery, 410–411
- SNP, *see* Single nucleotide polymorphism
- Split Firefly luciferase reporter fragments for
- protein fragment complementation, 339–341, 343–344
- SRY gene, 12
- Surface-enhanced laser desorption-ionization, 191–192
- T**
- Tandem affinity purification, 309–310, 315–320
- coupled with *in vivo* cross-linking, 309–310, 318–320
- Transcriptional profiling, 147, 149, 159–162, 174
- Transfection
- mammalian cells, 361
 - with small interfering RNA, 410–411
 - insect cells, 278–280
- Transformation, 1, 332, 360
- Two-Dimensional Difference Gel Electrophoresis, 211–212, 215–216, 219–221
- W**
- Western blotting, 142, 295, 299–300, 318, 320–321, 324, 356, 363, 407, 413–414
- WGA, *see* Whole genome amplification
- Whole genome amplification, 1, 2, 6, 10, 14