
Index

- Acetyl phosphate (AP), 4, 22, 39, 60, 65, 76
- Agarose gel electrophoresis,
analysis of PCR products, 85, 88, 100,
161, 176, 218, 261
for purification of PCR products, 168, 172
- Alkaline phosphatase,
expression in oocytes, 127
- Antibody,
for detection and removal of tagged
proteins, 154–155, 161–163
for detection of His-Tagged proteins,
141, 143, 147
for detection of protein expression in
oocytes, 119–120
for immunoprecipitation of in vitro-
expressed proteins, 220
- ATP
cytomim, 6, 8, 10–15
glucose, 10
regeneration (*see also* Energy sources)
PANOxSP, 6, 8, 10–15
- Autoradiography,
of SDS-PAGE, 153, 195
of 2D gel, 195–198, 200–202
- Bacillus anthracis*, 212
- Bacteriophage,
RNA polymerase, 23, 24, 106
- Biotin,
coupled to amino acids, 153, 154, 155,
157
coupled to DNA primers, 160, 162
- Brij 35, 59, 60, 70, 72
- Capping, of RNA, 40–42
- Cassettes,
for autoradiography, 192, 195
DNA, PCR amplification, 217
- Cat gene, 12, 15
- Cesium chloride, for plasmid purification,
215
- Chaperones, 19, 42, 46–48
- CHAPS, 61, 70, 192
- Chloramphenicol acetyl transferase, 12
- Collagenase,
use for oocyte isolation, 111, 114–115
- Continuous-exchange cell-free system
(CECF)
commercial version, 20
concept, 19–20
format, 21
preparation of *E. coli* S30 extract, 26
preparation of wheat germ extract, 31
procedure, using *E. coli* S30 extract, 26–
30
procedure, using wheat germ extract,
31–35
reactors, 21–22, 64
- Creatine kinase, 24, 45, 83, 89, 96, 103,
168, 259, 261
- Creatine phosphate (CP), 4, 24, 39, 45, 83,
89, 96, 97, 168, 259, 262, 263
- Cytomim, 6, 8, 10–15
- Detergents, for membrane protein synthesis,
42, 48, 58, 60, 67, 69–73, 76–77
- Dialysis
buffer, 62, 261, 262, 263, 268
membrane, for cap microreactors, 20–
22
S-30 Extract preparation, 61
- Dialyzers, 21, 261
- Digitonin, 49, 60, 70
- Dihydrofolate reductase,
expression in wheat germ extract, 49
- DNA
template, for transcription reaction, 13–
14, 80–81, 84–88, 100–101, 172–
173, 203, 218, 230, 259
- DNA ligase, 258, 259
- DNA polymerase
LA Taq DNA polymerase, 168, 173,
176
Pfu Turbo DNA polymerase, 82, 86, 87
Pwo DNA polymerase, 190, 194
TaKaRaExTaq DNA polymerase, 82, 85,
96, 100, 101

- Taq DNA Polymerase, 160
 T4 DNA polymerase, 258
 Disulfide isomerase, 42, 46
 Dodecyl- β -D-maltoside (DDM), 61, 70, 72
 Dodecylphosphocholine (DPC), 61, 69, 70
E. coli
 cysteine transporter YfiK (in vitro synthesis), 71
 lipids, 69, 71
 S30 extract, preparation, 26, 61–63
 ECL, 160, 161
 Electrophoresis,
 agarose gel for DNA analysis, 85, 100, 161, 172, 176, 214, 261
 SDS-PAGE, 152, 214, 228, 230
 2D-gel, 195–197
 Elongation factors
 EF-G, 5
 EF-Tu, 5
 Enantioselectivity, 166–167, 177–179
 Energy
 requirement, 5
 sources (for cell-free reactions), 5–11
 classes, 6
 energy-generating pathways, 8–9
 enzymatic reactions, 8
 Enolase, 8
 Enzymatic
 activity, HIV protease, 144, 174–178
 multi-step reactions, 3, 6, 8
 Fluorography, 161, 217, 218, 220, 227
 Folinic acid, 11, 13, 22, 28, 29, 45, 60, 65, 66, 83, 89, 168, 191
 French press, for cell disruption, 24, 26, 59, 62, 75, 189, 193
 Frog
 maintenance, 108
 Nucleoside transporter Tsx (in vitro synthesis), 72–73
 source, 108
 S30 extract, from *E. coli*
 preparation, 61–63
 solutions, 59–60,
 GABA transporter, rat type I (rGAT1), 237, 243, 245
 Galactosidase, conjugated to Fab, 140, 147
 GFP, synthesis, in wheat germ extract, 41, 43
 Glutamine synthase, 257, 270
 Glycerol,
 for bacterial growth, 59
 for 2D-gel, 192
 for enzyme storage, 23, 24
 for transcription and translation reaction, 32–36
 Glycolysis, 4, 10
 GST fusion, 99, 104, 105, 207
 HAART, 135, 136
 HIV, 136
 Human erythropoietin, 55
 Human placenta RNase, 23, 24
 Human proinsulin, 26, 27, 44, 46, 47, 48, 53
 Human telomerase, 48, 55
 Hybridization, of DNA, 159
 Inhibitor
 of oxidative phosphorylation, 10
 of proteases, 23, 60, 65, 112, 135, 136
 of protein synthesis, 4, 7, 39
 of RNase, 24, 60, 111, 258
 Immunoblot, 30
 Immunoprecipitation, 120–121, 156, 220, 226–227
 In vitro transcription and translation
 protein folding (*see also* Chaperones), 5–6, 42–48, 257
 temperature (role), 42–45, 90
 yield evaluation, 14–15
 Kinase,
 acetate, 8
 creatine, 24, 45, 83, 89, 96, 102, 168, 174, 177, 259, 261
 polynucleotide, 259
 pyruvate, 23, 59, 65, 66
 Kozak, consensus sequence, 40, 156, 162, 218
 Library,
 PCR library, 81, 86–88,
 Protein library, 79–81, 88–90, 172–177
 Lipase, *Burkholderia cepacia* KWI-96
 activity, 169, 174,
 gene mutation by PCR, 171–172
 structure (model), 169–170
 LMPG, 61, 69–71
 Luciferase (*Photinus pyralis*), in vitro synthesis, 43, 45, 46
 Magnesium
 concentration, 15, 35, 36, 38

- Membrane proteins (MP), *in vitro* expression,
 as soluble proteins, 72–73
 as insoluble precipitate, 68
 solubilization, 68–72
 strategy, 67
- Microinjection (*see also* Oocyte), 107–109,
 117–118, 123, 127, 236–238, 241,
 243, 244
- Microinjector, 109, 117
- Micromanipulator, 109, 118, 119, 244
- Mitochondrial rhodanese, 48
- Monoclonal antibody, 147
- Mouse, DNA immunization, 223, 228
- mRNA,
 concentration, 33, 38, 39, 40
 preparation, for oocyte injection, 115–117
 quantification, 259, 260
- Mutagenesis, 80, 84, 93, 166, 167, 236, 237
- NMR, protein labeling for studies, 257,
 262–263
- Nonidet P40, 48, 96, 97
- Nucleocapsid, 142, 146
- Octyl glucoside, 48
- Octyl- β -D-glucopyranoside, 60
- Oligodeoxynucleotides, *see* Primers
- Oocyte
 culture media, 110
 microinjection, 117–119
 preparation, 112–114
 protein expression, 119–121
- Oxalic acid, 8
- Phosphoenolpyruvate (PEP), 8, 10, 12, 13,
 15, 22, 28, 29, 39, 66, 76
- Phosphoglycerate, 8
- Phosphoglycerate mutase, 8
- Plasmids
 list of plasmids used
 pCI, 221
 pET, 23, 61, 84, 190, 194, 203, 259
 pET21, 60, 194, 203
 pEU, 98
 pEU3b, 259
 pIVEX, 37
 pK7CAT, 12–13
 preparation, 23, 146, 190, 221–223, 232
 pRSET, 167
 pSP64T, 126
 pXO1, 212
 storage, 23
- Polyadenylation, of RNA, 40–42
- Polyamines, 10
- Polymerase chain reaction (PCR)
 isolation of *Bacillus anthracis* genes (*see also* *B. anthracis*), 218
 cloning of PCR products, 203
 concept, 81
 mutation of lipase gene (*see also* Lipase), 170–171
 preparation of SM-PCR libraries, 86–88
 purification of amplified fragments, 116,
 229
 single-molecule PCR (SM-PCR)
 templates preparation, 85
 use of PCR fragments for *in vitro*
 transcription/translation, 40, 88–
 90, 98–105, 161, 173–174, 176–
 177, 203, 218–220, 230–231
- Potassium (K⁺) concentration, 38, 61, 64,
 65, 66, 76
- Primers,
 for amplifying *B. anthracis* genes, 218
 for amplifying *N. meningitidis* genes,
 202
 for lipase gene mutagenesis, 167–168,
 171–172
 for PTT analysis, 161–162
 for single molecule (SM) PCR, 84–86
 for tagging PCR products, 154–155
 split primers for direct transcription and
 translation in wheat germ, 100–102
- Proinsulin (human), *in vitro* synthesis, 27,
 44
- Protease
 activity, 144
 HIV protease (PR), 136
 inhibitor, 136
 substrate, 140
- Proteoliposomes, 72
- PTT (Protein Truncation Test)
 protocol, 160–163
 strategy, 158–159
- Purification of
in vitro synthesized proteins, 105
 PCR products, 116, 229
 plasmid DNA, 23, 221–223, 267

- Puromycin, 92
 Putrescine, 8, 10, 12, 13
 Pyruvate kinase, 23, 29, 30, 59, 65, 66
 Redox potential, 42
 Reverse transcriptase, 142
 Ribonuclease inhibitor
 from human placenta, 24
 porcine, 607
 RNA,
 preparation for oocyte injection, 115
 RNA polymerase,
 T7, *see* T7 RNA polymerase
 SP6, *see* SP6 RNA polymerase
 rRNA, 18S, in wheat germ, 42
- SDS-PAGE, *see* Electrophoresis
 Sephadex chromatography, 39, 96, 97
 Sepharose
 A sepharose beads, 220, 227
 A sepharose CL-4B, 112, 120, 215
 glutathione sepharose, 97, 105
 Q sepharose, 74
 Solubility of *in vitro* translated proteins, 44,
 46, 47, 48, 72
 SP6 RNA polymerase, 96, 103, 258, 262,
 267
 Spermidine, 8, 10, 12, 13, 24, 25, 33, 35, 36,
 96, 97, 215, 223, 258, 261–263
 Strains
 Bacillus anthracis Vollum, 218
 E. coli A19, 75
 E. coli BL21 DE3, 179
 E. coli JM109, 259
 E. coli MRE 600, 23
 Neisseria meningitidis MC58, 193, 194
 Streptavidin,
 coupled to magnetic beads for protein
 purification, 160–161
 Supercoiled DNA, 222, 232
 T4 DNA polymerase, 258
 T4 DNA ligase, 258
 T7 RNA polymerase
 purification, 12, 74
 use, 13–14, 27–29, 35–36, 58, 59, 61,
 65, 74, 82, 88, 89, 126, 168, 171,
 177, 205, 218
- Thesit, 60, 70
 TEVC (Two Electrode Voltage Clamp),
 236–237, 239, 242–243, 246–249,
 252
 Transaminase
 Triton X-100, 48, 60, 70, 72, 73, 112, 168,
 169, 174, 177, 214
 Tumor, diagnosis, 149, 152, 153, 157
 Tween-20, 61, 70, 161
 Untranslated regions (UTR) (*see also*
 mRNA)
 from alfalfa mosaic virus, 126
 for optimizing expression in wheat germ
 extract, 41, 42, 98
 Ubiquitin, yeast
 labeling in wheat germ extract for NMR
 analysis, 261
 mRNA preparation and quantitation, 260
 Vectors, *see also* Plasmids
 for DNA vaccination, 221
 for expression of HIV genes, 142
 Wheat-germ
 preparation of DNA template for
 transcription/translation reaction
 (*see also* PCR), 98–102
 preparation of embryos, 31, 97
 preparation of extract, 31, 97–98
 preparation of mRNA template for
 translation reaction (*see also*
 PCR), 40–42
- Xenopus levis*, *see also* Frog
 oocyte injection, 117–119
 oocyte preparation, 112
 oocyte protein expression, 119
 preparation of protein extracts from
 oocytes, 120
 source and maintenance, 108
 Yeast,
 cell-surface display, 80
 extract for bacterial growth, 59, 215
 tRNA, 24, 33
 ubiquitin,
 in vitro expression in wheat germ
 extract, 261
 isotope labeling, 262