

Note: The letters 't', 'f' and 'n' following the locators refer to tables, figures and note numbers respectively.

A

- Adenine sulfate56, 57, 58, 257
- Affymetrix
GeneChip system.....294
Technical Manual.....301
- Agarose
gel electrophoresis.....88, 122, 123, 128, 131, 213, 255,
262, 282 n5, 300, 303 n8, 344
plug formation and cell lysis168, 171
- Alkaline transfer buffer217, 227
- Allele-specific oligonucleotide (ASO).....328, 329, 338,
345–350, 346f, 356
- Allele-specific primers316, 319, 325f, 332–334, 336,
337, 339, 344f, 346f, 352 n2
primers used in crossover assays.....332t
- Anaphase I stage of meiosis.....77
- Arabidopsis thaliana*, detecting gene conversion in gametes of
backcrossing nonfluorescent alleles.....108–109
DNA preparation
from nonfluorescent mutant107–108
for sequencing104
- EMS mutagenesis of fluorescent transgenic
lines105–106
- heterozygous alleles, segregation of.....102f
- M1 FTLs for nonfluorescent alleles,
screening106–107
- microscopy104
- mutagenesis104
- nonfluorescent allele sequencing108
- plant material/growth104
- scoring GC events in the pollen tetrads110
- using GC test locus for mutant analysis110–112

B

- Background normalization160f
regions used for161t
- Benzoyl-naphthoyl DEAE (BND) cellulose.....144, 145f,
149, 154, 155, 160, 162
- β -mercaptoethanol (BME).....187, 190
- Binucleate cells.....272, 280 n3
- Biotinylated ASOs used in DEASH experiments347t
- Bivalent36, 37f, 80, 90, 91

- Bovine serum albumin (BSA).....123, 168, 170,
171, 199, 238, 248, 259, 270, 272, 282, 287,
289, 290, 297, 299, 301, 307, 308, 327, 328,
329, 348, 349

- Budding yeast (*S. cerevisiae*).....260

C

- C. elegans*, meiotic recombination analysis, *see* Meiotic
recombination analysis, *C. elegans*
- Cell lysis efficiency.....194 n5
- Cells, synchronization for entry into meiosis
liquid sporulation medium (SPM)58
synthetic complete (SC) powder58
yeast strains and genotypes57t
- Chiasma/Chiasmata.....36, 77–78, 79, 80, 81, 90, 92, 96
n17, 305
- ChIP, *see* Chromatin immunoprecipitation (ChIP)
- ChIP-chip
analysis31
protocol291, 292f
technique.....269
technology.....286
- ChIP grade antibody287
- ChIP on Chip (ChIPChip).....268
- Chi-Square distribution.....46
- Chi-Square “Goodness of Fit” test43
- Chi-Square test.....41, 43
- Chromatin analyses.....264f
- Chromatin immunoprecipitation (ChIP).....267–283, 285,
286, 291
- Chromatin structure at meiotic DSB sites in yeasts.....264f
cell culture
S. cerevisiae256–257
S. pombe257–258, 261
crude chromatin from *S. cerevisiae* cells and MNase
digestion.....260–261
gel electrophoresis
southern blotting analysis259
transfer to nylon membrane.....262–263, 263f
hybridization/detection.....264–265, 264f
isolation of chromatin fraction
from *S. cerevisiae* and MNase treatment.....258
from *S. pombe* cells and MNase digestion262
probe preparation.....263–264

- Chromatin structure (*continued*)
 probe/restriction site to analyze DSB hotspots263^f
 procedure (protocol)255–256, 255^f
 recombination hotspots, regulation of.....254–255
 synchronous sporulation of *S. cerevisiae*.....260
- Chromosome interference, *see* Methods for analysis of
 crossover interference, *S. cerevisiae*
- Chromosome-level DSB mapping,
S. cerevisiae121–123, 125–128
 hybridization and wash128
 PFGE.....122, 126–127
 plug DNA preparation121
 preparation of agarose-embedded DNA (“plugs”)....126
 southern blot hybridization/washing.....122–123
 southern blotting.....122, 127–128
- Coefficient of coincidence36, 42–43, 44, 45
 at varying linkage map distances.....45^f
- Colony-forming unit (C.F.U.)59, 60
- Counting model.....37, 41, 42, 45, 46,
 47, 48, 49
- Cre/loxP61
- Cre recombinase56, 58, 60
- Crosslink Reversal Buffer219, 230, 231
- Crossover interference in *S. cerevisiae*, methods for
 analysis of
 cytological and genetical methods36
 data collection38
 four-factor data45–46
 interference in meiotic mutants48–49
 detection of significant change49
 multifactor data and models46–48
 counting model to yeast, complications.....48
 densely spaced markers47–48
 sparse genetic markers47
 recording, organizing and evaluating data.....38
 three-factor data.....42–45
 random spore data.....42–43
 tetrad data for chromatid interference.....43–44
 tetrad-specific “coefficient of coincidence” for
 chromosome interference.....44
 tetrad test for chromosome interference.....44–45
 two-factor data.....38–42
 detecting and quantifying interference via
 internet.....42
 improvised two-factor test39–41
 NPD ratio39
 quantifying interference.....41
- Crossovers
 breakpoint mapping, principle of340^f
 frequencies for each SNP interval, calculating343^t
 and nonexchange gene conversions by the half-crossover
 assay, detecting.....344^f
 scoring, example of341–342
 tetrads.....44
- Crossovers/noncrossovers in mouse germ cells
 CO and NCO at the Psmb9 hotspot in B10×R209
 ovaries, parallel detection.....309^f
 detection of specific molecules by allele-specific nested
 PCRs.....311–312
 determination of
 CO frequency.....313, 316 n2
 concentration of amplifiable molecules312
 frequencies of CO/NCO in B10×R209 oocytes315^t
 frequencies of recombinant products,
 calculation314–315
 mapping recombination breakpoints315–316
 nested PCR procedure.....311^f
 parallel analysis of CO/NCO313–314
 preparation of DNA307, 310–311
 preparation of ovary samples...307, 308–310, 309^f, 310^f
 primers used for detection of
 recombinant products at Psmb9 hotspot in
 B10×R209 hybrid317^t
 reaction mixes for nested PCR.....308
- Culture media21, 67, 139 n1, 255, 256
- Custom primers169
- CyTM3/CyTM5 monoreactive dye pack150
- D**
- Data collection, interference analysis.....38
- DEASH, *see* DNA Enrichment by Allele-Specific
 Hybridization (DEASH)
- DEASH and detection of recombinants, principle of..319,
 345, 346^f, 347, 351
- Densely spaced markers47–48
- Diakinesis stage of meiosis prophase I77–78
- Dithiothreitol (DTT)168, 187, 198, 199, 288, 328
- D-loop.....100
- DNA
 binding.....16, 28, 29, 31, 267, 268
 breaks in testis cells from strains 9R (panel A) and 13R
 (panel B), detection of174, 174^f
 breaks (rare), method to detect.....167, 167^f
 double-strand break27, 65, 77, 91, 100, 165, 183,
 236, 306, 326
 extraction.....169, 172
 from a meiotic induction, analysis239
 mismatch repair system.....101
 –protein interaction285
 recombination intermediates, *S. pombe*
 cell culture236–237
 cell culture for meiotic induction.....242–243
 crossover products, gel electrophoresis245–246
 2D branched DNA intermediaries246–247
 DSB analysis, gel electrophoresis for.....244–245
 gel electrophoresis.....238, 243–244
 meiotic induction/DNA
 extraction.....237–238, 243–244

- southern blotting/hybridization/analysis 238–239
See also S. pombe, DNA recombination intermediates
- DNA Enrichment by Allele-Specific Hybridization
 (DEASH) 319 n14
- Double-Holliday junction intermediate 100
- Double-strand break (DSB) 4, 8, 10*f*, 65, 100, 119–141,
 209, 269
 frequencies 28
Drosophila 37*f*, 42, 45, 46
- DSB, *see* Double-strand break (DSB)
- DSB-cold 145
- DSB formation
 reduction methods
 chromatin-associated factors, modification of 28
cis-acting element, deletion of 28
 inducing transcription 28
 stimulation methods
 insertion/substitution of foreign DNA
 fragments 28–29
- E**
- Ea* gene 166, 167, 174, 177 n1
- EKRB (Enriched Krebs Ringer Buffer) 168
- EKRB-CO2-BSA buffer 168, 170
- EMS, *see* Ethylmethane sulfonate (EMS)
- End-labeling
 and analysis of Spo11 oligonucleotide complexes
 in *S. cerevisiae*, *see* SPO11-oligonucleotide
 complexes
 indirect 126, 256, 262, 263
- Endonucleolytic release of Spo11 from DSB ends 185*f*
- Enriched Krebs Ringer Buffer 168
- Equilibration Buffer 219, 230, 231
- Erlang, Agner Krarup 46
- Erlang distribution 37*f*, 45*f*, 47
- Ethidium bromide running buffer 228
- Ethylmethane sulfonate (EMS) 102, 104, 105, 106, 112
 n2, 113 n7
- Eukaryotes 27, 77, 306
- Eukaryotic DNA 253
- Eukaryotic hybridization control kit 289
- External environment, *S. cerevisiae* meiosis
 nutrient composition 10–11
 temperature 8–10
- F**
- Fisher Exact test 43
- Fission yeast 65, 165, 166, 260
- Fluorescence microscopy 59, 61, 102, 112, 220
- Fluorescenttagged lines (FTLs) 102, 103*f*
- Four-factor data 38, 45–46
- FTLs, *see* Fluorescenttagged lines (FTLs)
- G**
- Gal4 DNA binding domain (Gal4BD) 29–31
 DSB targeting by 30
 GAL2 promoter 30*f*
- Gamma distribution 46, 47
- GCOS, *see* GeneChip Operating Software (GCOS)
- GC tetrad frequency 110, 111
- Gel electrophoresis assays, DNA DSB in *S. cerevisiae*
 chromosome- level DSB mapping 121–123, 125–128
 hybridization and wash 128
 PFGE 122, 126–127
 plug DNA preparation 121
 preparation of agarose-embedded DNA
 (“plugs”) 126
 southern blotting 122–123, 127–128
 high-resolution mapping 124, 131–132
 denaturing PAGE gel electrophoresis and semi dry
 transfer 131–132
 PCR-mediated probe labeling 132
 polyacrylamide gel electrophoresis 124
 southern blotting 124
 media for yeast culture 121
 medium- resolution mapping 123, 128–131
 agarose gel electrophoresis 123
 genomic DNA extraction 123, 129–130
 quantification of DSB frequency 130–131
 restriction enzyme digestion and 130
 southern blotting 123, 130
 nucleotide resolution mapping 124–125, 132–138
 denaturing polyacrylamide gel
 electrophoresis 135–136
 electrophoresis 124
 preparation of probes/sequence standards/
 samples 134–135
 probe preparation and hybridization 125
 sample preparation 124
 southern blotting 125, 136–137
 strand-specific probe preparation 136–138
 synchronous meiotic cultures 125
- GeneChip Operating Software (GCOS) 301
- Gene conversion (GC) 99
 LD mapping 100
- Genetic heterogeneity, *S. cerevisiae* meiosis 5–8
 strain backgrounds 5–6
 msh4 mutation in 6*f*
- Genetic mapping techniques 78
- Genetic markers 5, 30, 47, 78, 82, 86, 87, 92, 93*f*
- Genome-wide high-resolution CHIP of meiotic
 chromosomal proteins in *S. cerevisiae*
 amplification of DNA 288–289, 299–300
 cell culture/*inVivo* crosslinking 286, 295–296
 cell lysis/fragmentation of genomic DNA 287,
 296–297

Genome-wide high-resolution ChIP (*continued*)

- ChIP-chip protocol 292^f
 - fragmentation/end-labeling 289, 300
 - hybridization 289, 300–301
 - Immunoprecipitation (IP) 287–288
 - oligonucleotide tiling array, design of 293
 - preparation of antibody
 - binding beads 297
 - conjugated beads 287
 - preparation of DNA 288, 298–299
 - washing/staining/scanning 290, 301–302
- Genome-wide mapping of meiotic DNA DSB in
S. cerevisiae
- amplification/labeling of ChIP-associated sequences
and ssDNA enrichment material by random
primer extension 155–157
 - background normalization of ChIP-chip and
ssDNA-chip microarray data 159–161, 160^f
 - batch enrichment of ssDNA on BND
cellulose 148–149, 154–155
 - high-density microarray analyses of DSB-associated
sequences 150–151, 157–159
 - meiotic cultures 146–148, 151
 - meiotic DNA preparation by phenol chloroform
extraction 148, 153–154
 - quantitative PCR of DSB-associated sequences 149
 - regions used for background normalization 161^t
 - Spo11 ChIP in break processing defective
mutants 148, 152–153

H

- Haploid parents, mass matings of 49 n1
- Hawaiian strain 86, 95 n10
- Heteroduplex DNA 100, 101, 213, 214, 315, 320 n18
- Heterozygous mice, assay for detecting meiotic CO and
NCO at another locus in 316 n2
- High-resolution mapping, *S. cerevisiae* 124, 131–132
 - denaturing PAGE gel electrophoresis and semi dry
transfer 131–132
 - PCR-mediated probe labeling 132
 - polyacrylamide gel electrophoresis 124
 - southern blotting 124
- HIS4LEU2 physical assay system 210, 210^f
- Holliday junction stabilization and detection on 2-d
gels 211^f
- Homologous recombination 144, 209, 236
- Hotspots 15, 48, 118, 141, 146, 159, 160,
239, 254, 259, 260, 263^f; 306, 316, 324, 325^f,
336, 353
- Human sperm, analysis of meiotic recombination
 - allele-specific PCR to detect crossover
molecules 334–335
 - amplification 334–337
 - designing, primers 332

- determining linkage phase of donors 333–334
- optimizing, primers 333
- primers used in assays, examples of 332^t
- crossover
 - breakpoint mapping, principle of 340, 340^f
 - frequencies for each SNP interval,
calculating 343^t
 - scoring, example of 341–342
- DEASH 328–329
 - DNA enrichment by 345–352
and subsequent detection of recombinants,
principle of 346^f
- DNA extractions 326–327, 329–332
 - from blood 331
 - quantification of genomic DNA 331
 - from semen 330–331
- gene conversions/crossovers using half-crossover
assay 323–345
- linkage phasing, principle of 334^f
- mapping crossover breakpoints by ASO
 - hybridization 328
- mapping locations of crossovers 327
 - ASO hybridization 338–339
 - mapping crossover breakpoints 338–339
 - third round of PCR 337
 - transferring reactions onto dotblots 337
- optimal yield/degree of purification post-DEASH,
determining 350^f
- recipe for 11.1× PCR buffer 327^t

I

- Indirect end labeling 126, 256, 262, 263
- Interference
 - definition 35
 - indices of 49
 - in meiotic mutants 48–49
 - detection of significant change 49
 - map-length/interference phenotype combinations,
ambiguity 49
- Intragenic recombination 75
- Isogenic strains 8

J

- Joint molecule (JM) 209–211, 210^f; 213,
230, 236

L

- Labeling assay, demonstrating sensitivity of 192^f
- “Late nodules” 36, 77, 79, 90, 92, 305
- LAT52 (pollen-specific promoter) 102, 104, 112 n3,
113 n9
- Linkage disequilibrium (LD) mapping 100
- Linkage phasing, principle of 334^f

- Linker DNA254, 256
- Lysis buffer80, 87, 89, 90, 148, 152, 153, 168, 171,
198, 199, 237, 243, 257, 258, 261, 260, 270, 272,
273, 287, 296, 298, 307, 310
- M**
- Mass matings of haploid parents49 n1
- M9 buffer80, 91, 96 n16
- Media, *S. cerevisiae* meiosis
liquid 14
solid 11–14
- Medium-resolution mapping, *S. cerevisiae* 123, 128–131
agarose gel electrophoresis 123
genomic DNA extraction 123, 129–130
quantification of DSB frequency 130–131
restriction enzyme digestion and electrophoresis 130
southern blotting 123, 130
- Meiosis, distribution of exchanges among acts of37f
- Meiosis, *S. cerevisiae* 3–5
sources of variability
external environment, *see* External environment,
S. cerevisiae meiosis
genetic heterogeneity, *see* Genetic heterogeneity,
S. cerevisiae meiosis
intracellular environment 14–15
media, *see* Media, *S. cerevisiae* meiosis
- Meiotic divisions, progression of281
- Meiotic DNA breaks in mouse testicular germ cells
agarose plug formation and cell lysis 168, 171
analysis of gel electrophoresis results176
DNA breaks (rare), method to detect 167f
DNA extraction 169, 172
primer extension and PCR 172–173
of dGTP-Tailed DNA fragments 169
primer labeling for PCR 169
rare DNA breaks, method to detect 167
restriction enzyme digestion
of PCR product/detection of DNA breaks.. 169–170
of second-round PCR products and detection
of DNA breaks 173–176
fluorescence-based detection of DNA
breaks 174–176, 175f
radioactive detection of DNA breaks 173–174
terminal deoxynucleotide transferase (TdT)
tailing 168–169, 171–172
testis dissection/germ cell preparation 170–171
- Meiotic recombination analysis, *C. elegans*
determination of chiasma frequency 80, 90–91
measuring crossing over using recessive phenotypic
markers 79–80, 81–85
males 85
during spermatogenesis/oogenesis in
hermaphrodites 82–83
- measuring crossovers over using snip-SNP
markers 86–90
in males 89–90
during oogenesis in hermaphrodites 88–89
during spermatogenesis and oogenesis in
hermaphrodites 87–88
through ionizing irradiation 81, 91–92
- Meiotic recombination analysis, human sperm, *see* Human
sperm, analysis of meiotic recombination
- Mendelian segregation 102
- MESA Green 274
- Microarray Suite 5.0 (MAS5) 301
- Micrococcal nuclease (MNase) 254, 258
- Mismatch (MM) probe 294
- Molecular cloning 206 n13
See also Mouse
- Morphological markers 78
- Mouse
DSBs 166
germ cells, crossovers/noncrossovers in, *see* Crossovers/
noncrossovers in mouse germ cells
Psmb9 hotspot 306
SPO11-oligonucleotide complexes, labeling of 202
testicular germ cells, meiotic DNA breaks in, *see* Meiotic
DNA breaks in mouse testicular germ cells
- Mre11 184, 268–269, 277–278
- Multifactor data and models 46–48
counting model application to yeast, complications... 48
heterogeneity of crossover process 48
densely spaced markers 47–48
sparse genetic markers 47
- M value calculation from tetrad data, *see* Stahl Lab Online
Tools
- N**
- NanoDrop spectrophotometer 157 n9
- Noncrossovers 4, 46, 47, 100, 101, 143,
305, 345
- Nonparental ditypes (NPDs) 38
- NPD ratio 36, 39
- Nucleosome 253, 254, 255, 256
- Nucleotide resolution mapping,
S. cerevisiae 124–125, 132–138
denaturing polyacrylamide gel electrophoresis... 135–136
of DSB at the YCR048w promoter region 138f
electrophoresis 124
preparation of probes/sequence standards/
samples 134–135
probe preparation/hybridization 125, 136–138
sample preparation 124
southern blotting 125, 136–137
strand-specific probe preparation
and hybridization 137–138

O

- Oligonucleotide tiling array, design of 293^f
- Optical filters 112 n4
- Optimal antibody concentration 195
- Optimal spheroplasting 162 n8
- Optimal yield and degree of purification post-DEASH,
determining 350^f
- Outcross progeny identification 94 n6
 - classes 84
 - miscounting of 94 n7
- Owen's model 47

P

- Papazian 39
- Parental ditypes (PDs) 38
- PCR, primer extension
 - dGTP-tailed DNA fragments 169
- PCR, primer extension and 172–173
- PCR, primer labeling for 169
- PCR (qPCR) 146
- Perfect match (PM) probe 294
- Perkins's equation 44
- PFGE, *see* Pulsed-Field Gel Electrophoresis (PFGE)
- PFGE for DSB analysis
 - 20 kb–1 Mb DNA fragments 244–245
 - whole chromosomes 244
- Poisson distribution 36, 46
 - Poisson law 320 n18
- Polyacrylamide gel electrophoresis 124
- Positive interference 35–36
- Potassium phosphate buffer 79, 80, 87
- Pre-sporulation medium (SPS) 269
- Primers 80, 177 n1
 - gene amplification 112 n3
- Probe preparation and microarray choice 145
- Promoter (pGPD-loxP) and reporter constructs
(loxP-reporter), orientation and location 56^f
- Proteinase K 80, 104, 121, 123, 129, 133, 148,
153, 168, 215, 216, 217, 222, 223, 237, 243,
244, 261, 262, 270, 273, 288, 302 n4, 307,
326, 330, 331
- Protein–DNA interactions during meiosis by qChIP
 - chromatin immunoprecipitation 272–273
 - DNA purification 270, 273–274
 - immunoprecipitation 270
 - interpretation of qChIP results 275–279
 - Mre11-HA interaction with meiotic chromosomes 269
 - Mre11 interacts transiently with hotspot/cohesin
 - binding site in meiosis 278^f; 279^f
 - preparation of cell extracts 270
 - qChIP, quantitative 267–269
 - real-time PCR
 - amplification chart 276^f

- analysis 274–275
 - quantitative analysis 271
 - synchronized yeast sporulation 271–272
 - yeast culture/sample collection 269–270
 - yeast extract preparation 272
- Psoralen cross-linking 250 n6, 251 n11
- Pulsed-Field Gel Electrophoresis (PFGE) 122

Q

- qChIP, *see* Quantitative chromatin Immunoprecipitation
(qChIP)
- qPCR (quantitative, real-time PCR) 146, 153, 155,
163 n9, 267, 268, 269, 274–278, 283 n11, n10
- QQR 31
- Quantitative chromatin Immunoprecipitation
(qChIP) 267–283

R

- RAD-51 protein/DNA complexes 92 n1
- rad50S*-like mutants 144
- Random-priming kit 263
- Random spore analysis, *S. pombe*
 - intragenic (gene conversion) recombinant
 - frequency 71–72
 - spore viability 71
 - spore yield viability 71
- Rare DNA breaks, method to detect 167
- Rate parameter (λ), 46–47
- Recombination frequency calculation 83, 88
- Recording, organizing and evaluating data 38
- Reporter genes 56
- Restriction enzyme digestion
 - PCR product and detection of DNA
 - breaks 169–170
 - second-round PCR products and detection of DNA
 - breaks 173–176
 - fluorescence-based detection of DNA
 - breaks 174–176
 - radioactive detection of DNA breaks 173–174
- See also* Meiotic DNA breaks in mouse testicular germ
cells
- Robust quantile normalization 303 n5
- Rosy locus 46

S

- S. pombe*, DNA recombination intermediates
 - cell culture 236–237
 - cell culture for meiotic induction 242–243
 - conditions for pat1+ strains 242–243
 - conditions for pat1-114 strains 242
 - gel electrophoresis 238, 243–244
 - crossover DNA products 245–246

- 2D electrophoresis of branched DNA intermediaries.....246–247
- for DSB analysis244–245
- meiotic induction/DNA extraction237–238, 243–244
- southern blotting/hybridization/analysis238–239
- S. pombe*, genetic analysis of meiotic recombination
 - cell growth and crosses.....70
 - culture media.....67–68
 - life cycle of66–67
 - meiosis and viable spore yield on plates, analysis.....69–70
 - meiotic tetrad analysis.....72–73
 - random spore analysis70–72
 - spore preparation and analysis68
- S. pombe* pat1-114 haploid or diploid cells236
- Saline-sodium phosphate-EDTA buffer (SSPE)289
- S. cerevisiae*
 - gel electrophoresis assays, DNA DSB in, *see* Gel electrophoresis assays, DNA DSB in *S. cerevisiae*
 - genome-wide high-resolution chromatin immunoprecipitation of meiotic chromosomal proteins in
 - genome-wide mapping of meiotic DNA double-strand breaks in
 - high-resolution mapping, *see* High-resolution mapping, *S. cerevisiae*
 - medium-resolution mapping, *see* Medium-resolution mapping, *S. cerevisiae*
 - meiosis.....3–5
 - external environment, *see* External environment, *S. cerevisiae* meiosis
 - genetic heterogeneity, *see* Genetic heterogeneity, *S. cerevisiae* meiosis
 - intracellular environment14–15
 - media, *see* Media, *S. cerevisiae* meiosis
 - meiosis, Media, *see* Media, *S. cerevisiae* meiosis
 - methods for analysis of crossover interference
 - cytological and genetical methods36
 - data collection38
 - four-factor data45–46
 - interference in meiotic mutants48–49
 - multifactor data and models46–48
 - recording, organizing and evaluating data.....38
 - three-factor data.....42–45
 - two-factor data.....38–42
 - modulating and targeting meiotic DSB
 - local modification of Spo11-dependent DSB
 - frequencies28–29
 - methods to reduce DSB formation28
 - methods to stimulate DSB formation28–29
 - methods to globally modify Spo11-dependent DSB frequencies29
 - site-specific targeting of Spo11 DSBs.....29–31
 - DSB targeting by Gal4BD-Spo1130–31, 30^f
 - targeting Spo11 with the artificial QQR zinc finger array31, 32^f
 - yeast strains31–32
 - optimizing sporulation conditions
 - pre-sporulation media.....24–25
 - results22
 - sporulation media.....25
 - standardized test protocol in liquid/plates.....25
 - spatial proximity/accessibility of chromosomal loci using Cre/loxP site-specific recombination
 - cre induction and return-to-growth58, 59–60
 - method analysis.....60–61
 - synchronization of cells for entry into meiosis.....56–58, 59
 - SDS extraction buffer189, 194 n5
 - SeaKem Gold agarose.....232 n9
 - Sequencing gel loading buffer199, 203, 206 n12
 - Sex chromosomes of *Drosophila*45^f
 - Shape parameter (ν).....46–47
 - Singer instruments.....38
 - Single-end invasion intermediate (SEI).....209
 - Single-strand DNA (ssDNA)144, 145, 148, 149, 153, 154, 155, 159, 160^f, 162
 - Single-Worm Lysis buffer.....80
 - Snip-SNP markers.....83
 - measuring crossovers during spermatogenesis and oogenesis in hermaphrodites using.....87–88
 - PCR analysis87
 - Sonic ZD-dismembrator58
 - Sparse genetic markers.....47
 - Sperm DNA, amplification of crossover molecules from
 - bulk recovery of crossover molecules337
 - calculating crossover frequencies336–337, 336^t
 - first round of PCR.....335
 - reciprocal crossover assays.....336, 337
 - second round of PCR336
 - S1 nuclease digestion335–336
 - Sperm typing.....306, 316, 324
 - Spheroplast buffer...147, 148, 153, 162, 215, 222, 237, 243
 - Spo11
 - DSBs, site-specific targeting of29–31
 - targeting by Gal4BD-Spo1130–31, 30^f
 - targeting Spo11 with the artificial QQR zinc finger array.....31, 32^f
 - enzyme77
 - FLAG/Mre11-FLAG/Rec8-FLAG during meiosis
 - on chromosome V of budding yeast, distribution of295^f
 - HA3His6, 195 n13
 - oligonucleotide complexes, detection of labeled mouse198, 201, 202^f, 205 n8, n9
 - protein27
 - by western blotting, detection of199, 201–202, 203^f

- Spo11 (*continued*)
 detected by Western blotting.....203*f*
See also Mouse
- SPO11-oligonucleotide complexes
 -associated oligonucleotides
 determining length of.....199, 202–204
 in wild-type mice, size of.....204*f*
 -dependent DSB frequencies
 local modification of.....28–29
 methods to reduce DSB formation.....28
 methods to stimulate DSB formation.....28–29
 methods to globally modify.....29
 in mouse testes
 detection of labeled mouse SPO11-oligonucleotide
 complexes.....198, 201, 202*f*, 205 n8, n9
 detection of SPO11 protein by western
 blotting.....199, 201–202, 203*f*
 determining length of SPO11- associated
 oligonucleotides.....199, 202–204
 immunoprecipitation.....198, 200
 preparation of testis lysates.....198, 199–200
 TdT labeling.....198, 200–201
 in *S. cerevisiae*
 from meiotic yeast cells, scheme for detecting ..186*f*
 potential problems.....192–193, 193*f*
 preparation of denatured yeast cell lysate.....187
 cell lysis in TCA.....189
 preparing cells for synchronous yeast
 meiosis.....188–189
 protein extraction from denatured yeast
 lysate.....189–190
 in *S. cerevisiae*, detection of.....194 n8
 in *S. cerevisiae*, end-labeling of.....187–188
 immunoprecipitation of Spo11-Oligo complexes...190
 sampling multiple time points.....192
 using TdT.....190–191
 western detection of Spo11 (Optional).....191–192
 in *S. cerevisiae*, end-labeling of detection.....191, 192*f*
- Spore viability.....29
- Sporulation.....21–22
- Sporulation efficiency
 different strains tested for.....22*t*
 effect of temperature.....22, 24*t*
 cold-sensitive.....22, 24*t*
 in liquid culture.....23*t*
 on plates.....23*t*
- Sporulation media, supplements for.....147*t*
- Sporulation medium (KAc).....147, 214
- Sporulation medium (SPM).....58, 119, 269, 270, 286
- Stabilization/electrophoretic analysis of meiotic
 recombination intermediates in *S. cerevisiae*
 DNA extraction
 guanidine/sarkosyl preparation of Psoralen cross-
 linked DNA.....215, 222–223
 yeast DNA preparation in agarose
 plugs.....216–217, 225
 yeast DNA preparation with
 CTAB.....215–216, 223–225
 DNA purification.....211–212
 preventing holliday junction isomerization and
 branch migration.....211*f*, 212
 Psoralen crosslinking.....212
 electrophoresis
 1-d agarose gels to detect JMs.....217, 226–227
 2d native/denaturing
 electrophoresis.....217–218, 228–229
 1d native gel electrophoresis.....218, 229
 2-d native/native gel electrophoresis....218, 229–230
 2-d native/native to detect JMs.....217, 217*f*,
 227–228
 removal of Psoralen cross-links and 2d native/
 denaturing electrophoresis.....219, 230–231
 JM detection and analysis using agarose gel
 electrophoresis.....213, 214*f*
 pregrowth and sporulation.....219–220, 221
 restriction enzyme digestion
 Psoralen cross-linked DNA.....225–226
 yeast DNA preparation in agarose plugs.....226
 yeast DNA prepared with CTAB.....226
 sporulation in 1% potassium acetate.....213–214
 Stahl Lab Online Tools.....39, 41, 42, 42*f*, 45
 Strain backgrounds.....5–6
 BR.....5–6
 S288c.....5–6
 selection basis.....6
 SK1.....5–6
 Y55.....5–6
 Strand-exchange.....209
 SYBR Green.....274
 Synchronous meiotic cultures.....125
 Synthesis-dependent strand annealing.....100
- T**
- Tailing buffer.....168, 171, 172
- TBE electrophoresis buffer.....122
- TBS, *see* Tris-buffered saline (TBS)
- TdT, *see* Terminal deoxynucleotide transferase (TdT);
 Terminal transferase (TdT)
- TdT labeling.....198, 200–201
- TE buffer.....168
- Terminal deoxynucleotide transferase (TdT).....168, 169,
 171, 185, 188, 197, 198, 289, 291, 300
- Terminal transferase (TdT).....197–198, 289, 291, 300
 tailing.....168–169
See also Terminal deoxynucleotide transferase (TdT)
- Testis
 dissection and germ cell preparation.....170–171
 lysates, preparation of.....198, 199–200

Tetrad analysis	100	Wicking process.....	95 n15, 96 n16
Tetratypes (TTs).....	38	Winkler, Hans (botanist).....	100
Three-factor data	42–45	Y	
random spore data.....	42–43	Yeast (budding), identification of.....	144
tetrad data for chromatid interference.....	43–44	Yeast cell lysate (denatured), preparation of.....	187
tetrad-specific “coefficient of coincidence” for		cell lysis in TCA	189
chromosome interference.....	44	preparing cells for synchronous yeast meiosis... 188–189	
tetrad test for chromosome interference.....	44–45	protein extraction from denatured yeast	
crossover effect in <i>URA-KAN</i> , 45 <i>t</i>		lysate.....	189–190
PD, TT, or NPD scores in <i>URA3-KAN</i>		Yeast culture	
interval	44 <i>t</i>	equipments used.....	24
Topo VI	183, 184	media for.....	121
Tris-buffered saline (TBS)	147, 286	Yeast extract agar (YEA)	236
Two-factor data	38–42	Yeast extract liquid (YEL).....	236
detecting and quantifying interference via		Yeasts, meiotic DSB sites	
internet.....	42	chromatin structure, <i>see</i> Chromatin structure at meiotic	
improvised two-factor test.....	39–41	DSB sites in yeasts	
NPD ratio	39	Yeast strains	31–32
quantifying interference.....	41	YPD (yeast extract, peptone, dextrose) medium	286
<i>fNexp vs. fTexp</i> plot.....	41 <i>f</i>	YPG (yeast extract, peptone, glycerol) medium.....	286
tetrad types and their genotypes	38 <i>t</i>	Z	
W		ZF-SPO11 fusion	31
Watson–Crick base pairing	212	“ZMM” epistasis group.....	8, 48
WCE, <i>see</i> Whole cell extract (WCE)		Zymolyase-treatment.....	265 n3
Whole cell extract (WCE).....	269		
Whole-genome tiling array	302 n5		