

Index

A

Abdominoperineal resection (APR), 111
cocygectomy, 121
coccyx spare, 118
colostomy formation, 119
controlled dissection, 119
da Vinci® surgical system, 112
ELAPR, 111
exploratory laparoscopy, 113
extraction site, 113
extralevator approach, 121
genitourinary structures, 120
inferior mesenteric pedicle, 113–114
intraabdominal levator transection, 115–116
laparoscopic rectal resection, 112
low anterior resection, 111
omental flap infarction, 120
operative room setup, 112–113
patient positioning, 112
perineal closure, 120
perineal dissection, 119–120
perineal hernia, 120–121
peripheral neuropathy, 120
port placement, 113
rectal mobilization, 114–115
RILT, 116
robotic arms collision, 121
tissue prevention, 118
traditional perineal dissection, 115
Anastomotic leaks (AL), 18

C

Circumferential resection margins (CRM), 111, 112, 121, 133
Combined endoscopic and laparoscopic surgery (CELS), 1–3, 5–7

D

da Vinci Xi robotic platform, *see* Robotic total colectomy

E

Endoscopic mucosal resection (EMR), 9
Endoscopic submucosal dissection (ESD), 9
bleeding, 13
costs, 14
elderly patient, 13
en-bloc resection, 13
hydroxyethyl starch, 12
hypromellose, 12
indications, 10
learning curve, 14

lesion identification, 11

lesion injection, 12
operative room setup, 11
over-the-scope clips, 13
patient positioning, 11
perforation, 13
preoperative planning, 11
principles, 10
quality of life, 14
retraction methods, 12
submucosal dissection, 12

Extracorporeal *anastomoses*, 45

Extralevator abdominoperineal resection (ELAPR), 111

F

Fluorescence angiography (FA), 18
Fluorescence, colorectal surgery
equipments, 18
high spatial resolution, 21
indocyanine green, 17
methylene blue, 17, 20
optical and haptic cues, 17
perfusion assessment, 18–20
anastomotic leaks, 18
ICG, 19
PILLAR II trial, 19
PILLAR III trial, 19
TatME, 19
utilisation of fluorescence angiography, 18
peritoneal carcinomatosis, 20
PSW nodes, 20
sentinel node, 20
targeted fluorophores, 21
urinary tract identification, 20

H

Hartmann reversal, 85
anastomosis, 89
colostomy takedown, 88–89
inadequate colon length, 90
intraabdominal colostomy mobilization, 87
morbid obesity, 90
operative room setup, 86
patient positioning, 86
perioperative antibiotics, 86
port placement, 86
postoperative care, 89
preoperative preparation, 85
rectal stump mobilization, 88
splenic flexure mobilization, 87–88

HookKnife™, 12
 HybridKnife®, 12
 Hydroxyethyl starch (HES), 12
 Hyperthermic intraperitoneal chemotherapy (HIPEC), 20
 Hypromellose, 12

I

Ileal pouch-anal anastomosis (IPAA), 19, 105
 anastomosis construction, 109
 anterior dissection, 108
 diverting loop ileostomy, 109
 familial adenomatous polyposis, 105
 ileal mesentery, 107
 patient positioning, 105
 port placement, 105
 posterior dissection, 108
 pouch construction, 106
 superior mesenteric artery, 107
 superior rectal artery, 107–108
 testing anastomosis, 109
 three stage approach, 105
 ulcerative colitis, 105

Indocyanine green (ICG), 17

Inferior mesenteric artery (IMA), 76, 102

Intracorporeal anastomosis (ICA), left colectomy, 55

Alexis retractor, 59
 anvil securing, 64
 anvil to proximal bowel, 61
 bowel preparation, 63
 continuous pressure pneumoperitoneum, 63
 distal bowel lumen, 61
 distal level of resection, 58
 distal margin, 59
 end to end anastomosis, 62–63
 extraction process, 64
 lateral peritoneal attachments, 57
 mesentery division, 57
 operative room setup, 55
 patient positioning, 55
 port placement, 55–56
 proximal level of resection, 57
 proximal margin, 58
 rectal cuff, 64
 specimen extraction, 64
 transrectal delivery, 60–61
 transrectal extraction, 59–60
 transrectal insertion, 59

Intracorporeal anastomosis, right hemicolectomy, 45
 diagonal port arrangement, 47
 duodenum, 49
 exploratory laparoscopy, 49
 ileocolic anastomosis, 51–52
 ileocolic vessels
 identification, 49
 ligation, 49
 ileum and colonic mesentery, 50–51
 ileum and transverse colon, 51
 lateral attachments, 49
 operative room set up, 45–46
 patient positioning, 45
 preoperative planning, 45
 specimen extraction, 52–53
 suprapubic port arrangement, 47–48
 terminal ileum and hepatic flexure, 50

L

Laparoscopic-assisted polypectomy, 1
 air leak test, 6
 bleeding, 7
 CELS approach, 1
 colonoscopic-assisted laparoscopic partial cecectomy, 6
 colonoscopy, 3
 contraindications, 7
 electrosurgical snare polypectomy, 3
 foley catheter, 2
 full thickness CELS, 5–6
 mobilization, 3
 morbid obesity, 7
 nasogastric drainage tube, 2
 operative room setup, 2
 patient positioning, 2
 perforation, 7
 polypectomy, 3–5
 port placement, 3
 post-operative care, 6–7
 preoperative planning, 2
 submucosal dilute dye injection, 5
 thermal colon injury, 4

Laparoscopy, 25

Low anterior resection (LAR), 111

P

Pelvic sidewall (PSW) nodes, 20

R

Rectal mobilization
 anterior dissection, 115
 anterior rectal mobilization, 103
 lateral dissection, 115
 lateral rectal mobilization, 103–104
 posterior dissection, 114–115
 posterior rectal mobilization, 103
 rectal division, 104

Rectus abdominis muscle flap, pelvic reconstruction, 123
 complications, 127
 deep inferior epigastric pedicle, 126
 gracilis muscle flap, 124
 inset of rectus muscle, 127
 mesh placement, 126–127
 myocutaneous and muscle flaps, 123
 patient poistioning, 125
 patient selection, 124
 port placement, 125
 posterior rectus sheath, 126
 postoperative care, 127
 rectus abdominis muscle, 124
 robotic approach, 124

Rectus muscle division, 126

Robotic colorectal surgery, 31
 clinical scenarios, optimization, 37–40
 components, 32
 credentialing, 40
 crowdsourcing, 42
 da Vinci platform, 33
 laparoscopic techniques, 33
 learning equipment and system components, 34–37

- clinical sales representative, 35
online modules, 34–35
simulator, 35
recredentialing, 42
simulators, 40
social media outlets, 42
3D systems, 40
trainee proficiency, 40
Robotic intra-abdominal levator transection (RIILT), 116
Robotic surgery
 current FDA approved platforms, 26
 da Vinci® SP, 27
 da Vinci® X, 27
 da Vinci® Xi, 26–27
 description, 26
 DiLumen C™ by Lumendi, Ltd., 28
 Flex® by Medrobotics® Corporation, 27–28
 Senhance™ by TransEnterix Surgical, Inc, 27
Robotic total colectomy, 99
 ascending colon lateral mobilization, 101
 descending and sigmoid colon mobilization, 103
 identification and ligation of IMA and ureter, 102
 ileocolic pedicle isolation, 101
 lateral ascending colon mobilization, 101
 left sided and rectal mobilization, 102
 operative room setup, 99
 patient positioning, 99
 port placement, 100
 preoperative planning, 99
 rectal mobilization, 103–104
 right sided mobilization, 101
 splenic flexure mobilization, 102–103
 terminal ileum transection, 101–102
 transverse colon lateral mobilization, 101
 transverse mesocolic dissection, 101
Robotic ventral mesh rectopexy (RVMR), *see* Ventral mesh rectopexy
- S**
Superior mesenteric vein (SMV), 69
- T**
Total mesocolic excision (TMCE) technique, 67
 da Vinci Si platform, 68
 da Vinci Xi platform, 68
 distal transverse colon cancer, 79–82
 colon transection, 81
 IMA dissection and division, 79
 IMV dissection and division, 79
 intracorporeal anastomosis, 81–82
 middle colic vessels, 80
 specimen extraction, 81–82
 splenic flexure, 80
 hypothesis, 67
 operating room setup, 68
 patient positioning, 68
 robotic anterior and left colectomy
 IMA dissection and division, 77
 IMV identification and dissection, 77
 port placement, 75
 proximal colon, 79
 rectosigmoid colon, 78
 sigmoid and descending colon, 77–78
 splenic flexure, 78
superior rectal artery, 78
suprapubic Pfannenstiel incision, 79
robotic right and extended right
 colic vessels, 71–72
 cranio-caudal approach, 74–75
 distal ileum, 72–73
 exploratory laparoscopy, 69
 Henle's trunk, 71
 hepatic flexure, 73
 ileocolic vessels dissection and division, 70–71
 intracorporeal anastomosis, 73–74
 port placement, 68
 SMV to ileocolic pedicle junction, 69–70
 specimen extraction, 73–74
Total mesorectal excision (TME), 143
 abdominal dissection, 146
 abdominal field, 145
 anastomotic techniques, 150
 connection of abdominal and perineal plane, 149
 dissection technique, 143
 equipment, 144
 GelPOINT Path access sleeve, 146
 laparoscopic transanal dissection, 148–149
 operative room setup, 144–145
 patient positioning, 145
 perineal dissection, 146
 port placement, 145
 preoperative planning, 144
 pure laparoscopic approach, 146
 single surgeon approach, 144
 abdominal dissection, 146
 perineal dissection, 151–152
 specimen extraction, 150
Transanal endoscopic microsurgery (TEM), 131, 135
Transanal minimally invasive surgery (TAMIS), 131–132, 135
 closure of defect, 140–141
 exploratory anoprostoscopy, 139
 full thickness dissection, 139
 older vs newer robotic platform, 141–142
 operative room setup, 136–138
 operative tools, 138
 patient positioning, 136
 postoperative followup, 141
 preoperative planning, 136
 robotic docking, 138
 technical Feasibility, 141
 trocar placement, 138
 variations in patient positioning, 141
Transanal total mesorectal excision, inflammatory bowel disease, 19, 132–133, 153
 abdominal colectomy, 156
 anastomosis, 157
 bilateral laparoscopic TAP, 157
 connection of abdominal and perineal dissection, 157
 equipments, 154
 GelPOINT® path, 156
 laparoscopic, 156–157
 Lone Star® retractor, 156
 operative room setup, 153
 patient poistioning, 153
 port placement, 155
 proctectomy, 156
 specimen extraction, 157
 terminal ileal mesentery, 156
Transversus abdominis plane (TAP), 157

V

Ventral mesh rectopexy, 91
anesthesia preparation, 92
anterior longitudinal ligament, 93
clearing pelvis, 92–93
complications, 96
goals, 91
mesh fixation, 94–95
operative room setup, 92
outcomes, 95–96

patient positioning, 92
patient selection, 91
peritoneum closure, 95
peritoneum flaps, 93
polypropylene mesh/biological graft, 94
port placement, 92
pouch of Douglas, 94
preoperative preparation, 91
rectovaginal septum, 94
vaginal and rectal examination, 92