

Index

A

- Additive method, 120
- Ageing, 54–55
- Aniridia, 151–153, 244
- Anophthalmia. *See* Congenital anophthalmia
- Anophthalmic patient
 - children, psychological issues for, 12–13
 - eye loss
 - aetiology of, 3–4
 - anophthalmic population, estimated size of, 2
 - biosocial and psychological aspects, 1
 - body image loss, 8–9
 - causes, 4–6
 - novelty eyes, 9–10
 - perceptual changes accompanying, 6–7
 - personal accounts, 13–21
 - phantom eye pain, 11
 - prosthetic eye wearers, 11–12
- Anophthalmic socket
 - conjunctiva of, 57–59
 - forniceal region, 43–44
 - GPC, 43
 - inflammation of, 110–112
 - tears
 - distribution, 53–54
 - film, 51–52
 - function of, 50–51
 - glands, 49
 - Jones test, 104, 106
 - nasolacrimal drainage system, 108
 - ocular tear ferning test, 108, 109
 - output, 52–53
 - phenol red thread test, 104
 - Schirmer tests, 104
 - TBUT test, 109, 110
 - volume, 106–108
 - types, 99, 102–104
- Arcus senilis, 55, 128, 130

B

- Bartisch, Georg, 84, 86
- Bioeye, 82
- Blood vessels, 38
- Body image, 8–9, 12, 13
- Boissonneau, Auguste, 291, 294
- Bulbar region, 43

C

- CMYK subtractive colour, 120, 121
- Collarette, 129
- Colour theory
 - characteristics, 121–122
 - rods and cones, 119–120
 - wheel, 120–121
- Cones, 119, 120
- Congenital anophthalmia
 - cases, 68
 - CT image, 68, 69
 - eye association, 68
 - treatment, 70–71
- Congenital microphthalmia
 - CT image, 68, 69
 - eye association, 68
 - treatment, 70–71
- Conjunctiva
 - of anophthalmic socket, 57–59
 - bulbar region, 43
 - forniceal region, 43–44
 - palpebral region, 41–43
 - sensitivity of, 44–46
 - single continuous mucous membrane with regions, 41, 42
 - structure of
 - epithelial layer, 46–47
 - goblet cells, 47, 48
 - mucus, function of, 48
 - substantia propria layer, 46

- Conjunctiva (*cont.*)
 surface papillary texture, assessment of, 113–115
- Contact lens-induced papillary conjunctivitis (CLPC), 252
- Cornea
 arcus senilis, 55
 corneal unit (*see* Iris/corneal unit)
 premanufactured iris discs and, 124–125
 sclera, 33
 sensitivity of, 44–46
 tattooing, 73
- Corneal opacities
 aetiologies and clinical characteristics of, 5
 densities and dimensions, 153, 154
- Corneal tattooing, 72, 73
- Corrugator muscles, 39
- Coulomb, Robert, 301, 303
- Cranium, bones, 30
- Custom-fit prosthetic eyes, 118–119
- D**
- Deposit build-up
 inter-palpebral zone, 222, 224
 rate
 deposit formation, intensity and extent of, 221, 222
 equal interval perceptual grading scales, 221, 223
 photographic record, 221, 224
 regression analysis, 222, 225
 retro-palpebral zone, 222, 224
 surface polish effect, 222–224, 226
 wettability, 227–228
- Disfigurement, 67, 71–73
- Driver and Vehicle Licensing Authority (DVLA), 268–269
- Dry eye syndrome, 93, 108, 244
- E**
- Ectropion, 207–208
- Elli, 13, 14
- Entropion
 cicatricial entropion, 208
 involutional entropion, 208
 prosthetic technique, 209
 surgical technique, 209
 ‘windscreen wiper’ effect, 208
- Enucleation, 73–76
 of eye, 74–76
 vs. evisceration, 73–74
 PESS, 61–64
 postsurgical care following, 77–80
- Evisceration
 enucleation vs., 73–74
 postsurgical care following, 77–80
 surgical removal of, 73, 76–77
- Exenteration. *See* Orbital exenteration
- Existing prosthetic eye in situ assessment, 95–96
- Extraocular muscles, 6, 32, 34–36, 57, 74–76, 81, 83, 191, 197
- Eyelid malposition
 ectropion, 207–208
 entropion
 cicatricial entropion, 208
 involutional entropion, 208
 prosthetic technique, 209
 surgical technique, 209
 ‘windscreen wiper’ effect, 208
- lagophthalmos
 anterior lamellar deficiency, 214–215
 middle lamella deficiency, 215
 posterior lamellar deficiency, 215
 prosthetic technique, 214, 215
- ptosis
 levator function, 210
 prosthetic technique, 210–212
- Eyelid-related pathology
 blepharitis, 243–244
 dry eye syndrome, 244
 lagophthalmos, 246
 MGD, 244–246
- Eyelids
 anatomical features, 28, 40–41
 muscles of
 corrugator, 39
 facial musculature of, 38, 39
 frontalis, 39
 orbicularis oculi, 39–40
 procerus, 39
 pupil, 28, 29
 upper eyelid angles, margin of, 29
 upper eyelid crease, 29, 30
- Eye loss
 biosocial and psychological aspects, 1
 children, psychological issues for, 12–13
 epidemiology of
 aetiology of, 3–4
 anophthalmic population, estimated size of, 2–3
 causes, 4–6
 perceptual changes, 6–7
 personal accounts of, 13–21
 psychosocial issues
 body image loss, 8–9
 novelty eyes, 9–10
 prosthetic eye comfort and convenience, 10–11
- Eye of Horus, 284, 285

F

Face

- ageing of, 54–55
- architecture, 27–28
- skeleton of, 30
- sound appreciation of the
 - characteristics of, 26

Facial symmetry, 4, 68

FD OCT. *See* Fourier domain optical coherence tomography (FD OCT)

Fick, Adolf, 310, 311

Forehead, muscles of

- corrugator, 39
- facial musculature of, 38, 39
- frontalis, 39
- orbicularis oculi, 39–40
- procerus, 39

Forniceal region, 43–44

Fourier domain optical coherence tomography (FD OCT), 107–108

Frontalis muscle, 39, 210, 212

G

Giant papillary conjunctivitis (GPC), 43, 113–115, 180, 234–235, 252

Goblet cells, 47–49, 51, 52

GPC. *See* Giant papillary conjunctivitis (GPC)

H

History

- ancient Egypt, 283–285
- contact lenses and scleral shell prostheses, 310–311
- eighteenth century, 290
- nineteenth century
 - artificial eyes, 291, 294
 - callipers, 295, 296
 - coloured glass rods, 295, 298
 - creating pupil, 295, 299
 - cryolite glass eye, 293, 295
 - European ocular prosthetics, 291, 293
 - glass blowing instruments and materials, 295, 296
 - glass eye industry, 291–294, 299, 302, 303
 - glass eye prescription form, 295, 297
 - glass sphere implant, 296–297, 301
 - handling glass eye instruments, 295, 297
 - importing stocks of glass eyes, 299, 302
 - partially eye with glass rods, 295, 299
 - partially finished glass, 295, 300
 - posterior aspect of the eye, 295, 300

preformed glass eyes, 295, 300

‘reform’ eye, 295

seal tube, 295, 298

softened glass, 295, 300

softening pre-tinted hollow glass tube, 295, 298

Traité pratique de l’oeil artificiel, 290, 291

William Mackenzie’s *A practical treatise on the diseases of the eye*, 291, 292

seventeenth century, 289

sixteenth century, 287–288

3000 BC, 284, 285

twentieth century

Coulomb, Robert *L’oeil artificiel*, 301, 303

custom-made PMMA eye, 304, 306

German glass eyes, 304, 307

L’oeil artificiel, 301

maxillofacial prosthetics, 304, 308

moveable eye article, 304, 305

plastic artificial eyes, 304, 306

stock glass eyes, 304, 308

twenty-first century, 309–310

2000 BC, 285–287

Hydrogel, 70–71, 227, 252

Hydroxyapatite implants, 81, 82

I

Impact-resistant glasses, 12, 13

Imperial Chemical Industries (ICI), 301

Inferior oblique muscle, 35–36, 74

InflammaDry™ technology, 110, 111

Intraorbital implants

Bioeye, 82

coral, 81

material, 82–83

orbital implant, 82

shape, 83

size, 83

titanium, 82

wrapping, 83–84

Iris/corneal unit

alternative methods, 125–127

painting

collarette, 129

individual component, 128, 129

limbus, 130

magnifying lamp, 128

pupil, 128–129

stroma, 129–130

premanufactured iris discs and corneas, 124–125

prosthetic eye, components, 122

- Iris/corneal unit (*cont.*)
 trying, wax pattern and positioning
 central gaze, rod aligned in, 139
 compromises and trade-offs, 139, 140
 fitting bilateral prosthetic eyes,
 139–140
 process, 138
 vernier gauge, 123
- J**
 Jones test, 104, 108
- K**
 Keratoconjunctivitis sicca. *See* Dry eye syndrome
 Krause's glands, 44, 46, 49, 55
- L**
 Lacrimal apparatus, 48–50
 Lacrimal gland, 37–38, 48, 49, 55, 93
 Lacrimal sac, 38
 Lactoferrin, 50
 Lagophthalmos
 anterior lamellar deficiency, 214
 middle lamella deficiency, 215
 posterior lamellar deficiency, 215
 prosthetic technique, 214, 215
 Levator muscle, 37
 Limbus, 44, 74, 130, 210
 Lysozyme, 50, 252
- M**
 Marginal zone, 42, 43
 Medial canthal tendon (MCT) laxity, 38, 186
 Meibomian gland dysfunction (MGD),
 243–246, 273
 Meibomian gland loss assessment, 112
 Methylmethacrylate and N-vinylpyrrolidone,
 70–71
 Mucoïd discharge
 CLPC/GPC, 252
 managing non-specific mucoïd discharge,
 254–257
 non-specific causes, 251–252
 polish improvement, 253
 range of socket characteristics, 242
 re-polishing artificial eyes, 251, 253
 specific causes
 cleaning prosthesis with solvent,
 249, 250
 environmental allergens, 246–247
 environmental irritants, 247
 extruding implants, 247, 248
 eyelid-related pathology, 243–246
 eye stress, 247
 gritty deposits, 249, 250
 mixed conjunctival membrane,
 247, 249
 pegged hydroxyapatite implant,
 247, 248
 scratches and dull patches, 249, 250
 silvery discoloration, 249
 thinly covered implants, 247
 undissolved sutures with/without
 granulomas, 247, 248
 viral/bacterial infections, 242–243
 tear protein deposition, 252
 wetting angle vs. surface finish, 254
 Mucoïd discharge assessment, 112–113
 Mucus, 48
 Mules, Phillip Henry, 296–297, 301
 Müller, Albert, 310
 Müller, Friedrich, 310
 Müller-Ur, Ludwig, 293, 295
- N**
 Nasolacrimal drainage system, 108
- O**
 Ocular globe, loss of
 anophthalmic socket, conjunctiva of,
 57–59
 empty socket, 55, 56
 PESS, 61–64
 prosthetic eye motility, 57, 58
 restore, 55, 56
 superficial anatomical features, 55, 56
 tear distribution, 60–61
 tear film, 59, 60
 tear output, 60
 Ocular tear ferning test, 108, 109
 Ocular tray impression technique, 131–133
 Optic nerve, 34
 Orbicularis oculi muscle, 39–40
 Orbit
 contents of
 blood vessels and nerves, 38
 extraocular muscles, 34–36
 eye, 32–33
 lacrimal sac, 38
 levator muscle, 37
 optic nerve, 34
 orbital fat, 38
 floor of, 31

- lateral wall of, 31
- medial orbital wall, 31
- pyramidal shape of, 31
- rim, 31
- roof, 31
- Orbital exenteration
 - additional diseased bone, removal of, 84, 86
 - adhesive-retained prosthesis, 87
 - Bartsch, Georg, 84, 86
 - content removal, 84, 85
 - eyelid skin and orbicularis muscle tissue, retaining, 84, 85
 - full-thickness skin graft, 85, 86
 - indication, 84
 - inserting magnets, 87
 - in mould, 88
 - prosthesis, non-existent sockets, 85, 86
- Orbital fat, 38
- Orbital volume deficit
 - expanded orbital volume, 204–205
 - orbital volume replacement
 - prosthetic technique, 200–201
 - secondary orbital implant, 201–203
 - subperiosteal implant, 203–204
 - PESS, 200
- Orbital zone, 43

- P**
- Palpebral fissures, 27, 140
- Palpebral region, 41–43
- Patient evaluation
 - anophthalmic sockets assessment
 - nasolacrimal drainage system, 108
 - ocular tear ferning test, 108, 109
 - TBUT test, 109, 110
 - tear volume, 106–108
 - tests, 104, 106
 - types, 99–104, 110–112
 - different patients, 91, 92
 - existing prosthetic eye in situ assessment, 95, 96
 - medical history, 93
 - meibomian gland loss assessment, 112
 - mucoid discharge assessment, 112–113
 - personal history, 91, 92
 - prosthetic eye ex situ assessment, 96–99
 - psychological assessment, 93–94
 - remaining sighted eye, health of, 94
 - surface papillary texture, assessment of, 113–115
 - visual perception assessment, 94
- Personal socket hygiene, 272–273
- Phantom eye pain, 11
- Phenol red thread test, 53, 104, 107
- Phenol red thread testing kit, 107
- Plaster mold casting, 137, 138
- (Poly)methyl methacrylate (PMMA)
 - conformer shells, 80
 - delamination, 98, 101
 - intraorbital implants, 81–82
 - spherical implant, 104
 - tear film, 59–60
 - wax pattern
 - alternative method, 137, 138
 - applying, second layer of iris colours, 141, 142
 - cooled down and removed, 135, 136
 - fine veins teasing, 141, 143
 - iris painting and scleral colouring, set up for, 141, 142
 - mould preparation, 134, 135
 - PMMA, clear veneer of, 143
 - preheated white ocular wax, 134–136
 - process completion, 135, 137
 - rod attaches, 141
 - sectioning, 134, 135
 - shape, 135, 137
 - trial packed with white PMMA, 141, 142
- Polyvinylsiloxane, 133
- Post-enucleation socket syndrome (PESS), 61–64, 200
- Postsurgical conformer shells, 80–81
- Procerus muscle, 39
- Prosthetic eye making and fitting
 - final clinical session, 146
 - impression
 - impression mixing gun technique, 131, 133–134
 - ocular tray impression technique, 131–133
 - polyvinylsiloxane, 130, 131
 - polishing, process for, 144–146
 - prosthetic eye making and fitting
 - clinical session, 122–123
 - components, 122
 - iris/corneal unit (*see* Iris/corneal unit) step for, 122
 - stock vs. custom-fit, 118–119
 - in twentieth century, 301–304
- Prosthesis retention
 - expanded conformer and tarsorrhaphy
 - custom-made conformer, 186, 190
 - eyelid-sparing exenteration, 186, 189
 - self-retentive prosthetic eye, 187
 - inadequate fornix/fornices
 - adhesions, 184
 - surgical technique, 185

- Prosthesis retention (*cont.*)
- inadequate socket lining
 - custom-made pressure conformer, 181
 - dermis-fat graft, 184
 - mucous membrane graft, 182–183
 - non-retentive contracted socket, 180–181
 - skin/split-thickness skin grafts, 184
 - lower eyelid laxity
 - anophthalmic socket, 185
 - inadequate lid tension, 185
 - prosthetic technique, 186
 - surgical technique, 186, 188
- Prosthetic contact lenses
- albinism, 155, 156
 - aniridia, 151–153
 - cataract appearance, 155
 - corneal dystrophy, 157
 - corneal leucoma, 151, 153
 - corneal opacities/scars, 153, 154
 - designs, 151, 152
 - Gundersen flap procedure, 150
 - heterochromia, 155–156
 - in-house colouring system, 157–158
 - iris coloboma, 154
 - soft hand-painted lenses, 159
 - soft opaque lens, 158, 159
- Prosthetic eyes
- altered depth perception
 - ball approaches, 265
 - parallax, 266, 267
 - perspective, contrast and overlap objects, 267, 268
 - relative motion, 263–266
 - senses, 267, 268
 - application, 237–238
 - cleaning prosthetic eyes, 276–277
 - ex situ assessment, 96–99
 - glasses selection, 278–279
 - handling social interactions, 260–262
 - managing mucoid discharge, 272
 - monocular vision
 - driver safety technology, 272
 - driving, 270–271
 - legal requirements, 268–271
 - parking, 271
 - motility, 57, 58
 - psychological importance of, 10–11
 - reduced peripheral vision
 - discrete use of mirrors, 263, 264
 - preplanning, 262, 263
 - strategically position, 263, 264
 - removal, cleaning and reinsertion
 - prosthesis
 - bacteria and foreign material, 230
 - caregiver guide, 273–274
 - disturbed socket environment, 230–231
 - patient guide, 275–276
 - physical stress, 229–230
 - processes, 229
 - surface characteristics, 231
 - resources, 280–281
 - supero-nasal notches in, 35
 - tear distribution, 60–61
 - tear film, 59, 60
 - tear output, 60
 - tear protein deposits
 - analogous, 220, 221
 - contact lens research, 220–221
 - deposit build-up (*see* Deposit build-up)
 - temporal limbus area, 220
 - three-phase model (*see* Three-phase model)
 - wearers, 2, 3, 8, 9
 - concerns of, 11–13
 - personal accounts, 13–21
 - population of, 2–3
- Ptosis
- levator function, 210
 - prosthetic technique, 210–212
- R**
- Rectus muscles, 35, 57, 74
 - Retina, 33, 120
 - Rigid gas-permeable (RGP) contact lenses, 227
 - Ritterich, Friedrich Philipp, 291–294
 - Rods, 119, 120
 - Rosacea, 243
 - Rose Bengal, 96, 99
- S**
- Schirmer test, 53, 106
 - Scleral shell prostheses
 - completed medium thickness, 161
 - custom metal moulds, 162
 - distorted cornea, 160
 - final scleral shell construction
 - edges polishing, 171, 172
 - fitting scleral shell prosthesis, 171, 172
 - iris and scleral colours, 170
 - PMMA veneer process, 171
 - polyurethane sheet, 170–171
 - pupil, collarette, stroma and limbus colours, 170

- remove white, clear/semitranslucent shell, 169
 - rubber suction devices, 171, 172
 - Gundersen flap procedure, 150
 - history, 310–311
 - left strabismic eye, 160, 161
 - making and fitting thin scleral shell prostheses, 163
 - manufacturing process, 163–165
 - opaque cornea, 160
 - polyvinylsiloxane impression, 166, 167
 - premanufactured two-dimensional curved iris discs, 162
 - thin scleral shell design, 163, 164
 - topical anaesthetic eye drops, 166
 - trial/test shell
 - construction of, 167
 - fenestration, 168–169
 - plaster separating solution, 167
 - two-part mould, 167, 168
 - UV and white light sources, 167
 - Seborrhoeic dermatitis, 243
 - Skull, 30
 - Socket complications, 177, 215–216
 - eyelid malposition (*see* Eyelid malposition)
 - lesions
 - chemosis, 179
 - GPC, 180
 - granulomas, 178–179
 - malignancy, 178
 - orbital volume deficit (*see* orbital volume deficit)
 - prosthesis motility
 - axes of movement, 197
 - excessive prosthetic eye motility, 198–199
 - inadequate prosthetic eye motility, 197–198
 - prosthesis retention (*see* Prosthesis retention)
 - socket stability (*see* Socket stability)
 - superior sulcus deformity
 - dermis–fat graft and fillers, 207
 - prosthetic technique, 205–206
 - surgical techniques, 205
 - upper eyelid blepharoplasty, 205, 207
 - Socket stability
 - displacing prosthetic eye, 190, 192
 - inadequate prosthesis retention, 190, 192
 - orbital implant exposure
 - prosthetic technique, 193
 - surgical technique, 194–196
 - symptoms, 192, 194
 - orbital implant migration, 191–194
 - rotational instability, 196
 - Stereopsis, 6
 - Stock prosthetic eyes, 118–119
 - Stroma, 129–130
 - Substantia propria layer, 46
 - Subtractive colour system, 120, 121
 - Superior oblique muscle, 35
 - Surface papillary texture assessment, 113–115
 - Syringing technique, 273
- T**
- Tarsal zone, 43
 - Tear Break-Up Time (TBUT) test, 109, 110
 - Tears
 - assessment, anophthalmic socket
 - nasolacrimal drainage system, 108
 - ocular tear ferning test, 108, 109
 - TBUT test, 109, 110
 - tear volume, 106–108
 - tests, 104, 106
 - distribution, 53–54
 - ferning test, 108, 109
 - film, 51–52, 59, 60
 - function of, 50–51
 - glands, 49
 - output, 52–53, 60
 - Tearscope, 109, 110
 - Three-phase model
 - breakdown phase, 232
 - epitheliopathy of Marx's line, 234, 235
 - GPC, 234–236
 - over time deposits thicken and encroach, 234
 - physiological homeostasis, 236, 237
 - scanning electron microscopy, 234
 - stagnation, 236
 - equilibrium phase, 231, 233
 - establishment phase, 231, 232–233
- V**
- Vascular choroid, 33
 - Visual perception assessment, 94
- W**
- Wax pattern, 134–139