

Index

A

- Adaptive evolution
 - climate change, 134
 - heritable traits, 134
 - management implications, 147
 - neutral genetic variation, 139–141
 - new genotypes, 134
 - research, 148–149
- Adaptive management, 438, 444, 453
- Adaptive management framework
 - biophysical system, 345
 - evaluation, 348–350
 - feedback loop, 345
 - management goals and restoration strategies, 345
 - planning, 346–348
- Agate Fossil Beds National Monument, 38–40
- AIM. *See* Assessment, inventory and monitoring (AIM) strategy
- Albedo, 74, 75
- Allelic richness, 122, 125
- Allozymes (enzyme electrophoresis), 104
- Alternative stable states, 439
- Annual grass and tree invasion disturbances, 402
- Arbuscular mycorrhizal fungi (AMF), 82
- Asian Carp Prevention and Control Act of 2010, 418
- Assessment, inventory and monitoring (AIM) strategy, 360
- Attitudes
 - change strategies, 411, 415–416
 - expressions of people's values and beliefs, 411
 - toward invasive plants, 413–415
 - toward management options, 413–415

B

- Badlands National Park, 38
- Behavioral intentions, 416–417
- Beliefs about exotic annual grasses, 412–413
- Beliefs, human dimensions
 - beliefs about exotic invasive plants, 412–413
 - and management options, 412–413
- Benefit-cost ratios, 431, 433, 434, 444, 445, 449
- Bio-economic modeling
 - analyze management issues, 434
 - biological invasions, 434
 - catastrophic wildfire, 432
 - decision-maker, 431
 - decision-maker's objective function, 432
 - design, 434
 - and dynamic, 433
 - individual (or firm) whose primary concern, 431
 - management, 434
 - market rate of interest, 433
 - mathematical relationships, 432
 - objective function, 432
 - private decision-makers, 431
 - private decision-making models, 431, 432
 - public decision-makers, 431
 - public-good nature, 431
 - risk-averse decision-maker, 433
 - risk-averse ranchers, 433
 - risk-neutral decision-maker, 433
 - social rate of time preference, 433
 - trade-offs, 434
 - uncertainty and risk, 432

- Bio-economic models
 development, 430
 simulation and optimization, 430
- Biological crusts, 291–292
- Biological invasions
 and communities, 107
 post-immigration evolution, 108
 preadaptation, 108
 propagule pressure, 107, 108
- Blackbrush shrubland, 18
- Bleach blonde syndrome, 214
- BLM. *See* Bureau of Land Management (BLM)
- Boissiera*, 157, 158
- Bromeae, 157
- Bromopsis*, 157, 163, 165, 171, 173,
 176–178, 184
- Bromus alopecurus*, 184
- Bromus andringitensis*, 163
- Bromus arenarius*, 184
- Bromus arvensis*, 16, 180, 184, 185, 342–343
- Bromus auleticus*, 171, 178
- Bromus benekenii*, 171
- Bromus berterioanus*, 184
- Bromus biebersteinii*, 178
- Bromus briziformis*, 184
- Bromus carinatus*, 180
- Bromus catharticus*, 171, 178, 184
- Bromus cebadilla*, 184
- Bromus commutatus*, 184
- Bromus danthoniae*, 184
- Bromus diandrus*, 172, 176, 184, 342
- Bromus diandrus* Roth (ripgut brome), 16
- Bromus erectus*, 171, 178, 184
- Bromus fasciculatus*, 183
- Bromus grossus*, 171
- Bromus hordeaceus*, 171, 172, 180, 184, 185,
 341–342
- Bromus hordeaceus* L. (soft brome), 16
- Bromus inermis*, 171, 178, 183–185, 419, 420,
 423, 425
- Bromus interruptus*, 171
- Bromus japonicus*, 16, 171, 184, 185
- Bromus lanceolatus*, 184
- Bromus lepidus*, 172, 184
- Bromus madritensis*, 17, 184
- Bromus madritensis* L. (compact brome), 16
- Bromus marginatus*, 180
- Bromus mollis* (syn. of *Bromus hordeaceus*), 180
- Bromus pectinatus*, 184
- Bromus pumpellianus*, 171, 178
- Bromus racemosus*, 184
- Bromus ramosus*, 171
- Bromus rigidus*, 184
- Bromus riparius*, 171, 178
- Bromus rubens*, 16, 17, 156, 170, 176, 180,
 184, 185, 263, 343, 410, 411, 419,
 420, 425
- Bromus scoparius*, 184
- Bromus secalinus*, 171, 176
- Bromus sericeus*, 183, 184
- Bromus squarrosus*, 184
- Bromus sterilis*, 184
- Bromus tectorum*, 16, 156, 157, 170, 172, 176,
 180, 183–185, 293, 343–344, 390,
 391, 393, 410–412, 414, 415,
 418–420, 423, 425
- abiotic factors, 195
- A. tridentata* spp. *vaseyana* model, 389,
 394, 395
- A. tridentata* spp. *wyomingensis*, 394, 396
- Beetle (mountain big sagebrush), 388
- bleach blonde syndrome, 217
- climate change, 388, 394, 397
- colonize bare soil, 218
- cyclic succession, 195
- die-off/stand failure, 195
- dispersal and lose dormancy, 194
- ecological processes
 CO₂ levels, 391
 fertilization, 393
 forcing factors, 391
 future time series replicates, 391
 GCM, 391
 invasive annual grass and tree
 expansion, 393
 PDSI, 393
 shrubland–woodland, 393
 spatial averaging methods, 391
 STSMs, 393
 temporal multipliers, 390
- ecological systems, 388
- environmental tolerance, 137–138
- environmental values, 388
- Epicoccum*, 219
- epidemic disease, 220
- fall seed bank densities, 194
- florets, 194
- fungal pathogens, 194
- fungistasis, 218
- GCMs and MC1, 397
- genetic analyses, 140
- genetic variation, 135
- germinating seeds/preemergent
 seedlings, 195
- global circulation models, 389
- granivores, 194
- habitats, 135
- ILAP, 397

- investigations, 218
 - landscape-level approach, 220
 - life cycle, 216
 - litter dynamics, 218
 - management implications, 219–220
 - microenvironmental factors, 220
 - monocultures, 194
 - multiple research approaches, 218
 - mycoherbicidal biocontrol, 219
 - NEPA, 388
 - Path Landscape Model (Path) software, 389
 - pathogens, 216, 217
 - productivity, 194
 - range shifts, 390, 397
 - reciprocal transplant studies, 135
 - remote sensing data, 394
 - restoration tool, 195
 - seedling emergence, 194
 - soilborne pathogens, 194
 - soil moisture, 397
 - state-and-transition models, 389
 - STM transitions, 394
 - top-down and bottom-up approach, 397
 - vegetation classes, 394
 - winter annual grass, 134
 - winterkill, 195
 - B. tectorum* in North America, 108
 - distribution, 110
 - distribution of MLGs, 109
 - Eastern US populations, 109
 - genetic diversity, 109
 - herbarium specimens, 109, 113, 120
 - history and the concatenation, 109
 - Lancaster County, Pennsylvania, 109
 - MCG, 109, 110 (*see* Most common genotype (MCG))
 - MLGs, 112 (*see* Multilocus genotypes (MLGs))
 - multiple introductions, 109, 117, 121, 122
 - 60 populations, 112
 - B. tectorum* L.
 - invasion, 100
 - Bromus tectorum* L. (cheatgrass or downy brome), 434, 442
 - B. tectorum* research
 - assess, 356–357
 - BLM approach, 356
 - burning, 353–354
 - “command and control” approach, 356
 - defoliation, 354
 - design, 357–359
 - herbicide application, 354–355
 - implementation, 358, 359
 - literature review, 350–352
 - monitoring, 360
 - revegetation, 355
 - treatment outcomes, 352
 - wildlife and native plant communities, 355
 - Bromus tectorum* ssp. *lucidus*
 - (syn. of *Bromus sericeus*), 183
 - Bromus texensis*, 165
 - Bromus trinitii*, 184
 - budget, 432, 443, 444, 447
 - Bullae/bunts, 202
 - Bureau of Land Management (BLM),
 - 316, 418
- C**
- Calcium (Ca), 81, 82
 - Carbon dioxide (CO₂) concentration, 145
 - C:N ratios, 145
 - environmental conditions, 144
 - phenology, 145
 - precipitation, 146
 - warming-induced changes, 145
 - water availability, 146
 - Carbon sequestration, 76
 - Cation-exchange capacity (CEC), 245
 - Cattle prices, 436, 440, 450
 - CEC. *See* Cation-exchange capacity (CEC)
 - Centrocercus urophasianus* Bonaparte, 3
 - Ceratochloa*, 157, 163–165, 171, 174, 177, 178, 184
 - Cercocarpus ledifolius*, 135
 - Chloroplast spacer regions *trnT-trnL* and *trnL-trnF*, 102
 - Cleistogamous flowers, 104
 - Climate change, 262, 263
 - adaptive management practices, 270
 - anthropogenic greenhouse gas emissions, 258, 259
 - Bromus*, 258, 262
 - B. tectorum* and *B. rubens*, 267, 269
 - CO₂ concentration, 259
 - ecological responses, 143
 - ecosystems, 260–262
 - emissions, 259
 - expansion/contraction, 259
 - experimental and modeling, 271
 - fall/spring precipitation
 - B. tectorum*, 262
 - El Niño events, 263
 - germination and growth, 263
 - global change factors, 263
 - IPCC, 263
 - sagebrush steppe ecosystems, 262
 - seasonal congruence, 263

- Climate change (*cont.*)
- semiarid ecosystems, 262
 - snowmelt/spring rains, 262
 - wet winter/spring conditions, 262
- GAP analysis plots, 267
- global change manipulations, 144
- IPCC, 259
- management implications, 270
- models projections, 267
- native and invasive plants, 258
- phenology, 258
- physiological tolerances, 258
- PRISM, 267
- propagules, 258
- radiative forcing, 259
- regional species distribution modeling, 264
- representative concentration pathway/
RCP4.5, 259
- research needs, 271
- season precipitation, 268–270
- soil moisture, 260
- species persistence, 143
- substantial influence, 143
- summer precipitation, 263, 264
- temperature and precipitation, 258
- warmer temperatures, 258
- water stress, 259
- Winter and spring temperatures, 266–267
- winter precipitation, 266
- Climatic regimes
- native ecosystems, 2
- CMAQ. *See* Community Multiscale Air
Quality (CMAQ)
- C:N, 79
- Codominant genetic markers, 104
- Coefficient of inbreeding, 105
- Cold deserts, 32–34
- ecosystem threats, 34–35
 - environmental conditions, 31
 - invasion potential
 - ENSO events, 32
 - low to mid elevations, 33
 - moderate climates, 33
 - resilience to disturbance, 32, 33
 - resistance, 32, 33
 - salt desert vegetation types, 32
 - seasonality of precipitation, 33
 - soil temperature regimes, 33
 - upper elevations, 33
 - water availability, 33
 - management strategies, 35–37
 - temperature and precipitation conditions, 31
 - tolerant, 31
 - winter conditions, 31
- Collaborative management
- build trust, 425, 426
 - citizen-driven, 421
 - and education programs, 410
 - emergence of multiparty, 424
 - exotic plant management, 420
 - forming and maintaining, 425
 - land management issues, 420
- Common garden, 135, 149
- Communication, 448
- Community multiscale air quality (CMAQ), 240
- Computer simulation models, 441–443
- Conceptual models, 376, 377
- Cool-season grasses, 38
- Cool-season species, 38
- Coordination
- exotic annual invasive grass management,
447–448
 - and incentives, 450
- Cost-effective, 436, 444–447, 449, 452
- Cost minimization models, 436
- Cost-minimizing strategy, 435
- Cost-sharing across institutions, 424
- Cost-sharing programs, 448
- Creosote bush communities, 21
- Creosote bush scrub, 18, 19
- Cropland
- homestead laws, 315
 - roadsides, ditches and field margins, 316
 - technological improvements, 315
- Crusts
- biotic crust microsites, 63
 - cryptobiotic, 81
 - physical soil crusts, 71
 - soil crusts, 71, 79
- D**
- Damages, 432, 434, 436, 446, 447, 450
- Database, 159, 163, 164, 178–180
- categorization, 165
 - chromosome counts, 176
 - creation and population, 159
 - diploid/polyploid, 176, 177
 - frequency distribution, 181, 182
 - histograms, 164
 - human use and cultivar availability
 - categorization, 179
 - cultivar polar, 178
 - logistic regressions, 178
 - northern meadow types, 178
 - reseeding and burn rehabilitation, 180
 - soil stabilization, 180
 - taxonomy, 178

- management implications, 183
 - multiple logistic regressions, 165, 169
 - ploidy level, 176, 177
 - seed awns and mass, 172, 174–176
 - statistical analysis
 - biogeographic realms, 159
 - chromosome count and ploidy level, 163
 - crop weed, 163
 - cultivars and no cultivars, 163
 - dependent variables, 159
 - environmental weed, 163
 - forage/hay/revegetation, 163
 - GrassPortal, 164
 - independent variables, 163, 164
 - life span, 163
 - logistic multiple regressions, 164
 - negative impact, 159
 - polyploid, 163
 - ruderal weed, 163
 - seed mass values, 163
 - temperature and precipitation data, 164
 - univariate linear logistic regressions, 164
 - wide introduction, 159
 - taxonomy and life span, 165–172
 - temperature and precipitation, 180
 - univariate logistic regressions, 165–168
 - weediness, 165
 - Western United States, 184
 - year, 172, 173
 - Deep soil, 77
 - Defensible space, 448
 - Defoliation, 354
 - Dense layers, 65
 - Desertification
 - pattern, 87
 - Developmental plasticity, 134
 - Die-offs, 195
 - B. tectorum*, 65
 - eradication, 87
 - Discounting, 433, 434
 - Disturbance, 436, 439, 441–443, 450
 - Disturbance regimes
 - Cold Deserts, 32, 34, 35
 - Cold Deserts ecoregion, 12
 - ecological effects, 16
 - management implications, 47
 - Mediterranean California, 25–27, 30, 31
 - resilience to, 12
 - Warm Deserts, 20–22, 24
 - Western Forests, 44–45
 - Western Great Plains, 38–41
 - Diversity
 - and abundance of insects, 83
 - AMF fungi, 82
 - deep/tap rooting and shallow/fibrous-rooting plant species, 65
 - native species richness, 63
 - and pollination activity of insect pollinators, 84
 - DNA sequence data, 101
 - Drought disturbance, 401
 - Dryland, 228
 - Dynamic optimization
 - analyze, 434
 - control variables, 435
 - cost minimization models, 436
 - decision-maker, 435
 - description, 434
 - and dynamic simulation models, 434
 - and imperfect information, 438–439
 - management actions, 434
 - management of *Bromus*, 434
 - natural processes, 434
 - OC methods, 435, 436
 - ranch management models, 436–437
 - SDP, 435, 436
 - state variables, 435
 - stochastic processes, 437–438
 - time horizon, 435
- E**
- Ecological genetics, 134. *See also* Adaptive evolution
 - Ecologically-based invasive plant management (EBIPM) program, 422, 423
 - Ecological process
 - abiotic and biotic attributes, 12
 - Ecological resilience, 441
 - Ecological site description (ESDs), 373
 - Ecological states, 430, 434, 437, 439, 440, 444, 445, 450
 - Ecological threshold, 443
 - computer simulation models, 441–443
 - critical point, 439
 - economics analyzes, 439
 - natural resource management problems, 439
 - and optimization models, 440–441
 - rangeland policy, 439
 - resilience, 441
 - self-organizing processes, 439
 - spatial interactions (*see* Spatial interactions)
 - uncertainty, 443
 - Economic benefits, 443, 448–450, 453
 - Economic costs, 412

- Economic model, 434–439, 451–453
 and ecological, 430
 and ecological dynamics (*see* Dynamics optimization)
 bio-economic, 431–434
 Efficient *Bromus* Management, 449
 incentives and coordination, 450
 research
 management issues, 452–453
 modeling assumptions, 451–452
- Ecoregions, 17–47
 cold deserts, 12 (*see also* Cold deserts)
 Mediterranean California
 (*see* Mediterranean California)
 research, 48–49
 Warm deserts (*see* Warm deserts)
 Western forests (*see* Western forests)
 Western great plains
 (*see* Western great plains)
 in western US, 12–14, 17
- Ecosystem goods and services, 452
- Ecosystems, 63–68, 73–85
 animals dependent, 83
 changes in bird, 86
 fire frequency, 62
 fluxes and energy partition
 Bowen-ratio approach, 74
 Ecosystem surface temperatures, 74
 energy balance, 73
 Kansas soils, 76
 latent heat and CO₂ on *BRTE*-dominated/sagebrush-steppe communities, 74, 75
 lower ET, 74
 NCE and ET, 76
 net radiation balance, 74, 75
 net radiation fluxes, 73
 plot-level chamber measurements of NCE and ET, 74
 reduced NCE, 74
 soil C storage, 76
 solar radiation, 74
 spatial and temporal scale, 74
 forage quality, 83
 habitat transformation, 83
 invertebrates and small vertebrates
 ants/ground squirrels, 83
 bromus-dominated plots, 84
 burrow densities, 85
 ground squirrel relationships, 85
 locomotion, 85
 loss of native forbs and pollinator community, 84
 negative relationships, 84
 nutrition, 85
 omnivorous species, 84
 seed-harvesting genera, 83
 shelter, 85
 landscape disturbance, fire, 68–69
 large-mammal communities, 86
 management implications, 86–87
 microfaunal communities, 82–83
 native- and *Bromus*-dominated communities, 62
 plant communities (*see* Plant communities)
 research needs, 87–88
 soil hydrology and water balance, 71–73
 soil microbiological, 82–83
 soil nutrients (*see* Soil nutrients)
 soil stability, 69–71
 structure and function, 62
- Ecosystem services, 431, 439
- Ecosystem threats
 cold deserts, 34–35
 Mediterranean California, 28–30
 Warm deserts, 21–23
 Western great plains, 42
- Education, 421–422
- Education programs, 410, 412, 422
- Effectiveness, 432, 437, 444, 453
- El Niño–Southern Oscillation (ENSO) events, 32
- Energy balance, 73, 74
- Environmental risks, 412
- Environmental tolerance
 genotypes, 136
 phenotypic plasticity, 141–142
- Enzyme electrophoresis (allozymes), 104, 105, 109, 112, 117, 121, 125
- Eradication, 437, 446, 451
- Erosion
 direct effects, 70–71
 grazers, 76
 infiltration, 87
 topsoil, 72, 77
- ESDs. *See* Ecological site description (ESDs)
- Evapotranspiration, 70, 71, 73, 74
- Evolution. *See* Adaptive evolution
- Executive Order 13148, 418
- Exotic annual *Bromus* and human dimensions
 annual grasses, 325
 Euro-American settlements, 308
 function of, 308
 governmental regulations and policies, 309
 invasive species, 325
 (*see* Land uses and policies)
 rapid population growth, 308 (*see also* Sagebrush-dependent wildlife)
 structure and food resources, 325

- Exotic annual *Bromus* colonization, 315–316, 318–321
B. tectorum and *B. rubens*, 315
 cropland (*see* Cropland)
 ecosystem resilience and resistance, 314
 energy developments, 324
 extracted natural resources, 324
 land management policies, 317–318
 livestock grazing (*see* Livestock grazing)
 Montana Greater Sage-grouse Habitat
 Conservation Advisory Council, 324
 power lines, 325
 resilient lands, 314
 (*see also* Transportation corridors and vehicles) (*see also* Urban, suburban and exurban development)
 vegetation clearing is, 325
 weed policies and exotic annual *Bromus* (*see* Weed policies and exotic annual *Bromus*)
 well pads, 324
- Exotic annual *Bromus* grasses. *See also*
 Adaptive management framework
 adaptive management frameworks, 340
 agroecosystems, 340
 arid and semiarid ecosystems, 340
Bromus arvensis, 342–343
Bromus diandrus, 342
Bromus hordeaceus, 341–342
Bromus rubens, 343
Bromus rubens L. (red brome), 4
Bromus tectorum, 4, 343–344 (*see also* *B. tectorum* research)
 characteristics, 344–345
 human and economic dimensions and management options, 6
 invasions in Western USA, 341
 management implications, 360–361
 net primary production and annual precipitation, 341
 organismal traits, 5
 patterns and impacts of invasion, 5
 research needs, 361–362
 research, planning and management, 2
 resilience, 2–4
 resistance, 2–4
 soil, climate and plant community controls, 5
 spectrum, 1–2
 US Department of Agriculture's *Bromus* REENet, 6
- Exotic annual grass management
 policy influences, 418–420
- Exotic invasive species management, 420–421
 institutions (*see* Institutions)
- F**
- Fall Imazapic treatment
B. diandrus, 23, 24
- Feedback
 climate of ecosystem, 74
 self-perpetuating, 82
- Fire, 402–403
 wildfire, 12
- Fire regimes
 annual grass/fire cycle, 312
 cold deserts ecoregion, 34
 ecoregions, 311
 ecosystems, 310
 environmental/productivity gradients, 310
 fire-dependent ecosystems, 30
 frequency, 20, 21, 25, 32
 and fuelbed changes, 311
 grass/fire cycle, 16, 21, 22, 29, 48
 high-intensity fires, 31
 history, 27
 and human disturbance, 28
 and lower post-fire resilience, 28
 management strategy, 24
 and management treatments, 36
 Mediterranean California region and Palouse Prairies, 310
 perennial herbaceous species, 313
 perennials, 310
 post-fire landscapes of eastern Mojave Desert, 20
 post-fire seeding, 24
 severity, 26
 short-interval low-intensity fires, 29
 size, 21, 33
 Warm desert ecoregion, 311
 Western forests, 47
 Western great plains, 38, 39
- Fitness
 environmental factors, 141
 unfavorable and favorable environments, 142
- Floral characteristics, 103, 104
- Florets, 194
- Forage, 83, 85, 86, 431, 432, 435, 439, 447, 452
- Forest, 14
 chaparral and mixed conifer, 25
 clear-cut harvests, 16
 fire-sensitive tropical forests, 29
 inland mountain, 25
 oak woodlands and mixed conifer, 28
 ponderosa pine, 31
 Western forests ecoregions (*see* Western forests)
- Founder effects, 107, 122, 123
- Fuel treatment, 445, 452

Fuels

- accumulations, 28
- fire, 69
- flashy, 29
- fuelbeds, 46
- herbaceous, 29
- load, 31
- loads, 21
- loss of woody, 69
- moisture, 29
- in Mojave Desert, 68
- perennial, 68
- pre-fire fuels manipulations, 47
- production, 68
- structure, 16
- wildfire, 66
- woody, 35

Funds, 433, 441, 442, 448–450, 452

Fusarium

- ascomycete, 210
- B. tectorum* seeds, 210
- die-off, 210
- distribution and epidemiology, 213
- fungal organisms, 210
- host ranges, 213
- life cycle, 211–213
- macroconidia, 211
- molecular-genetic characterization, 210
- natural ecosystems, 210
- seed rot, 210

Fusarium nivale, 195

G

GCM. *See* Global circulation models (GCM)

Genea, 157, 163–165, 170–172, 183, 184

- Bromus*, classification, 101
- diploid species, 102
- evolutionary history, 101
- genus *Bromus*, 100
- hypothesized origins of the polyploid species, 102
- invasiveness of *B. tectorum*, 101
- monophyletic, 102
- patterns of speciation, 101
- phylogenetic relationships, 101
- phylogenetic tree, 102
- Poaceae, 101
- polyploid series, 100

Gene flow, 134

Genetic admixtures, 107, 108, 113, 115, 117, 118, 120, 122–124

Genetic diversity, 139, 146

B. tectorum in NA, 109

- enzyme loci, 105
- level and structure, 101
- and MLGs, 109
- native and invasive populations, 121
- populations, 103, 108, 115
- populations of *B. tectorum*, 122–123

Genetic surveys

population, 105

Geographic origins (source populations), 121–122

Global circulation models (GCM), 391, 392

Grass

- array, 100
- destructive, 100
- fire cycle, 21–23, 29, 47, 48
- immigrants, 100

Grassland

- annual, 22
- California, 29
- and chaparral, 43
- desert grasslands, 20
- Exotic annual, 26
- grazing, 16
- Mediterranean California, 25
- native perennial, 30
- perennial, 25
- semiarid desert perennial, 17
- serpentine, 28
- Western great plains, 40

Grazing, 430, 442, 450

Great Basin surveys, 414

Greening the Government Through Leadership in Environmental Management, 418

H

Herbarium specimens, 109, 113, 120

Herbicide application, 354–355, 416

Herbicides, 23, 35, 36, 43, 48, 414–416, 423–425

Herbivory

- B. tectorum*, 293
- and community resistance, 294
- disturbance regimes and community resilience, 294–295
- soil and plant community disturbance, 292

Heterozygous, 105

Human dimensions, 411–421

- attitudes (*see* Attitudes)
- behavior change strategies, 411
- behaviors (*see* Behaviors, human dimensions)
- Beliefs (*see* Beliefs, human dimensions)
- biology and management of exotic invasive species, 410

- classification, 409
- directly/indirectly affect, 410
- education, 421–422
- institutions (*see* Institutions)
- management implications, 422–425
- natural resource management centers, 411
- policies (*see* Policy)
- policy sciences, 410
- published studies, 410
- research, 425, 426
- social-psychological research, 410, 411
- society invests, 409
- Hydrology
 - soil, 71–73
- I**
- ILAP. *See* Integrated Landscape Assessment Project (ILAP)
- Imperfect information
 - and dynamic optimization, 438–439
- Imperfect knowledge, 438, 451
- Incentives, 431, 434, 447, 448
 - and coordination to enhance management, 450
- Infiltrations
 - collaboratives, 420
 - and erosion, 87
 - mobilizing citizen groups, 421
 - and nutrients, 88
 - HYDRUS to partition rain, 70
 - precipitation, 71
 - public agencies, 420
 - and retention in semiarid soils, 77
 - and runoff in California grasslands, 72
 - soil, 70
 - and soil hydraulic conductivity, 71
 - volunteer groups, 420
 - weed programs, 421
- Integrated Landscape Assessment Project (ILAP), 397
- Interdependencies, 430
 - spatial (*see* Spatial interdependencies)
- Interviews, 413
- Invasion potential
 - annual minimum and maximum temperature and precipitation, 14, 15
 - cold deserts, –, 32, 34
 - Mediterranean California, –, 25, 28
 - resilience, 12–16
 - resistance, 12–16
 - Warm deserts, 19–21
 - Western forests, 44–45
- Invasiveness, 135
 - breeding and cultivation, 185
 - C3 photosynthetic pathway, 157
 - cereal grasses, 157
 - climatic factors, 185
 - databases, 157
 - distribution and evaluation, 183
 - environments, 183
 - genetic and demographic characteristics, 157
 - hypotheses, 157, 160–162
 - Mediterranean climate, 157
 - natural environments and human agricultural systems, 156
 - patterns, 185
 - plant taxa and types, 156
 - and plant traits, 183
 - prediction, 156
 - REENet project, 157
 - sagebrush ecosystems, 156
 - scientific principles/reliable procedures, 156
 - section *Bromus*, 185
 - taxonomists, 157, 158
- Invasive species
 - costs, 434
 - hypothetical, 448
 - imperfect information and dynamic optimization, 438–439
 - management costs, 436
- Invertebrates
 - and small vertebrates, 83–85
- K**
- Kansas soils, 76
- L**
- Landscape disturbance, fire
 - fire-return intervals, 68
 - fuel (litter) production, 68
 - mechanisms, 68–69
- Landscape Fire and Resource Management Planning Tools Project (LANDFIRE), 378
- Landscape-scale management approach, 47
- Land treatment digital library (LTDL), 360
- Land uses and policies
 - Bromus* invasion, 309
 - community dominance, 309
 - ecosystems, 309
 - vegetation management tool, 309
- Learning model
 - experience-weighted attraction, 438

- Legacy effects
Bromus, 66
- Litter accumulation
 community structure, 65–68
- Littledalea*, 157, 158
- Livestock, 16, 17, 21, 28, 30, 33–35
 attitudes, 414
 grazing, 416
 grazing and herbicide application, 414
 operations, 86
 preferences, 86
 wildlife and domestic, 62
- Livestock grazing
 clonal growth, 319
 mismanagement of, 320, 321
 morphological adaptation, *B. tectorum*, 319
 native grasses, 319
 plant communities, 319
 plant species and communities, 318
 tolerance, 318, 320
 trampling, 320
 uncontrolled livestock, 318
 in western USA, 318
- Local adaptation
 genetic variation, 139–141
 habitats, 138
- Low copy nuclear gene *Waxy*, 101
- LTDL. *See* Land treatment digital library (LTDL)
- M**
- Magnesium (Mg), 82
- Management
 belief-attitude-behavior research, 411
 public lands, 411
- Management, human dimensions, 413
 attitudes (*see* Attitudes)
 and beliefs about exotic invasive plants,
 412–413
- Management strategies
 cold deserts, 35–37
 landscape-scale management approach, 47
 Mediterranean California, 30–31
 prevention, 48
 restoration, 48
 Warm deserts, 23, 24
 Western forests, 47
 Western great plains, 43
- Manganese (Mn), 82
- Maternal effects, 141
- Mating systems, 103–108
 Baker's list, 103
 biological invasions
 (*see dee* Biological invasions)
- biology and genetics of plant
 populations, 103
- Bromus tectorum*
 floral characteristics, 103, 104
 genetic markers, 104, 105
 outcrossing rate, 105–107
 seed set, 103
 sex allocation theory, 104
- The evolution of weeds, 103
 genetic diversity, 122–123
 management implications, 123–124
 plant mating systems, 103
 research needs, 124–126
 selfing mating system, 103
 source populations, 121–122
 weed's potential mating system, 103
- MCG
 high frequency in Central USA, 112
- Mediterranean annual grasses, 1
- Mediterranean California, 25–29
 ecosystem threats, 28–30
 grasslands, 25
 invasion potential
 anthropogenic factors, 26
 atmospheric pollution in nitrogen
 deposition, 27
 canopy shrublands, 27
 chaparral sites, 26
 chaparral with fire histories in southern
 California, 27, 29
 ecosystems, 28
 exotic annual grasslands, 26
 fire history, 27, 29
 meta-analysis, 28
 natural disturbance factors, 28
 nitrogen addition, 28
 oak savannas, 28
 oak woodlands and savannas, 25
 resilience to historical and altered fire
 regimes, 25, 26
 resistance, 28
 resistance to common *Bromus*, 25, 26
 sage scrub, 25
 management strategies, 30–31
 soil temperatures, 25
- Mice
 deer mice, 84
 and reptiles, 85
- Microbes
 C:N of organic substrates, 79
 live, 82
 phosphatases, 81
- Microdochium nivale*, 195
- Microfaunal communities, 82–83

- Microsatellite, 138
 Microsatellite DNA, 104
 Missoula residents, 415
 MLGs
 in Central USA, 112
 detection, 112
 Eastern Canada, 112
 Got-4c, 112
 low frequency in populations in the Central USA, 112
 recombinant genotype, 112
 Model uncertainty, 384
 Moisture regimes
 classification, 25
 and drier soil temperature, 43
 ecosystem resistance and resilience, 17
 resistance, 13
 soil, 17
 temperature, 31, 32
 Monitoring, 445
 Monte-Carlo random sampling methods, 385
 Most common genotype (MCG), 109, 113
 California and the American Southwest, 118
 in East Coast populations, 109–110
 Eastern Canada, 121
 high frequency (52 of 54 populations) in the mid-continent USA, 115
 high frequency in the Central USA, 112
 northern portion of the mid-continent USA, 115
 Western US populations, 117
 Multilocus genotypes (MLGs)
 B. tectorum populations, 117
 detected in Eastern US populations, 112
 distribution, 109
 Got-4c, 112
 identified, Western US populations, 117
 low frequency in populations in the Central USA, 112
 lower frequency, 117
 Mdh-2b & *Mdh-3b*, 109
 in Northern California, 118
 Pgm-1a & *Pgm-2a*, 109, 112, 115
 Pgm-1ab & *Pgm-2ab* heterozygous, 115
 populations of *B. tectorum* from California and the American Southwest, 118
 25 populations of *Bromus tectorum* in British Columbia, 119
 38 populations of *Bromus tectorum* in the Eastern USA, 110
 40 populations of *Bromus tectorum* in the Western USA, 116
 51 populations of *Bromus tectorum* in Canada, 114
 60 populations of *Bromus tectorum* in California and the American Southwest, 118
 60 populations of *Bromus tectorum* in the Central USA, 111
 60 populations of *Bromus tectorum* in the mid-continent USA, 113
 restricted distributions, 115
 South Dakota or Eastern Wyoming, 115
 Tpi-1b, 118, 120
 Multiple introductions, 101, 107–109, 117, 121–124, 134
 genetic diversity, 101
 Mycorrhizal fungi, 82

N
 NAISM. *See* North American Invasive Species Management Association (NAISM)
 National Environmental Policy Act (NEPA), 388
 Native American land uses
 Euro-American colonization, 314
 evidences, 314 (*see also* Exotic annual bromus colonization)
 paleo-botanical and archeological records, 313
 tree removal, 313
 wildlife, 313
Neobromus, 157, 163–165, 171, 184
 Net ecosystem exchange, 73
 Neutral genetic variation, 139, 141
Nevskiella, 157, 158, 163–165
 Nitrogen (N)
 addition, 25, 28
 atmospheric deposition, 21
 B. tectorum, 241
 deposition, 21, 27, 63
 fertilization, 240, 241
 perennial grasses, 241, 242
 pollution, 27
 shortgrass steppe, 242
 soil nitrogen cycling, 78–81
 soil phosphatase, 242
 Wyoming big sagebrush community, 241
 Nitrogen (N) deposition
 adaptive and evolutionary responses, 146
 ecological effects, 146
 Nongovernmental organizations (NGOs), 424
 North American Invasive Species Management Association (NAISM), 316
 Novel genotypes
 creation, 147
 implications, 148

Noxious Weed Control and Eradication Act, 418
 Noxious weed designation, 419, 424
 Nuclear ribosomal DNA (rDNA) regions,
 101–102
 Nutrients, 242

O

Objective function
 bio-economic model, 432
 decision-maker's, 432, 435
 time horizon, 435
 Objective function/decision-maker's, 430
 Oklahoma cooperative extension service, 39
 Opportunity cost, 433
 Optimal control (OC), 435, 436, 440, 441
 Optimal control (OC) methods, 435
 Optimization, 435
 dynamic (*see* Dynamic optimization)
 Optimization models
 and ecological thresholds, 440–441
 Organic matter, 65, 69, 71, 72, 77
 Outcrossing mating system, 103–108, 112

P

Palmer drought severity index (PDSI), 393,
 400, 401
 PDSI. *See* Palmer drought severity index
 (PDSI)
 Peavine Mountain population, 106
Pennisetum ciliare, 381, 415
 Perennial grass, 20, 449
 and forbs, 449
 and sagebrush, 439, 440
 Perfect foresight, 433, 452
 Phenology, 65, 72, 79
 growth and reproduction, 142
 and seed weight, 141
 Phenotypic plasticity
 germination, 142
 habitats, 142
Tamarix ramosissima, 142
 Phosphorus, 77, 81–82, 242
 Pinyon-juniper, 448, 453
 Plant communities, 64–68
 disturbances, 63
 Mojave basin and range, 63
 monocultures or exotic annual/biennial, 63
 sagebrush steppe, 63
 spatial and temporal structure, 63
 structure
 accumulation, 65–68
 annual or biennial exotics, 65

deep/tap rooting and shallow/
 fibrous-rooting plant species, 65
 distribution of *Bromus tectorum*'s
 dominance, 65, 66
 instability, 65
 phenology, 65
 soil interspaces, 64
 spatiotemporal effects, 65
 Wyoming big sagebrush site, 63, 64
 Plant community resistance, 278–284
Bromus, 276, 277
 competition/facilitation, 284
 disturbance and stressors, 277
 environmental factors, 276
 arid and semiarid ecosystems, 279
 competitive interactions, 280
 deep soil water, 277
 evapotranspiration, 277
 higher resource availability and plant
 productivity, 278
 plant growth, 278
 seasonality of precipitation and
 temperature, 278
 soil temperature and moisture
 regimes, 281
 summer precipitation, 278
 water availability, 278
 woody and herbaceous species, 277
 environmental gradients, 278
 forbs, 284–285
 management implications, 295
 perennial grasses
 bunchgrasses, 281
 ecoregions, 281
 forage production, 282
 gap size, 281
 life stages, 282
 sandy loam soils, 282
 short-grass steppe, 282
 winter/spring precipitation, 281
 plant traits and resource dynamics
 canopy and litter effects, 280
 growth/maintenance resource
 pool, 280
 nitrogen, 280
 nitrogen availability, 280
 nutrients, 279
 perennial grasses, 280
 root length and biomass, 280
 seasonal priority advantage, 279
 seed entrapment and germination, 280
 shallow soil water, 280
 soil resource, 279
 spatial and temporal patterns, 279

- winter/spring precipitation, 279
 - woody plants, 279
 - research needs, 296
 - resilience, 276
 - resistance to invasion, 276
 - woody plants
 - B. rubens*, 284
 - biomass and seed production, 283
 - bunchgrass, 283
 - cold desert, 283
 - factors, 283
 - seeding experiment, 282
 - shrubs, 283
 - temperatures limitation, 283
 - trees and forest canopies, 284
 - Plant invaders
 - negative consequences, 100
 - Plant Protection Act of 2000, 418
 - Poaceae, 156
 - Policy
 - Asian Carp Prevention and Control Act of 2010, 418
 - BLM, 418
 - categorization, 419
 - discrepancies, 420
 - federal land management
 - agencies, 418
 - futility, 420
 - mechanisms, 418
 - Noxious Weed Control and Eradication Act, 418
 - noxious weed designation, 419
 - noxious weed lists, 419
 - Plant Protection Act of 2000, 418
 - policy-makers, 451
 - and programmatic decisions, 445
 - Public, 450
 - question, 430
 - rangeland, 431, 437, 439
 - regulatory policy, 419
 - SAFE Transportation Equity Act of 2005, 418
 - source populations, 419
 - US Department of Agriculture, 420
 - US Geological Survey, 418
 - utility, 420
 - utility and futility, 420
 - Polymorphic marker system, 106
 - Pooideae, 157
 - Postfire revegetation, 442, 449
 - Post-immigration evolution, 107–108, 125
 - Potassium (K), 82
 - Preadaptations, 103, 107–108
 - Preadapted genotypes, 108
 - Precipitation
 - and evapotranspiration, 146
 - N-limited habitats, 146
 - winter and spring, 147
 - Private land, 430, 448, 450
 - Profit maximization, 452
 - Profits
 - average annual, 433
 - higher ranch, 450
 - maximization, 452
 - maximize, 431
 - rancher's profits, 432
 - streams, 433
 - Progeny array analyses, 104
 - Propagule pressure, 107–109, 122, 123
 - Public and private land managers, 430
 - Public land management agencies, 430
 - Public land managers, 415, 447, 451
 - Pyrenophora semeniperda*, 204–209
 - black fingers of death pathogen, 204, 205
 - death genetics, 209–210
 - distribution and epidemiology
 - annual bromes, 207
 - B. tectorum*, 208
 - cytochalasin B, 209
 - inoculum addition experiments, 209
 - mean killed seed density, 207
 - mortality, 208, 209
 - mycelial growth and toxin
 - production, 209
 - pathosystem, 208
 - seed bank, 208
 - host ranges, 206–207
 - life cycle
 - aerial inoculum spray method, 205
 - B. tectorum* plants, 206
 - floral infection, 206
 - mature nondormant seeds, 204
 - race for survival, 204
 - ring spot, 206
 - seed germination rate, 204
 - sexual structures (perithecia), 204
 - toxic compounds, 206
- Q**
- Quantitative trait, 135
- R**
- Race for survival, 204
 - Radiation, 66, 69, 73, 74
 - longwave, 74
 - net radiation, 73

- Radiation (*cont.*)
 net radiation balance, 75
 radiation balance, 69
 solar radiation, 74
 terrestrial radiation, 66
 thermal radiation, 74
- Ranchers, 430–433, 435, 437, 440, 441, 448, 450–452
- Ranch management models, 436–437
- Range expansion, 139, 148
- Rapid ecoregional assessments (REAs), 356, 357
- REAs. *See* Rapid ecoregional assessments (REAs)
- Reciprocal transplant
 adaptation, 138
 adaptive differentiation, 135
 local adaptation, 135
- Recombinant genotypes, 107, 112, 117, 120, 125
- Regulatory policy, 419
- Rehabilitation, 430, 432, 434, 437, 440, 442–445, 449, 450, 452
- Reproductive traits. *See* Plant community resistance
- Research, Education, and Extension (REEnet) project, 157
- Resilience, 21–24, 26, 28–30, 32–36, 39, 41, 276
 cold deserts
 ecosystem threats, 34, 35
 invasion potential, 32–34
 management strategies, 35, 36
 definition, 3
 disturbances, 2
 ecological, 441
 ecosystem, 3
 environmental factors and ecosystem attributes and process, 4
 invasion potential, 12–16
 management strategies, Western forests, 47
 Mediterranean California
 ecosystem threats, 29, 30
 invasion potential, 25–28
 management strategies, 30
 native ecosystems, 3
 Warm deserts
 ecosystem threats, 21–23
 management strategies, 24
 Western great plains
 ecosystem threats, 42
 invasion potential, 38–41
- Resistance, 20–22, 24, 25, 28, 32–36, 38–40, 42, 44–47
 and abiotic and biotic attribute functions, 3
 cold deserts
 ecosystem threats, 34, 35
 invasion potential, 32–34
 management strategies, 35, 36
 definition, 3
 ecosystem attributes and processes, 2
 environmental factors and ecosystem attributes and process, 4
 invasion, 3
 invasion potential, 12–16
 invasive *Bromus* grasses, 2
 Mediterranean California
 invasion potential, 25–28
 plant community, 5
 Warm deserts
 ecosystem threats, 21, 22
 invasion potential, 20, 21
 management strategies, 24
 Western forests
 ecosystem threats, 46
 invasion potential, 44, 45
 management strategies, 47
 Western great plains
 ecosystem threats, 42
 invasion potential, 38–41
 management strategies, 43
- Restoration, 416, 417, 421–426
 active, 30
 fire regimes, 47
 guidelines in Warm Deserts, 24
 landscape-scale management approach, 47
 passive, 30
 postfire, 445
 post-wildfire, 442
 practices, 438
 rangeland, 447
 resilience and resistance, 48
 restoration-based fuel treatment, 445
 success rate, 445
 success rates, 443
 technology, 439
 treatments, 441
 wildfire, 452
- Restoration success rates, 443, 445
- Reticulate pattern of polyploidy speciation, 102
- Revegetation, 355, 442, 449
- RFLP types, 82
- Risk aversion, 433
- Robust management, 441
- Root exudates, 81
- Rooting, 65, 71, 77
- Rooting systems, 72
- Rule of thumb, 432, 452

- Runoff, 70–72
- Rutstroemiaceae
- ascomycete pathogen, 214
 - bleach blonde syndrome, 214, 215
 - distribution and epidemiology, 216
 - greenhouse pathogenicity tests, 214
 - host ranges, 216
 - life cycle, 214–216
 - molecular-genetic techniques, 214
 - sterile plants, 214
- S**
- SAFE Transportation Equity Act of 2005, 418
- sagebrush, 439, 440, 442, 445, 448–450, 453
- Sagebrush-dependent wildlife
- communities, 326
 - C. urophasianus* populations, 327
 - habitat, 326
 - life stages, 326
 - management implications, 327–329
 - perennial herbaceous plants, 327
 - reduction treatments, 327
 - research, 329–330
 - woody plant reductions, 326
- Sage grouse, 83, 86
- Saltbush scrub, 20, 21
- Salt desert vegetation types, 32
- Scotts Bluff National Monument in
northwestern Nebraska, 38
- Search, 447
- Seed banks
- abiotic and biotic factors, 289
 - burns and wildfires, 289
 - dispersal, 289
 - germination, 289
 - growing seasons, 290
 - livestock grazing, 290
 - Mojave desert community, 290
 - peak densities, 289
 - propagule pressure, 289
 - soil surface/litter, 288
 - Wyoming big sagebrush community, 290
- Seed characteristics
- adhesion, 287
 - animals, 287
 - birds eat seeds, 288
 - caryopses, 288
 - diaspores, 287
 - herbivores, 288
 - heteromyid rodents, 288
 - lemma awns, 286
 - microsite conditions, 288
 - moisture and temperature conditions, 287
 - physical and biotic factors, 287
 - sterile florets, 287
 - wind and water dispersal, 287
 - world planted regions, 288
- Seed dispersal and germination, 291–292
- Seed germination, 290–291
- Seeding
- perennial herbaceous species, 35
 - post-fire, 24
- Seed production and propagule supply,
285–286
- Seed set, 103
- Selfing, 103–106, 108, 125
- Selfing rates, 106
- Self-pollinating (selfing) mating system, 103
- Sex allocation theory, 104
- Shortgrass steppe, 36, 38
- Shortgrass steppe ecosystem, 40
- Shrubland
- blackbrush, 18, 24
 - canopy, 27
 - chaparral, 25
 - coastal, 30
 - cold deserts, 34, 45
 - ecosystems, 27
 - fire-prone shrubland ecosystem, 29
 - and grasslands, 16, 43
 - Mediterranean, 26
 - native, 22, 28
 - semiarid, 27
 - soil carbon storage, 29
 - vegetation types, 17
- SIC, 77
- Single nucleotide polymorphisms (SNPs),
106, 139
- SOC, 77, 78
- Social benefits, 431
- Social costs, 431
- Social desirability bias, 413
- Social-psychological research, 410
- Soil aggregates, 72, 77, 82
- Soil biogeochemical factors
- annual grasses, 228
 - arid and semiarid ecosystems, 228
 - B. tectorum* and *B. rubens*, 229
 - Chihuahuan desert, 248
 - climate and soil factors, 247, 248
 - dryland, 228
 - Great Basin/Columbia Plateau, 249
 - management implications, 250
 - Mojave desert, 249
 - P-limitation model, 250
 - regression model, 249
 - research needs, 250

- Soil biogeochemical factors (*cont.*)
 - soil organic matter, 249
 - soil texture and depth, 250
 - SOM levels, 249
 - W/T ratio and total winter precipitation, 248
 - water and nutrient availability, 229
 - Western USA, 229
- Soil C in topsoil vs. subsoil, 77–78
- Soil controls
 - B. rubens* and *B. tectorum*, 232
 - Chihuahuan desert, 232
 - climatic regions, 232
 - coastal sage scrub, 239
 - Colorado Plateau Desert, 234, 238
 - Great Basin/Intermountain Region and Columbia Plateau, 238, 239
 - Mojave desert, 233
- Soil food web
 - Bromus*-invaded *P. jamesii* grasslands, 83
 - components, 82
- Soil hydrology and water balance
 - burned and invaded areas, 72
 - infiltration, 72
 - microsite types, 72
 - percolation, 71
 - physical properties, 72
 - pivotal model parameters, 71
 - reduced plant cover, 71
 - spatial and temporal patterns, 72–73
- Soil microbiological, 82–83
- Soil moisture, 243
- Soil nitrogen cycling, 78–81
- Soil nutrient and water
 - calcium/magnesium carbonates, 230
 - decomposition and nutrient transformations, 230
 - depositional zones, 231
 - desert soils, 230
 - dryland soils, 230
 - erosional processes, 231
 - fire/heavy grazing, 232
 - K/Mg ratios, 230
 - macronutrients, 229
 - micronutrients, 229, 230
 - nitrogen, 230
 - patchwork pattern, 231
 - plant nutrition, 230
 - shrub canopies, 231
 - soil texture, 230
 - SOM, 231
- Soil nutrients
 - calcium (Ca), 81, 82
 - disturbances and soil factors, 76
 - elemental cycling, 76
 - nitrogen cycling, 78–81
 - phosphorus, 81–82
 - Soil C in topsoil vs. subsoil, 77–78
- Soil organic matter (SOM), 231
- Soil resources, 67
- Soil stability
 - annual horizontal flux, 70, 71
 - B. tectorum* die-off in the Northern Basin and Range, 67, 69
 - direct effects on erosion, 70–71
 - ecosystems, 69
 - herbicide Oust®, 69
 - stabilization, 69
 - temporary losses of plant cover, 69
 - wind erosion, 69
- Soil texture and biochemistry
 - alkaline soils, 246
 - CaCO₃ solubility, 245
 - calcareous soils, 245
 - climate factors, 245
 - disparate organisms, 246
 - disturbance/management treatments, 244
 - exotic annual grasses, 246
 - factors, 244
 - germination and establishment phases, 244
 - H₂CO₃ formation, 245
 - hotter and cooler regions, 244
 - low pH, 244
 - micronutrients, 246
 - osmoregulation, 246
 - photosynthesizing tissues, 246
 - potassium, 246
 - root CEC, 245
 - root growth, 245
 - sandstone, 245
 - shallow sandy soils, 247
 - soil depth and texture, 247
 - soil factor controlling, 244
 - soil fungi and plants, 245
- Soil water, 73, 79, 81
- SOM. *See* Soil organic matter (SOM)
- Source populations, 121–122
- Southwest survey, 414
- Spatial interactions
 - coordination of exotic annual invasive grass management, 447–448
 - dynamic ecological and economic processes, 443
 - interdependencies between locations, 444
 - interdependency of biophysical processes, 446–447
 - level and importance of spillover effects, 444
 - management actions, 444
 - management benefits and costs, 444

- management efforts, space with resource constraints, 444–445
 - management resources, 443
 - spatial interdependencies in biophysical systems, 444
 - Spatial interdependency
 - biophysical and human systems, 430
 - biophysical processes, 446–447
 - biophysical systems, 444
 - and unique features for specific sites, 445
 - Spillover effects, 444
 - Spread, 430, 434, 438, 444, 446–448, 452
 - Squirrels, 83, 85
 - Squirreltail, 70
 - Stand failure, 65, 67, 69, 70
 - State-and-transition models (STM), 441, 442
 - biotic/abiotic thresholds, 375
 - Bromus* invasion and community dominance, 376
 - Bromus tectorum* L. (cheatgrass/downy brome), 373
 - disturbance/stress, 373
 - ecological system, 372
 - ESDs system, 373
 - graphical representation, 376
 - natural disturbances, 375
 - natural resources, 376
 - NRCS, 376
 - phases, 373
 - plant communities, 373
 - plant community, 373
 - soil resource, 373
 - thresholds, 373
 - vegetation community, 374–375
 - State-and-transition simulation modeling (STSM), 380–388
 - approaches, 377, 378
 - conceptual developments and terminology, 377–378
 - cost-effective approach, 380
 - ecological and economic, 372
 - ecological research needs, 399, 400
 - ecological systems, 372
 - LANDFIRE, 378
 - management implications, 398, 399
 - model uncertainty, 380
 - model uncertainty and conservation ecosystem dynamics, 384, 386
 - Eucalyptus* forests, 384
 - expert opinion, 384
 - experts, 385, 386, 388
 - imperfect knowledge, 385
 - ineffective/wasted management, 384
 - Monte-Carlo random sampling methods, 385
 - R software, 385
 - replicate models, 387
 - sources, 384
 - system stochasticity, 385, 387
 - temporal transition multipliers, 387
 - variance components analysis, 385, 387
 - variation among experts, 385
 - VDDT model, 385
 - VDDT models, 387
 - natural resource management problems, 377
 - non-*Bromus* systems, 380
 - nonspatial and spatial model approaches, 378, 379
 - plant monitoring and treatments
 - aerial photography, 381
 - budgets, 382
 - buffelgrass, 381
 - control measures, 380
 - exotic plant invasions, 380
 - land management programs, 380
 - land managers lack tools, 380
 - limited and unlimited budgets, 382
 - management strategy, 382
 - P. ciliare*, 381
 - resource allocation, 380, 381, 383
 - TELSA, 381
 - trade-off, 383
 - resource allocation, 372
 - spatial/nonspatial analyses, 377
 - spatial/non-spatial analyses, 372
 - spatially explicit, 378
 - state classes, 377
 - ST-Sim, 377–, 378
 - usage and benefits, 379, 380
 - VDDT, 377–, 378
 - States, 373–375
 - STM. *See* State-and-transition model (STM)
 - Stochastic dynamic programming (SDP), 435–438, 440
 - Stocking rates, 432, 433, 437, 438, 441, 447, 450
 - STSM. *See* State-and-transition simulation modeling (STSM)
 - Subsidies, 440, 441
 - Surveys, 413
 - cost-effective survey, 445
- T**
- Taraxacum officinale*, 246
 - Taxonomic classification, 100

- Taxonomy**
 annual/perennial, 165, 170
 archaeophytes, 170
Bromopsis, 171
 cereal crops, 171
Genea and *Bromus* crop weeds, 171
Genea and *Bromus* species, 170
 and life span, 165, 170, 171
 Pliocene, 165
 rhizomes, 171
 section *Bromus*, 170
 seed-cleaning techniques, 171
- Teliospores**, 197
- Temperate grasslands in NA**, 101
- Temperature regime**
 colder soil, 33
 cool to cold, 44
 soil, 17, 18, 43
- Temporal multipliers**, 390
- Temporal transition multipliers**, 387
- The evolution of weeds**, 103
- Thresholds**, 430
 ecological (*see* Ecological thresholds)
- Tilletia bromi***, 200–203
 chestnut bunt pathogen, 200
 distribution and epidemiology, 203
 host ranges, 203
 life cycle
 bullae/bunts, 202
 cyclic pattern, 202
 dikaryotic secondary basidiospores, 202
 ecology, 202
 spikelets, 202
 teliospores, 201–202
- Transgenerational plasticity (trans-generational plasticity)**, 141
- Transitions**, 375–376
- Transportation corridors and vehicles**
 corridors, 323
 dispersal agents, seeds, 323
 herbicide spraying, noxiousweeds, 323
 invasive species dispersal, 322
 management, 323
 routes, 323
- Trophic levels**, 82
 Bird and Large-Mammal Communities, 86
 Invertebrates and Small Vertebrates, 83–85
- Trust**
 collaborative group processes, 425
 federal agencies, 415
 in government, 424
 land managers' ability, 416
 management ability of public land agencies, 416
- Trust in agencies and organizations**, 415
- U**
- Uncertainty**
 dynamic optimization, 437–438
 and risk, 432
- Ungulates**, 86
- Urban, suburban and exurban development**
 atmospheric gases, 322
 atmospheric nitrogen deposition, 322
 detection and control, invasive, 322
 population spread, 322
- US Department of Interior (USDI) agency**, 418
- US Department of Interior (DOI) Secretarial Order**, 3
- US Forest Service**, 316
- US Geological Survey**, 418
- Ustilago bullata***, 195–198
 distribution, epidemiology and genetics, 199, 200
 head smut pathogen, 196
 host ranges, 198, 199
 life cycle
 inoculum levels, 197
 lose dormancy, 197
 mycelium, 198
 seed germination and seedling growth, 197
 sporidial cells, 197
 sterilizing/castrating fungi, 198
 teliospores, 196, 197
- V**
- Vegetation dynamics development tool (VDDT)**, 377–378
- Vegetation management tools**, 415
- Vernalization**, 142, 143
- Vertebrate**, 62, 83–85
- Volunteering**, 415–417, 420, 421, 424, 425
- W**
- Warm deserts**, 19–21
 ecosystem threats, 21–23
 environmental conditions, 17
 geographic distribution, 18
 hyperthermic and xeric regions, 18, 19
 invasion potential
 blackbrush, 20
 characterization, 19
 creosote bush scrub, 19
 desert grasslands, 20
 livestock grazing, 21
 Mojave desert, 20
 nitrogen deposition, 21
 perennial grasses, 20

- saltbush scrub, 20
 - vegetation type, 20
- management strategies, 23–24
- restoration guidelines, 24
- soil moisture regimes, 17, 18
- soil temperature regimes, 17, 18
- ustic aridic, 17
- wind and solar energy developments, 17
- Water balance
 - and soil hydrology, 71–73
- Water status, 73
- Weed policies and exotic annual *Bromus*
 - ecoregions, 316
 - noxious, 316
 - regulatory programs, 316
- Western forests, 44, 45
 - chaparral and grasslands, 43
 - coolest and wettest conditions, 43
 - ecosystem threats, 45–46
 - invasion potential, 44–45
 - cool to cold temperature regimes, 44
 - Drier and warmer mixed conifer, 44
 - fire, 44
 - fire events, 44, 45
 - high and low severity human
 - disturbances, 45
 - Jeffrey pine, 44
 - lower temperature limits, 45
 - ponderosa pine forests, 44
 - removal of litter and coarse woody
 - debris, 45
 - resistance, 44
 - rising CO₂, 45
 - invasive plants, 43
 - management strategies, 47
 - precipitation, 43
 - ranges, 43
 - resilience, 44
 - resistance, 44
 - shrublands/grasslands, 43
 - soil temperature, 43
- Western great plains, 37–43
 - Bromus gracilis*, 38
 - cool-season species, 38
 - ecosystem threats, 42
 - invasion potential
 - cool-season grasses, 38
 - drought, 39
 - fire, 40–41
 - grazing of ungulates, 40
 - resistance, 38, 39
 - southern Northern Great Plains
 - Network prairie parks, 38, 41
 - management strategies, 43
 - precipitation, 36
 - shortgrass steppe, 36, 38, 42
 - shortgrass steppe and northern and
 - southern mixed-grass prairie, 36
 - temperatures, 36
- Western USA, 2
 - exotic annual *Bromus* (*see* Exotic annual *Bromus* grasses)
- wildfire
 - catastrophic, 432
 - and drought, 442
 - and effectiveness of rangeland
 - rehabilitation treatments, 437
 - frequency, 431
 - frequency and severity, 439
 - homeowners' wildfire risk, 448
 - ignitions, 437
 - post-wildfire rehabilitation, 430
 - risk mitigation, 453
 - suppression cost savings, 442, 445, 452
- Wind and solar energy developments, 17
- Wind Cave National Park in South Dakota, 38
- Wind erosion, 65, 69, 70
- Winter and spring temperatures, 266
- Winter annuals, 180
- Winterkill, 195
- Winter/spring grazing, 30
- Woodland
 - moister, 25
 - oak, 25, 28
 - piñon–juniper woodland, 20
- Wright's Coefficient of Inbreeding, 104, 105
- Wyoming Basin soils, 82
- Wyoming big sagebrush, 388