

Index

A

- Abdominal aortic aneurysm (AAA)
 - balloon occlusion, 88
 - bowel ischemia, 89
 - centres and surgeon specialization, treatment in, 84–86
 - Cochrane review, 71
 - EVAR
 - percutaneous with femoral artery access, 87–88
 - in young patients, 88–89
 - guidelines, 69–73
 - intact AAA
 - open and endovascular repair, 76–78
 - registry data, 78–80
 - management of, 72–73
 - midline laparotomy, 89
 - risk stratification, 86–87
 - ruptured AAA
 - EVAR vs. OR, 80–82
 - registry data, 82–84
 - rupture rates, 90
 - screening
 - guidelines, 71–72
 - systematic reviews/meta analyses, 74–75
 - ultrasound screening, 75
 - surgical monitoring and indication, 69–70
- Access induced ischemia, 227–228
- ACCF. *See* American College of Cardiology Foundation (ACCF)
- Achilles tendon lengthening (ATL), 250
- Acute limb ischemia (ALI)
 - cancer patients with, 193–194
 - classification and prognosis, 181–182
 - controlled reperfusion, 194
 - endovascular and surgical revascularisation, 191–192
 - guidelines, 182–184
 - management algorithm, 195
 - paediatrics, 193
 - percutaneous endovascular thromboaspiration, 187–188
 - registry data, 189–191
 - surgical treatment
 - acute thrombectomy, 188
 - bypass surgery, 188–189
 - systematic reviews, 184–185
 - thrombolysis
 - catheter-directed, 185–186
 - dosage, 187
 - ultrasound-accelerated, 186–187
- Acute mesenteric ischemia (AMI), 132
- Acute thrombectomy, 188
- AHA. *See* American Heart Association (AHA)
- ALI. *See* Acute limb ischemia (ALI)
- American College of Cardiology Foundation (ACCF)
 - abdominal aortic aneurysm, 70, 72
 - acute limb ischemia, 182
 - chronic intestinal ischemia, 125–126
 - critical limb ischemia, 160–161
 - DTAA and TAAA, 47–49
 - extracranial carotid stenosis, 2–3
 - intermittent claudication, 135–138
 - renal artery stenosis, 97–99
 - visceral artery aneurysms, 113
- American Heart Association (AHA)
 - abdominal aortic aneurysm, 70, 72
 - acute limb ischemia, 182
 - aortic dissection, 29
 - chronic intestinal ischemia, 125–126

- American Heart Association (AHA) (*cont.*)
 critical limb ischemia, 160–161
 DTAA and TAAA, 47–49
 extracranial carotid stenosis, 2–4
 fibromuscular dysplasia, 100–101
 intermittent claudication, 135–138
 renal artery stenosis, 97–99
 visceral artery aneurysms, 113
- American Stroke Association, extracranial carotid stenosis, 4
- American Venous Forum (AVF), 256–257
- Aneurysmal degeneration, 32–33
- Angioplasty and Stenting for Renal Artery Lesions (ASTRAL), 103–104
- Antimicrobial therapy, for diabetic foot, 238
- Antiplatelet therapy, intermittent claudication, 154
- Aortic dissection, Stanford's type B
 acute complicated
 aortic fenestration, 37–38
 clinical studies and case series, 36–37
 endovascular and open repair, 35–36
 acute uncomplicated
 aneurysmal degeneration, 32–33
 conservative treatment failure, 33
 endovascular aortic repair, 34–35
 long-term survival, 34
 chronic
 case series, 40–41
 randomized study, 39–40
 registry data, 42–43
 systematic reviews, 39
 guidelines, 29–32
- Aortic fenestration, 37–38
- Aortoiliac occlusive disease (AIOD), in intermittent claudication, 140
- Arteriovenous access stenosis, 216
- Arteriovenous fistula (AVF)
 treatment guidelines, 211, 213–214
 vascular access, 214–215
- Arteriovenous graft (AVG), 211
- Artherectomy, 164
- B**
- Balloon angioplasty
 arteriovenous fistulas, 226
 for intermittent claudication, 148, 150
 renal artery stenosis, 98, 103
- Balloon occlusion, for abdominal aortic aneurysm, 88
- Biological prostheses, 222–223
- Bowel ischemia, abdominal aortic aneurysm with, 89
- Brachiobasilic AVF (BBAVF), 218
- Brachiocephalic AVF (BCAVF), 218
- C**
- Canadian Hypertension Education Program, 102
- Cardiac disturbance syndromes, 102
- Cardiovascular Outcomes in Renal Atherosclerotic Lesions (CORAL), 104–106
- Carotid artery stenosis (CAS)
 CEA *vs.*
 meta-analysis/systematic review, 9–10
 randomized studies, 6–9
 guidelines, 1–6
 registry data, 14–16
- Carotid endarterectomy (CEA)
 after IVT, 21–22
 anesthesia, 22
 and coronary artery bypass, 22
 guidelines, 1–6
 intraoperative shunting, 18
 patching, 18
 registry data, 13–14
 restenosis after CEA, 17
 volume outcome relationship, 16–17
vs. CAS
 meta-analysis/systematic review, 9–10
 randomized studies, 6–9
- Carotid Revascularization Endarterectomy *vs.* Stenting Trial (CREST), 8
- Catheter-based treatments, in CLI, 162
- Catheter-directed thrombolysis (CDT), 185–186
- CEA. *See* Carotid endarterectomy (CEA)
- Celiac artery, endovascular revascularization, 129–130
- Cerebrospinal fluid (CSF) drainage, 50–51
- Chest wall arteriovenous grafts (CWAVGs), 220
- Chronic mesenteric/intestinal ischemia
 endovascular revascularization, 129–130
vs. open revascularization, 130–131
 endovascular therapy, 128–129
 guidelines, 125–126
 open revascularization, 131–132
 registry data, 127–128
 systematic review, 126–127
- Claudication onset time (COT), 143–144
- Coil embolization, splenic artery aneurysms, 114
- Compression stockings (CS), 256, 270, 274
- Congestive heart failure (CHF), 106

- Critical limb ischemia (CLI)
 catheter-based treatments, 162–163
 classification, 159
 endovascular treatments
 arterectomy, 164
 drug-eluting balloons, 163–164
 drug-eluting stents, 163
 guidelines, 161
 registry data, 165–167
 endovascular vs. surgical revascularisation,
 174–177
 guidelines, 160–161
 infrainguinal endovascular treatment,
 164–166
 medical and pharmacological treatment,
 160–161
 peripheral endovascular interventions, 166
 prognosis, 159
 surgical intervention
 biological bypass material, 171–172
 bypass surgery, 170–171
 registry data, 167–170
 revascularisation, 174
 synthetic bypass grafts, 173–174
- Cyanoacrylate-embolization, for varicose
 veins, 273–274
- D**
- DeBakey classification, of aortic dissection, 38
- Descending thoracic aortic aneurysms
 (DTAA)
 cost analysis, 55
 degenerative, 51–52, 63
 endovascular *versus* open surgical
 approach, 49–50
 guidelines, 47–51
 meta-analyses, 51–52
 multicenter study, 55–56
 nonruptured, 55
 registry data, 52–55
 ruptured, 52
 spinal cord protection, 50–51
 surgical indications, 47–49
 volume-outcome relationship, 53–54
- Diabetic foot
 amputation risk, 243
 antimicrobial therapy, 238
 diabetic foot infection, 247–248
 guidelines, 233–238
 local therapy
 dressing products, 244–245
 hyperbaric oxygen therapy, 246
 negative pressure wound therapy, 245
 off-loading methods, 247
 platelet-rich plasma, 246
 skin substitutes, 245
 wound bed preparation, 244
 microbiological considerations, 237–238
 nerve decompression, 248–249
 prognosis, 241–242
 revascularization, 240–241
 severity assessment, 237
 surgical treatment, 238
 tendon lengthening and fascia release,
 249–250
 WIfI-classification-system, 239–240
- Diabetic foot osteomyelitis (DFO)
 diagnosis of, 234–235
 treatment guidelines, 237
- Diabetic foot ulceration (DFU)
 peripheral arterial disease and, 236
 prevention of, 233–234
 wound bed preparation, 244
 wound care for, 235–236
- Distal aortic dissection type Stanford B. *See*
 Aortic dissection Stanford's type B
- Drug-coated balloons (DCBs), 146
- Drug-eluting balloon, 163–164
- Drug-eluting stent (DES)
 critical limb ischemia, 163
 intermittent claudication, 146, 149
- DTAA. *See* Descending thoracic aortic
 aneurysms (DTAA)
- Dual antiplatelet therapy (DAPT), 154
- Duplex scanning, 256
- E**
- Early cannulation grafts, 223–224
- Endografts, 225–226
- Endothermal ablation (ETA), 270
- Endovascular aortic repair
 acute complicated aortic dissection, 35–36
 acute uncomplicated aortic dissection,
 34–35
- Endovascular balloon occlusion (EBO), 88
- Endovascular repair (EVAR)
 abdominal aortic aneurysm
 EVAR and open repair, 76–77
 percutaneous with femoral artery
 access, 87–88
 in young patients, 88–89
 critical limb ischemia
 arterectomy, 164
 drug-eluting balloons, 163–164
 drug-eluting stents, 163
 guidelines, 161
 registry data, 165–167

- Endovascular repair (EVAR)(*cont.*)
- intermittent claudication
 - indications/guidelines, 135–136
 - meta-analyses/systematic reviews, 145–148
 - and surgical intervention, 153–154
 - popliteal artery aneurysm, 203–208
 - renal artery stenosis
 - randomized studies, 103–106
 - systematic reviews/meta-analyses, 102–103
 - uncontrolled studies, 107–108
 - ruptured AAA, 80–82
 - visceral artery aneurysms, 113–115
 - vs. open repair
 - renal artery stenosis, 108–109
 - visceral artery aneurysms, 116–118
 - vs. surgical revascularisation
 - acute limb ischemia, 191–192
 - critical limb ischemia, 174–177
- Endovenous/endovascular thermal ablation (EVTA), 271
- clinical effectiveness and cost-effectiveness, 269–271
 - EVLV vs. open surgery, 264–266
 - meta-analyses and systematic reviews, 262–263
 - radiofrequency ablation, 268–269
 - SVF and AVF guidelines, 257
 - UGFS and conventional surgery, 266–268
 - venous leg ulcers, 263–264
 - vs. open surgery, 264–266
- European Association for Cardio-Thoracic Surgery (EACTS), 47–48
- European Society for Vascular Surgery (ESVS)
- abdominal aortic aneurysm, 69, 73
 - varicose veins, 257–259
- European Society of Cardiology (ESC)
- abdominal aortic aneurysm, 71, 73
 - acute limb ischemia, 183
 - aortic dissection, 30
 - carotid artery disease management, 1–2
 - chronic mesenteric ischemia, 125–126
 - descending aortic aneurysms, 48
 - intermittent claudication, 139
 - renal artery stenosis, 99–100
- Exercise training, intermittent claudication
- COT and PWT, 143–144
 - and endovascular therapy, 150–152
 - indications/guidelines, 135
 - meta-analysis/systematic reviews, 142–143
 - quality of life, 144
 - supervised exercise therapy, 144–145
- Extracranial carotid stenosis
- anesthesia, 22
 - asymptomatic, 23
 - CEA after IVT, 21–22
 - CEA vs. CAS
 - meta-analysis/systematic review, 9–10
 - randomized studies, 6–9
 - clinical practices, 23
 - coronary artery bypass and CEA, 22
 - early risk of stroke, 19
 - eversion/conventional technique, 18–19
 - guidelines, 1–6
 - intervention, early, 19–21
 - intraoperative shunting, 18
 - patching, 18
 - registry data, 10–16
 - restenosis after CEA, 17
 - screening, 5
 - systematic review, 5–6
 - volume outcome relationship, 16–17
- F**
- Femoral artery access, 87–88
- Femoral vein, 219–220
- Femoropopliteal occlusive disease, 140, 152
- Femoropopliteal restenosis, 149–150
- Fibromuscular dysplasia (FMD)
- endovascular vs. open surgery, 108
 - revascularization guidelines, 100–102
- G**
- Gastrocnemius recession (GR), 250
- German Task Force Clinical Nephrology, 213–214
- H**
- Hemodialysis reliable outflow (HeRO) graft, 224–225
- Hemodialysis, vascular access for. *See* Vascular access, for hemodialysis
- Heparin-bonded polytetrafluoroethylene (hepPTFE) grafts, 222
- Hyperbaric oxygen therapy (HBOT), 246
- Hypertension
- with aortic dissection, 31
 - fibromuscular dysplasia, 100
 - renal artery stenosis, 97, 98
- I**
- Infringuinal disease, 161, 164–165
- Intact AAA
- open and endovascular repair, 76–78
 - registry data, 78–80

- Intermittent claudication (IC)
 antiplatelet therapy, 154
 aortoiliac occlusive disease with, 140
 balloon angioplasty, 148
 drug-eluting stent, 149
 endovascular therapy
 indications/guidelines, 135–136
 meta-analyses/systematic reviews,
 145–148
 and surgical intervention, 153–154
 exercise training
 COT and PWT, 143–144
 and endovascular therapy, 150–152
 indications/guidelines, 135
 meta-analysis/systematic reviews,
 142–143
 quality of life, 144
 supervised exercise therapy,
 144–145
 femoropopliteal occlusive
 disease in, 140
 paclitaxel-coated balloon, 149–150
 reporting standards, 141
 revascularization, guidelines for, 139
 surgical guidelines, 136–138
- International Carotid Stenting Study
 (ICSS), 7–8
- International Registry of Acute Aortic
 Dissection (IRAD), 42
- International Working Group on the Diabetic
 Foot (IWGDF), 236–238
- Intraoperative shunting, during CEA, 18
- Ischemia. *See also* Acute limb ischemia
 (ALI); Chronic mesenteric/
 intestinal ischemia; Critical limb
 ischemia (CLI)
 access induced, 227–228
 bowel ischemia, 89
- Ischemic stroke
 AHA/ASA recommendations for, 4
 CEA after intravenous thrombolysis,
 21–22
- L**
- Laparoscopy, splenic artery aneurysms, 121
- Laparotomy, abdominal aortic aneurysm, 89
- M**
- Malperfusion, 31
- Mechano-chemical ablation, 271–273
- Mono antiplatelet therapy (MAPT), 154
- Motor evoked potentials (MEP), 51
- N**
- National Institute for Health and Care
 Excellence (NICE)
 intermittent claudication, 138–139
 for varicose veins, 255–256
- National Kidney Foundation, USA,
 212–213
- Negative pressure wound therapy (NPWT),
 245
- Nerve decompression, 248–249
- Nordic pole walking (NPW), 144
- O**
- Off-loading DFUs, 234
- Open surgical repair (OR)
 abdominal aortic aneurysm, 76–78
 acute complicated aortic dissection,
 35–36
 popliteal artery aneurysm, 202–208
 renal artery stenosis, 108–109
 thoracoabdominal aortic aneurysms,
 60–62
 visceral artery aneurysms, 116
 vs. endovascular repair
 renal artery stenosis, 108–109
 Ruptured AAA, 80–82
 visceral artery aneurysms, 116–118
- Optimal medical care (OMC), 151
- Osteomyelitis, diabetic foot. *See* Diabetic foot
 osteomyelitis (DFO)
- P**
- Paclitaxel-coated balloon, 147, 149–150
- Parenchymal loss, 108
- Patch angioplasty, during CEA, 18
- Peak walking time (PWT), 143–144
- Percutaneous access, 87–88
- Percutaneous angioplasty (PTA)
 acute limb ischemia, 187
 chronic mesenteric ischemia, 131
 critical limb ischemia, 164
 fibromuscular dysplasia, 101, 108
 intermittent claudication, 146, 152
 renal artery stenosis, 101
- Percutaneous endovascular thrombosuction
 (PET), 187–188
- Percutaneous transluminal angioplasty (PTA),
 213
- Peripheral arterial disease (PAD)
 and DFU, 236
 intermittent claudication, 147
- Platelet-rich plasma, 246

- Plethysmography, 256
- Popliteal artery aneurysm (PAA)
 - aneurysmal ligation and bypass grafting, 202
 - endovascular repair, 203–208
 - guidelines, 199
 - medial approach, 203
 - meta-analysis and systematic overviews, 200
 - open repair, 202–208
 - registry data, 200–202
 - thrombolysis, 207
 - women, intervention in, 208
- Prosthetic arteriovenous access, 221
- Pseudoaneurysms, 213, 225–226
- R**
- Radiocephalic AVF (RCAVF), 218
- Radiofrequency ablation (RFA), 268–269
- Recanalization, 114, 267
- Removable cast walker (RCW), 247
- Renal artery aneurysms
 - incidence and outcome, 118–120
 - natural history, 120–121
- Renal artery stenosis (RAS)
 - endovascular therapy
 - randomized studies, 103–106
 - systematic reviews/meta-analyses, 102–103
 - uncontrolled studies, 107–108
 - endovascular vs. open surgery
 - fibromuscular dysplasia, 108
 - registry data, 108–109
 - guidelines, 97–102
- Renal parenchyma protection, 108
- Ruptured AAA
 - EVAR vs. OR, 80–82
 - registry data, 82–84
- S**
- Sclerotherapy, 259–262
 - adverse events, 261
 - contraindications, 259–260
 - indications, 259
 - systematic review, 260
- Shunt-induced ischemia, 213
- Society for Cardiovascular Angiography and Interventions (SCAI)
 - for infrapopliteal arterial intervention, 164–165
 - renal artery stenosis, 102
- Society for Vascular Surgery (SVS)
 - diabetic foot, 233–236
 - varicose veins, 256–257
 - vascular access, 212
- Somatosensory evoked potentials (SSEP), 51
- Splenic artery aneurysms
 - endovascular repair, 114, 117
 - laparoscopic treatment, 121
 - open repair, 117
 - rupture and pregnancy, 121
- Stanford classification, of aortic dissection.
 - See Aortic dissection Stanford's type B
- Steal syndrome, 227–228
- Steam ablation, 271
- Stent revascularization (ST), 151
- Superior mesenteric artery (SMA)
 - endovascular revascularization, 129–130
 - visceral artery aneurysms, 116
- Supervised exercise therapy (SET), 142–145, 151, 152
- Synthetic bypass grafts, 173–174
- Synthetic vascular accesses (SVA), 219
- T**
- Thoracic aortic dissection, AHA
 - guidelines, 29
- Thoracoabdominal aortic aneurysms (TAAA)
 - endovascular repair, 57–58
 - endovascular vs. open surgical approach, 49–50, 56–57
 - guidelines, 47–51
 - hybrid technique, 58–60
 - open repair, 60–62
 - rupture and growth rates, 63
 - spinal cord protection, 50–51
 - surgical indications, 47–49
- Thrombectomy, acute, 188
- Thrombolysis
 - acute limb ischemia
 - catheter-directed, 185–186
 - dosage, 187
 - ultrasound-accelerated, 186–187
 - popliteal artery aneurysm, 207
- Thrombosed dialysis shunts, 215
- Thrombosed popliteal artery aneurysms (PAA), 207
- Total contact casting (TCC), 247
- Tunneled dialysis catheters (TDCs), 225

U

- UK Renal Association, vascular access, 211
- Ultrasound-accelerated thrombolysis (UAT), 186–187
- Ultrasound-guided foam sclerotherapy (UGFS), 270
- Unsupervised exercise therapy, 142–143
- US Preventive Services Task Force
 - abdominal aortic aneurysm, 74
 - screening for asymptomatic CAS, 5

V

- Varicose veins
 - compression stockings as initial treatment, 274
 - cyanoacrylate-embolization, 273–274
 - endovascular thermal ablation, 271
 - endovenous thermal ablation
 - clinical effectiveness and cost-effectiveness, 269–271
 - EVLT vs. open surgery, 264–266
 - meta-analyses and systematic reviews, 262–263
 - radiofrequency ablation, 268–269
 - UGFS and conventional surgery, 266–268
 - venous leg ulcers, 263–264
 - guideline recommendations, 255–260
 - mechano-chemical ablation, 271–273
 - sclerotherapy, 260–262
 - steam ablation, 271
- Vascular access, for hemodialysis
 - access induced ischemia, 227–228
 - arteriovenous access stenosis, 216
 - biological prostheses, 222–223
 - chest wall arteriovenous grafts, 220

- choice of, 214–215, 218–219
 - early cannulation grafts, 223–224
 - endografts, 225–226
 - femoral vein, 219–220
 - guidelines, 211–214
 - hemodialysis reliable outflow graft, 224–225
 - heparin-bonded polytetrafluoroethylene grafts, 222
 - jugular catheter, 220
 - in paediatrics, 220–221
 - percutaneous interventions, 226–227
 - prosthetic arteriovenous access, 221
 - registry data, 216–218
 - thrombosed dialysis shunts, 215
- Venous leg ulcers, 263–264
- Visceral artery aneurysms
 - endovascular and open repair, 116–118
 - endovascular repair, 113–115
 - guidelines, 113
 - incidence and outcome, 118–120
 - open repair, 116
 - symptomatic and asymptomatic, 115

W

- Wound, Ischemia, and foot Infection (WIFI)
 - classification-system, 239–240

Y

- Yale Center for Thoracic Aortic Disease, 48–49

Z

- Zenith Alpha Thoracic Endovascular Graft, 55