

Index

A

- Abiotic stress, 345, 346
Abomasum, 367
Acacia polyphylla, 46
Acetyl esterase, 377
Acidobacteria, 7
Acinetobacter, 77, 79
Acremonium, 43, 51
Acremonium vitellinum, 85
Actinobacteria, 6–8
Actinomycetes, 61
Advanced Genetics Science (AGS), 408
Aedes aegypti, 77
Aeruginosin, 156, 157
Agrobacterium tumefaciens, 409, 413
Alcanivorax, 115
Alcoholic fermentation, 392
Aliterella atlantica, 108
Allorhizobium, 276, 278, 280
Alternaria, 43
Amazon. *See* Brazilian Amazon
Amazon rainforest, 58
Amazonian Dark Earth (ADE), 84, 85
Amazonian fungi, 87–92
1-aminocyclopropane-1-carboxylate (ACC)
 deaminase, 211, 219, 232
Amplicon sequencing, 217
Amycolatopsis orientalis, 65, 66
Anabaena cylindrica, 157
Anabaena variabilis, 157
Anopheles, 79
Antarctic Bottom Water (ABW), 107
Arabidopsis thaliana, 347
Arbuscular mycorrhizae (AM)
 AMF spores, 246
 in cerrado soils, 257–258
 environmental factors, 246
 forest structuration and reforestation,
 258–261
 heavy metals and reclamation, 255–257
 mycorrhizosphere, 249–251
 parasitism, 254–255
 rock phosphate, 254
 taxonomy and functions, 247–248
 tropical soils, phosphate and mycorrhizae,
 251–254
 vascular plants, 246
Arbuscular mycorrhizal fungi (AMF),
 246–262, 324, 325, 332
Archaea, 374
Arthrospira fusiformis, 157
Arthrospira platensis, 154
Ascomycetes, 349
Aspergillus, 43, 75, 78, 82, 90, 91, 93, 94
Aspergillus flavus, 296, 309
Aspergillus niger, 51
Atlantic forest native plants, 152
Auxin production, 219
Average nucleotide identity (ANI), 271
Avicennia schaueriana, 152
Avicennia, 38, 40
 5-azacytidine, 171–176, 180, 181
Azadirachta indica, 323
Azorhizobium, 276, 279
Azospirillum, 273–275, 282, 283
Azospirillum brasilense, 273, 282–284, 306

B

- Bacillus*, 41, 45–48, 50, 110, 112, 113, 116
Bacillus aerophilus, 112
Bacillus agaradhaerens, 114

- Bacillus anthracis*, 83
Bacillus licheniformis, 114
Bacillus pumilus, 112
Bacillus stratosphericus, 112
Bacillus subtilis, 93, 309
Bacillus thuringiensis (Bt)
 toxin, 407, 412, 420
 Bacteriocins, 211
 Basidiomycetes, 349
Bauhinia guianensis, 93
Beijerinckia fluminensis, 327
Betaproteobacteria, 212, 213, 226, 228
 Bioactivity, 156
 Biological nitrogen fixation (BNF), 218, 219,
 271–272, 278–281
 diazotrophic bacteria (*see* Diazotrophic
 bacteria)
 diazotrophic diversity, 270
 N₂-fixing (diazotrophic) bacteria, 270
 nitrogenase constraints, 273–275
 rhizobial taxonomy (*see* Rhizobial
 taxonomy)
 stoichiometry, 270
 symbiotic diazotrophic rhizobia, 270
 Biological soil crusts (BSC), 149
 Biomass degrading enzymes, 379
 Bioproducts, 5
 Bioprospection, 61
 Bioremediation, 114, 115
 Biotic stress, 345
 Biotroph, 247
Bipolaris setariae, 190
Bradyrhizobium, 271, 275–277, 279
Bradyrhizobium diazoefficiens, 306
 Brassicaceae, 353
 Brazilian Agricultural Research
 Corporation, 295
 Brazilian Amazon, 72, 73, 77–81
 biodiversity, 74–81
 in aquatic environments, 77–79
 microbial diversity in soils, 80–81
 biological control, 81
 endophytic fungi, 82
 microbial chemistry, 85–94
 soil remediation, 84–85
 water bioremediation, 82–84
 Brazilian Atlantic forest, 281
 Brazilian biomes, 150, 156, 157
 Brazilian cacti species, 25
 Brazilian cyanobacteria, 150, 151
 Brazilian environments, 144–147, 154
 Brazilian savannah, 58, 59, 61, 62, 66
 Brazilian semiarid region, 5, 11
 Brazilian sugar-energy sector, 316
Brevibacterium, 110, 116
Bruguiera gymnorrhiza, 40
Burkholderia cepacia, 328
- ## C
- Caatinga, 5, 8–10
 Caatinga ecosystem, 26
 Caatinga semiarid region, 5
 Cactaceae family, 22
Camarosporium, 43
Camptotheca acuminata, 352
Candida albicans, 62, 65
 C1A *Orpinomyces* sp., 377
 Carbohydrate-active enzymes (CAZymes), 355
Cassia leptophylla, 64
 CAZy database, 355
 Cellodextrinase, 379
 Cellulosomes, 377
Centaurea stoebe, 345
Cephalosporium, 82
 Chemical characterization
 endophytic biochemical, 197–202
Chenopodiaceae, 43
 Chromatographic column (CC), 197–199
Chromobacterium violaceum, 78, 79
 Chroococcalean, 151
Chryseobacterium piscium, 111
Cladosporium, 43
Clavibacter xyli, 409, 412, 413
Clavicipitaceae, 212
 Clustered Regularly Interspaced Short
 Palindromic Repeats (CRISPR), 416
 Co-culture, 191–193
 Coevolution, 370
 Cold-stress tolerance, 346
Colletotrichum, 43, 75, 81, 82, 294–296, 299,
 300, 303, 304, 307–309
Colletotrichum acutatum, 307
Colletotrichum falcatum, 323, 325
Colletotrichum gloeosporioides, 294–296,
 298, 299, 307–309
 in actinomycetes, 181
 antibacterial activity, 175, 181
 5-azacytidine and sodium butyrate, 176
 bioactive compounds, 179
 biofabrication, 181
 cryptic compounds, 176
 disc diffusion assay, 175
 DNA methylation, 180
 epigenetic modifiers, 171–173, 180, 182
 HPLC analysis, 175–179

- human bacterial pathogens, 179
 - ITS rDNA sequence-based phylogenetic tree, 174
 - leaf segment of *S. cumini*, 179
 - metabolites, 173–176
 - microbes/endophytes, 182
 - secondary metabolites, 174, 180
 - Colletotrichum lindemuthianum*, 307
 - Colletotrichum tofieldiae*, 353
 - Colony forming units (CFUs), 216
 - Competent endophyte, 211
 - Compost, 257
 - Cordyceps confragosa*, 85
 - Cry1Ab gene, 420
 - Convertible bacteria, 108–119
 - diversity and biotechnological potential
 - bacteria isolated from other sources, 111
 - bioactive molecules, 116–117
 - bioremediation, 114–115
 - enzymes, 111–119
 - genomic studies, 117–119
 - seawater and sediments, 108–111
 - Cupriavidus*, 275, 281
 - Curtobacterium*, 41, 48
 - Curvibacter*, 79
 - Curvularia*, 81
 - Curvularia pallescens*, 51
 - Curvularia* thermal tolerance virus (CThTV), 346
 - Cyanobacterial morphotypes, 141
 - Cyanobacterial systematics, 142
 - Cyanopeptides, 156
 - Cyanopeptolin, 156, 157
 - Cyanotoxin, 156
 - Cylindrospermopsis raciborskii*, 151
 - Cytophaga*, 114
- D**
- Deinococcus-thermus*, 115
 - Desorption electrospray ionization (DESI), 200
 - Diaporthe*, 43, 50, 51
 - Diaporthe phaseolorum*, 51
 - Diazotrophic bacteria
 - Acetobacter diazotrophicus*, 283, 284
 - associative and endophytic diazotrophic bacteria, 282
 - Azospirillum*, 282–283
 - classification of, 272–273
 - diversity of, 271–272
 - Herbaspirillum*, 283
 - legume-rhizobia specificity, 276–278
 - phytohormones, 282
 - rhizobia, 275–281
 - rhizobia-legume symbioses, 276
 - sustainable agriculture, 284
 - Diazotrophic endophytes, 210
 - Diazotrophs, 327
 - Dichanthelium lanuginosum*, 346
 - Direct-fed microbials (DFMs), 379
 - Diversity, 22–31, 142
 - in biome, 153
 - Brazilian cyanobacteria, 151
 - cyanobacteria, 142
 - endolithic cyanobacteria, 143
 - taxonomic, 143
 - in Tucuruí hydropower plant, 153
 - DNA methyltransferase (DNMT) inhibitors, 171, 172, 180
 - Double labeling of oligonucleotide probes (DOPE), 216
 - Double-stranded RNA virus genome (dsRNA), 308
 - Drought stress, 7, 9, 10
 - DroughtGard® corn, 406, 407
 - Duguetia stelechantha*, 75, 86, 93
- E**
- Ecosystems, 140, 143, 148–150, 152
 - Ectomycorrhizae (ECM)
 - in Acacia, 262
 - in Brazil, 261
 - dyeing and microscopic examination, 261
 - “ecto”, 261
 - Hartig net, 261
 - time-course field experiment, 262
 - Edaphoclimatic factors, 316
 - Electrospray ionization (ESI), 200
 - Endo-glucanase, 379
 - Endoglucanase A (EgIA), 413
 - Endophyte, 247, 249, 252, 254, 258
 - endorrhizosphere, 210
 - Koch’s postulate, 210
 - microbial endophytes, 210
 - nonlegume plants, 210
 - rhizobia with legume plants, 210
 - siderophores and phytohormones, 211
 - Endophyte diversity assessment
 - biocontrol (antifungal activity), 222
 - culture-dependent methods, 214–216
 - culture-independent methods, 216–218
 - macro- and micronutrients, 220–222
 - nitrogen fixation, 218, 219
 - phytohormonal balance, 219, 220
 - plant tissues, 214
 - practical study, 223

- Endophyte fungi
 axenic culture metabolites, 186
 challenges, 186
 chemical characterization, 197–202
 chemical communication, 186
 column chromatograph, 198–199
 extraction, secondary metabolites,
 194–196
 gas chromatograph, 199–200
 HPLC, 199
 LC-MS/MS, 200–201
 metabolic compounds, 185, 188–191
 metabolic diversity, 191–202
 NMR, 202
 plants, 187, 188
- Endophyte isolation, 215
- Endophyte-epiphyllous organism
 interactions, 352
- Endophytes, 21–30
 asexual clavicipitaceous, 212
 bacterial community composition, 213
 bacterial endophytes, 212
 biological domains, 211
Clavicipitaceous, 212
 diazotrophic microbes, 212
 diversity of, 213
 facultative endophytes, 212
 gramineous plants, 212
 prokaryotic sequences, 213
 xenobiotics, 214
- Endophytic bacteria, 273
- Endophytic fungi, 82, 170, 176, 181, 182,
 186–191, 193, 194, 196, 198, 200
 Cactaceae family, 22
 Cacti, 24–26, 30
 dry environments, 22–24
 endophytism, 22
 endophytology, 21
 mycoendophytes, 22
- Endophytic microbiota in situ, 224
- Endophytic microorganisms, 21, 38–49
 agronomic and biotechnological potential,
 43–49
 antimicrobial activity, 49–51
 mangrove endophytic bacteria, 40–42
 enzyme production and bioremediation,
 47–49
 plant growth promotion, 43–47
 mangrove endophytic fungi, 42–43
- Endophytism, 22
- Endophytology, 21
- Endo- β -1,4-glucanase, 379
- 5-enolpyruvylshikimate-3-phosphate (EPSP)
 synthase, 406, 407, 415
- Enterobacter*, 41, 45, 46, 48, 49, 212, 221–224
- Enterococcus*, 221
- Enterococcus faecalis*, 65
- Entodinium* genus, 376
- Entomopathogenic fungi, 345
- Epicoccum* sp.
 biological control agent, 319
E. nigrum, 319, 322
 epicolactone, 322
- Epilithic and endolithic cyanobacteria, 148
- Epiphytic mycelia, 171
- Erythrobacter*, 110, 115, 117
- Erythrobacter atlanticus*, 114
- Escherichia coli*, 49, 62, 63, 67
- Esterases, 377
- Ethanol Production, Brazilian Process, 392
- Exopolysaccharides (EPS), 10, 12
- Exserohilum rostratum*, 93
- Extremophiles, 12
- F**
- Fa cai in mandarin, 154
- Farmyard manure (FYM), 323
- Fast atom bombardment (FAB), 200
- Fatty acid methyl ester (FAME), 216
- Feruloyl esterase, 377
- Firmicutes, 6, 7
- Fluorescence in situ hybridization (FISH), 216
- Fomitopsis pinicola*, 85
- Fungal endophytes
 abiotic stress, 345–346
 antibiosis, 344
 applications, 356, 357
 axenic culture, 344
 bioabatement, 351
 biotic stress, 345
 industrial enzymess, 348–349, 354–356
 morphological and molecular studies, 344
 nutrition management, 346–347
 in planta, 344
 plant fitness, 352–353
 pretreatment and saccharification, 349–351
 secondary metabolites, 347–348, 353–354
- Furaldehydes, 351
- Fusarium*, 43, 50, 75, 81, 82
- Fusarium oxysporum*, 47
- G**
- Gammaproteobacteria, 108, 110–112,
 115–117, 121
- Gas chromatograph (GC), 198–200
- Gastrointestinal microbiome, 378
- Genetically modified organisms (GMOs),
 404–415

- adverse effects, 404
 - definition, 404
 - development of, 404
 - field experiments and controlled environments, 416–423
 - GMA (see GM animals (GMAs))
 - GMMs (see GM microorganisms (GMMs))
 - GMPs (see GM plants (GMPs))
 - host plant and fungal communities interactions, 419–421
 - host plant and microbial communities interactions, 421–423
 - microorganisms, 415–416
 - plant and animal genomes, 404
 - risk assessment, 424–425
 - in tropical countries, 423–424
 - Genomics
 - G. diazotrophicus* PA1 5 and *Serratia* Sp. M24 T3, 229–232
 - Geodermatophilus*, 8
 - Gibberellins, 328
 - Glomerella cinctulata*, 308
 - Glomerella cingulata*, 295, 296
 - Gluconacetobacter*, 211, 212, 218, 220, 222, 223, 225, 226, 230
 - Gluconobacter diazotrophicus*, 284, 327, 328
 - β -glucosidase, 351, 379
 - GM animals (GMAs)
 - caseine, 405
 - GM cattle milk, 405
 - human glucocerebrosidase protein, 405
 - in nature, 404
 - in the United States, 405
 - GM Microorganisms (GMMs), 411–415
 - in agriculture
 - biocontrol, insects and disease, 412–413
 - biofertilizers, 412
 - in bioremediation, 413–414
 - in human disease, 415
 - in probiotics, 414
 - regulatory agencies, 411
 - applications, 409–410
 - in biofuels, 408–411
 - in bioreactors, 408
 - and biosecurity, 408
 - FrostBan®, 408
 - research, 408
 - GM plants (GMPs)
 - and animals, 406
 - and bacterial communities interactions, 417–419
 - Bt toxin acts, 407
 - chemical pesticides, 405
 - DroughtGard® corn, 407
 - EPSP synthase, 407
 - glyphosate herbicide, 407
 - herbicides, 407
 - in maize, 408
 - and microbial communities interactions, 417
 - nutrient improvements, 407
 - transgenic traits, 407
 - in the United States, 405
 - wood, 407
 - GM *Sphingomonas paucimobilis* UT26, 414
 - Gramineous plant, 212
 - Growth promotion, 246, 249, 250
 - Guarana, 294–309
 - bacteria in guarana leaves
 - endophytic bacteria, 300–304
 - simultaneous isolation, 302, 303, 305
 - bacteria isolated from the rhizosphere, 304–307
 - effect on plant growth, 305–307
 - Colletotrichum*, 294–295
 - metabolites of bacteria and fungi from endophytes, 309
 - mycovirus and anthracnose control, 307–308
 - nonpathogenic fungi and bacteria, 295–300
 - endophytic fungi in *P. cupana* leaves, 296–299
 - from guarana seeds, 299–300
 - Guignardia bidwellii*, 51
 - Gut metagenomics, 379
- ## H
- Halomonas*, 110, 117
 - Halomonas zincidurans*, 114
 - Heavy metals, 247, 255, 256, 259
 - Herbaspirillum*, 283
 - Herbaspirillum seropedicae*, 283
 - Hydrogenosomes, 377
 - High-performance liquid chromatography (HPLC), 197–201
 - Histone deacetylase (HDAC), 171, 172, 180
 - Horizontal gene transfer (HGT), 378
 - HPLC analysis
 - 5-azacytidine, 176
 - C. gloeosporioides*, 177–179
 - crude compounds, 175
 - RP-C18 column, 173
 - Human chronic myeloid K562 leukemia, 190
 - Hydralazine, 171
 - Hydrogenosome, 377
 - Hydroxymethyl furfural (HMF), 351
 - Hypoxylon*, 51

I

- In situ hybridization (ISH), 216
- Indole acetic acid (IAA), 38, 282, 307, 328
- Industrial yeast strains, 393–394
 - ethanol production in Brazil, 394
 - genome complexity, 394–395
 - main characteristics, 396–398
 - tailored for ethanol production, 395–396
- Inoculant, 273, 277, 281–284
- Inselbergs, 149

J

- Joostella*, 108

K

- Koeleria macrantha*, 345
- Köppen climate classification, 3
- Krokinobacter*, 108

L

- Laguncularia racemosa*, 38, 152
- Leeuwenhoekella*, 108
- Leishmania infantum/chagasi*, 66
- Length heterogeneity PCR (LH-PCR), 216
- Leptospherulina crassiasca*, 352
- Lignocellulosic biomass (LCB), 348–351, 356
- Lyophilization, 196

M

- Magnaporthe grisea*, 22
- Mameliella atlantica*, 114
- Mangrove endophytic bacteria, 40–42
 - enzyme production and bioremediation, 47–49
 - plant growth promotion, 43–47
- Mangrove endophytic fungi, 42, 43
- Mangrove forests, 38
- Marine-derived fungal
 - (MDF) endophytes, 350
- Marinobacter*, 111, 112, 114–117
- Martella*, 111
- Matrix-assisted laser desorption ionization (MALDI), 200
- Mauritia flexuosa*, 74, 86
- Mesorhizobium*, 276, 278
- Metabolic compounds, 185, 188, 190, 191
- Metagenomic surveys, 7
- Metagenomics, 378, 379
- Metarhizium anisopliae*, 326

- Metatranscriptome, 232
- Metatranscriptomic analysis, 380
- Methanobrevibacter*, 374
- Methanogens, 374, 375
- Methanogens metabolism, 374
- Methanomicrobium*, 374
- Methanoplasmatales, 375
- Methyl coenzyme M reductase (*mcrA*)
 - genes, 374
- Methylobacterium*, 40, 42, 49, 280
- Miconia albicans*, 59, 64–66
- Microbacterium*, 111
- Microbial chemistry, 85, 92–94
- Microbial diversity, 106, 124
- Microbiome, 250, 331, 332
- Microbiota, 299, 304, 309, 331
- Microbispora amethystogenes*, 64
- Micrococcus*, 41, 110
- Microcystin, 151, 156
- Microcystis aeruginosa*, 151
- Microginin, 156
- Micromonospora yulongensis*, 64
- Minimum inhibitory concentration (MIC), 50
- Most probable number
 - (MPN) technique, 216, 223
- Multilocus sequencing analysis (MLSA), 271
- Muricauda*, 108
- Mycelia Sterilia*, 51
- Mycoendophytes, 22
- Mycorrhizae, 246–263, 324, 325
 - AM (*see* Arbuscular mycorrhizae (AM))
 - ECM (*see* Ectomycorrhizae (ECM))
 - host plant, 246
 - orchid mycorrhizae (*see* Orchid mycorrhizae)
 - types of, 246
- Mycorrhizal root, 260
- Mycorrhizal spore, 260
- Mycorrhizosphere, 249, 250
- Myrciaria dubia*, 80

N

- Natural products, 156
- Neocallimastix frontalis*, 376
- Next-generation DNA sequencing (NGS), 271
- Nitrogen use efficiency (NUE), 219
- Nitrogenase, 270, 272–275, 282
- Nocardioides*, 8
- Nocardioides albus*, 64
- Noncultivable microorganisms
 - ecological studies, 120–125
 - methods to access, 119–130

- prospective metagenomics, 125–130
 Nonpathogenic microbes, 315
 Nonruminants, 368
 North Atlantic Deep Water (NADW), 107
Nostoc species, 155
 Nuclear magnetic resonance
 (NMR), 197, 198, 202
- O**
- Ochrobactrum*, 281
 Oligocene, 370
 Omasum, 367, 368
 Orchid mycorrhizae
 anthropogenic intervention, 263
 mycoheterotrophic phase, 263
 orchid mycorrhizal fungi, 263
Rhizoctonia, 263
 Organophosphorus, 380
 Oscillatorialean, 151
 Oscillatoriales, 151
 Osmolytes, 7–11
 Oxyphotobacteria, 140
- P**
- Paclitaxel, 194
Paecilomyces, 43
Paecilomyces variotii, 51
Paenibacillus polymyxa, 64
Paenibacillus terrae, 67
Pantoea, 41, 46, 48
Pantoea agglomerans 33.1, 329, 330
Paraburkholderia, 281
 Parasitism, 254
 Passenger endophyte, 211
Paullinia cupana, 294
 p-coumaryl esterase, 377
 PCR-DGGE technique, 422
Penicillium, 43
Penicillium avelanni, 86
Penicillium janthinellum, 355
Penicillium sclerotiorum, 94
Pestalotiopsis, 43, 51, 81, 82, 91, 93
Pestalotiopsis guepinii, 93
Phaeomoniellales, 352
Phoma, 43
Phomopsis, 43, 50, 51
Phormidium bohneri, 156
 Phosphate, 247, 251, 252, 254–257, 261
 Phosphorous, 220
 Photoassimilates, 317, 318
 Photodiode array detectors (PDAs), 173
Phyllobacterium, 275, 280
Phyllosticta, 43
 Phylogenetic diversity (PD), 6
 Phylum cyanobacteria, 156
 Phylum neocollimastigomycota, 377
 Phytohormone indoleacetic acid, 328
 Phytohormones, 220
Phytophthora palmivora, 81
 Pigmented endophytic fungi, 22
 Pine wilt disease (PWD)
 endophytic microbiomes, 225–226
 microbial community, 228–229
 portuguese nematode-carried bacterial
 community, 226–228
 Pinewood nematode (PWN), 225
Piriformospora indica, 346, 349
 Plant growth promoting bacteria
 (PGPB), 273, 282–284
Azospirillum spp., 329
Beijerinckia sp., 327
 in Brazil and India, 326
Burkholderia sp., 328
 endophytic bacteria, 329
Gluconacetobacter sp., 327–328
 growth-promoting processes, 327
Herbaspirillum spp., 329
 isolation, 327
Pantoea agglomerans
 strain 33.1, 329, 330
 Plant growth-promoting rhizobacteria
 (PGPR), 8, 9
 Plant stress homeostasis-regulating
 rhizobacteria (PSHR), 8
 Plant-endophyte interactions, 187
Polaribacter, 114
Polynucleobacter, 77
 Potassium, 221
 Potato dextrose agar
 (PDA) medium, 172–174
 Preparative thin-layer chromatography
 (PTLC), 197
Prochlorococcus, 108, 121
 Proteobacteria, 6, 7, 278, 281
Proteus morgani, 49
 Protoruminants, 371
Prototheca moriformis, 411
 Protozoa, 376
Pseudoalteromonas, 114
Pseudomonas, 41, 46, 47, 50
Pseudomonas aeruginosa, 49, 93
Psychrobacter, 111
Psychropotes longicauda, 111
Pythium aphanidermatum, 47

R

- Random amplified polymorphic DNA (RAPD), 216
- Recombinant human lactoferrin (rhLF), 405, 406
- Repetitive sequence-based PCR (Rep-PCR), 216
- Reverse transcription PCR (RT-PCR), 216
- Rhizobial taxonomy
 - Allorhizobium*, 280
 - Allorhizobium undicola*, 278
 - Azorhizobium*, 279
 - Bradyrhizobium*, 279
 - Cupriavidus*, 281
 - Mesorhizobium*, 278
 - Methylobacterium*, 280
 - Microvirga*, 280
 - Paraburkholderia*, 281
 - Phyllobacterium*, 280
 - Rhizobium*, 278
 - Sinorhizobium*, 279
 - α -Proteobacteria, 281
- Rhizobium*, 278
- Rhizoctonia solani*, 47
- Rhizophora mangle*, 38
- RNAi technology, 407
- Rumen biotechnology, 378–380
- Rumen microbiome
 - anaerobic fungi, 376–377
 - archaeal community, 374–375
 - bacteria, 372
 - bacteriophages, 378
 - bacteroidetes, 373
 - biotechnological perspectives, 378–380
 - “browsers” and “frugivorous”, 369
 - core bacterial microbiome, 369
 - fibrolytic, 373
 - mechanical mastication and enzymatic reaction, 368
 - Megasphaera elsdenii*, 373
 - metabolic activity, 373
 - and microorganisms, 370–371
 - molecular biology approaches, 380
 - nonruminants, 368
 - phylum Firmicutes, 373
 - protozoa, 375–376
 - rumen bacterial community, 372
 - rumen harbors, 368
 - ruminal microbial communities, 371
 - ruminants, 367
 - Selenomonas ruminantium*, 373
 - 16S rRNA gene, 373
 - Streptococcus bovis*, 373
 - subpopulations, 372
 - VFAs, 368
- Ruminants, 367, 369–372, 375

S

- Saccharification, 348
- Saccharomyces cerevisiae*, 392, 408, 409, 411
- Saccharomyces* strains, 394, 396, 398
- Salicornia brachiata*, 40
- Sclerotinia sclerotiorum*, 47
- Secretomes, 356
- Serratia*, 41, 50, 221, 224, 228–232
- Shewanella sediminis*, 111
- Shigella sonnei*, 65
- Shinella*, 275, 281
- Sinorhizobium*, 276, 279
- Siphophages, 378
- Soil fertility, 259
- Soil remediation, 84
- Soil structure, 253
- Solanum lycocarpum*, 59, 60, 62, 63
- Sonneratia caseolaris*, 50
- South atlantic ocean
 - oceanography, 107
 - sediments, 107
 - water movements, 107
- Soxhlet system, 196
- Spirulina, 154
- Sporormiella*, 43
- Sporormiella minima*, 43
- Staphylococcus*, 62–65, 67
- Staphylococcus aureus*, 50, 62, 63, 65, 67, 93
- State of Amazonas Research Foundation (FAPEAM), 77
- State of São Paulo Research Foundation (FAPESP), 77
- Strategies to endophytic metabolism exploration, 191–202
- Streptococcus pneumoniae*, 50
- Streptomyces*, 61–64
- Streptomyces argenteolus*, 64
- Streptomyces griseus*, 8
- Streptomyces peucetius*, 64
- Streptomycetes, 61
- Suaeda monoica*, 40
- Suberoylanilide hydroxamic acid (SAHA), 171, 181
- Sugarcane, 326–331
 - AM fungi, 332
 - clean and renewable energy, 314
 - Cryptococcus* yeasts, 326
 - cultivation of, 314–316
 - entomopathogenic fungi, 326
 - Epicoccum* sp., 319–322
 - ethanol production, 314
 - fungal and bacterial genera, 315
 - Fusarium* species, 326
 - harvest area and production, 315
 - microbiome, 331–332

- and microorganism interactions, 315
 - mycorrhizal fungi, 324–325
 - nonpathogenic interactions, 319
 - in nonpathogenic relationships, 320–322
 - P. agglomerans* 33.1, 330
 - PGPB (*see* Plant growth-promoting bacteria (PGPB))
 - production factors, 317–318
 - Trichoderma* spp., 322–324
 - World and Brazilian average yield, 317
 - Sugarcane endophytes
 - applications, 224–225
 - culture-dependent and -independent methods, 223–224
 - Sulfitobacter*, 111
 - Sulfitobacter pontiacus*, 114
 - Symbionts, 247, 263
 - Synechococcus*, 121
 - Syzygium cumini*
 - abovementioned properties, 171
 - 5-azacytidine, 181
 - cryptic compounds, 181
 - DNMT and HDAC inhibitors, 180
 - DNMT inhibitors, 171
 - endophyte, 170
 - endophytic *Alternaria* sp., 181
 - epigenetics, 171
 - host origin, 170
 - in India, 170
 - natural source, 170
 - pharmacological properties, 170
- T**
- Tailored-yeast strains, 395–397, 399
 - Talaromyces aculeatus*, 309
 - Tecuitlatl, 153
 - TempSym1 *Curvularia protuberata*, 346
 - Thalassospira*, 111, 115
 - Theobroma cacao*, 346
 - Thermophile, 13
 - Thermoplasmata, 375
 - Thermotolerant, 12–13
 - Thermotolerant cellobiohydrolase, 351
 - Thiamine pyrophosphate (ThPP), 232
 - Tolypocladium*, 81
 - Trametes versicolor*, 349
 - Trehalose, 8
 - Trichocoleus desertorum*, 149
 - Trichoderma*, 43, 51
 - Trichoderma martiale*, 81
 - Trichoderma* spp.
 - biological control, 323
 - biological control agents, 323
 - FYM/bioagents, 323
 - lignocellulosic organic material, 323
 - soil fungal community, 323
 - T. virens*, 324
 - T. viride*, 323
 - Tropical cyanobacteria
 - Bergey's Manual of Systematic Bacteriology*, 142
 - biogeochemical processes, 139
 - biotechnological applications, 153–157
 - botanical and bacteriological frameworks, 142
 - constituted of N₂, 140
 - cyanobacterial systematists, 142
 - on earth, 139
 - epi/endobiotic symbioses, 140
 - morphological characters, 141
 - Precambrian Era, 140
 - in tropical Brazil, 150–153
 - in tropics, 142–150
 - Tropical dry forest, 25
 - Tropical Savannah
 - bioactivity of endophyte species, 61–67
 - endophytic bacteria, 62–67
 - trees, 59–60
- V**
- Vibrio cincinnatiensis*, 116
 - Victoria amazonica*, 74, 86
 - Visceral leishmaniasis (VL), 66
 - Vitellibacter*, 108
- W**
- Water Bioremediation, 82, 83
- X**
- Xerophiles, 11–12
 - Xerotolerants, 11–12
 - Xylaria*, 43, 51, 75, 82, 89, 93
 - Xylocarpus moluccensis*, 50
 - Xylotrophic endophytes, 212
- Z**
- Zinc, 222
 - Zoosporogenesis, 377
 - Zunongwangia*, 108
 - Zunongwangia atlantica*, 114