

# Index

## A

Activities of daily living (ADLs), 38  
Acute brain injury, 80  
Acute concussion, 71  
Acute respiratory distress syndrome (ARDS), 11  
*Aid in cases of illness*, 116  
Akathisia, 45  
Allodynia, 28  
Alternating attention training, 110  
Alzheimer disease (AD), 18, 19, 87, 89  
American Congress of Rehabilitation Medicine (ACRM), 69  
American Professional Football Association (APFA), 75  
Amnesic syndrome, 107  
Amyloid pathology, 102  
Amyotrophic lateral sclerosis (ALS), 89  
Anamnesis, 118  
Anisotropic diffusion, 97  
Annegers' classification, 50  
Anterior nucleus of the thalamus (ATN), 55  
Apolipoprotein Eε4 (APOEε4) allele, 51  
Apparent diffusion coefficient (ADC), 81, 97  
Aquaporins, 18  
Arachnoid granulations, 17  
Arterial spin labeling (ASL), 100  
Atelectasis, 29  
Attention  
    alternating training, 110  
    and cognitive disorders, 108  
    in distracting/stressful situations, 109  
    divided training, 110  
    and memory deficit, 107  
    selective training, 110  
    sustained training, 110  
    and temporal-spatial orientation, 108  
Auditory discrimination, 107  
Axial diffusivities (AD), 98

## B

Ballism, 44–45  
Bed rest, 25–27, 29  
Biomarkers, 80, 81  
Blood brain barrier (BBB), 52  
Blood-oxygen-level-dependent (BOLD) signal, 100  
Bone density, 26–27  
Bone loss, 27, 30  
Boxing x martial arts, 83  
Brain contusions and lacerations, 95  
Brain Injury Association of America (BIAA), 35, 101  
Brain injury brain concussion, 70, 71  
Brain reorganization, 1

Brain-derived neurotrophic factor (BDNF), 3  
Brazilian Public Health System (SUS), 113, 116  
Brownian movement, 97

## C

Cannabinoid type-1 (CB1) receptor, 54–55  
Cardiovascular adaptation, 29  
Catastrophic brain injuries, 79  
Cavum septum pellucidum, 81  
Centers for Disease Control and Prevention (CDC), 75  
Cerebral blood flow (CBF), 100  
Cerebral blood volume (CBV), 100  
Cerebral palsy, 36  
Cerebral perfusion pressure (CPP), 11  
Cerebral vascular accident (CVA), 114, 116  
Chorea, 44–45  
Chronic brain injury, 80  
Chronic traumatic encephalopathy (CTE), 75, 103  
    aggression, 90  
    autopsy, 89  
    concussions, 89–90  
    depression, 90  
    irritability, 90  
    neuropsychiatric symptoms, 89, 91  
    and PCS, 91  
    suicidality, 90, 91  
Closed-head injury, 51  
Cognitive dysfunction, 35  
Cognitive function evaluation, 110  
Cognitive impairment  
    amnesic syndrome, 107  
    frontal lobe injury in post-TBI retain, 107  
    after hospital–post-acute/chronic phase, 108  
    language disorders, 107  
    long-term memory, 107  
    long-term neuropsychiatric deficits, 108  
    mechanisms of acceleration-deceleration, 107  
    subclinical findings, 107  
    type and degree, 107  
    working memory, 107  
Cognitive rehabilitation, 107  
    attention  
        alternating training, 110  
        divided training, 110  
        selective training, 110  
        sustained training, 110  
    memory, 110–111  
    training strategies outside the outpatient unit, 111  
Cognitive rehabilitation outpatient clinic, 117  
Computed tomography (CT), 96

Concussion, 70, 71, 87–88  
 knowledge and awareness, 76  
 National Concussion Surveillance System, 76  
 signs and symptoms, 76  
 sports, 75, 77  
 Consciousness, 21  
 Contracture deformities, 27  
 Controlled cortical impact (CCI) injury, 51, 54  
 Cough peak flow (CPF), 12  
 Craniectomy size, 18  
 Cranioplasty, 20, 22

## D

Daily life activities, 113  
 Data-driven types of analysis techniques, 100  
 Decompressive craniectomy (DC), 18, 20–23  
 Deep brain stimulation (DBS), 55  
 Deep vein thrombosis, 29  
 Deep venous thrombosis (DVT), 37  
 Deformity, 25–28, 30  
 Dementia pugilistica, 79  
 Depression, 90  
 Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV), 88  
 Diffuse axonal injury (DAI), 81, 96–99  
 Diffuse swelling, 95  
 Diffusion tensor imaging (DTI), 53, 97–100  
 Diffusion-weighted imaging (DWI), 97–100  
 Disability, 71  
 Disability-adjusted life year (DALY), 71  
 Disrupted-in-schizophrenia I (DISC-1), 5  
 Divided attention training, 110  
 Dynamic susceptibility contrast imaging (DSC), 100  
 Dysesthesia, 28  
 Dystonia, 42

## E

Early mobilization, 25, 28–30  
 Early rehabilitation, 30  
 Electrical activity  
 brain tissue, 61  
 EEG, 62, 63  
 LORETA, 63  
 parahippocampal, 64  
 Electroencephalography (EEG), 50, 52–54  
 advantages, 62  
 clinical assessment, 62  
 clinical electrophysiologic techniques, 62  
 cognitive disorders, 61  
 cognitive dysfunction, 61  
 contraindication, 62  
 cortical pyramidal neurons, 62  
 CT and MRI data, 62  
 dEEG, 63, 64  
 discovery, 61  
 electrical activity, brain tissue, 61  
 intrinsic properties, 62  
 mTBI, 63  
 neural mechanisms, 61  
 new neurological and psychiatric diagnostic tool, 61  
 oscillatory activity, 62  
 postconcussive symptoms, 65  
 post-traumatic epilepsy, 64  
 and sleep, 64–65  
 in TBI, 62–63

Encephalopathy, 103  
 Endocrine, 29  
 Endogenous neural recovery plasticity, 38  
 Ependymal cilia, 17  
 Ephrins and their receptors (Eph), 3  
 Epilepsy, 22  
 Epileptogenesis, 52–54  
 Eucapnia, 11  
 Exact low-resolution brain electromagnetic tomography (eLORETA), 64  
 Executive functions, 107, 110, 111  
 External stimulation, 35  
 Extra-axial hematomas, 95  
 Extracellular matrix (ECM), 17  
 Extra-pyramidal syndromes  
 dystonia, 42  
 hyperkinesias, 41  
 Parkinson disease, 42, 43  
 parkinsonism, 43, 44  
 tremor, 41

## F

Fatigue, 26  
 FDG-PET, 101, 102  
 Fluid percussion injury (FPI), 51, 52, 55  
 Fluid-attenuated inversion recovery (FLAIR), 96–99  
 Fractional anisotropy (FA), 98, 99, 101  
 Frequency bands, 62  
 Functional MRI (fMRI), 100  
 Functional neuroimage  
 classification, 95  
 CT, 96  
 DAI, 96–99  
 distinct/concurrent pathologic features, 95  
 DTI, 97–100  
 DWI, 97–100  
 fMRI, 100  
 GCS, 95, 97, 98  
 mechanisms, 96  
 and molecular, 101–103  
 MRI, 96  
 multiplanar T1-weighted images, 96  
 perfusion MRI, 100  
 scoring categorization, 95  
 SWI, 97  
 T2\*-GRE, 97

## G

GABAergic pathways, 52  
 Gamma-aminobutyric acid-ergic (GABAergic), 91  
 Glasgow Coma Scale (GCS), 70, 95, 97, 98  
 Glia cell, 18  
 Glial fibrillary acid protein (GFAP), 2  
 Glial glutamate transporter protein (GLT), 52  
 Glutamic acid decarboxylase 1 (GAD1), 51  
 Granular neurons, 5  
 Gross Motor Function Measures (GMFM), 37

## H

Head injury, 63, 64  
 Head trauma, 51, 53, 54, 71, 72  
 Herniations, 95–96  
 Heterotopic ossification (HO), 27  
 Hippocampal atrophy, 81

Hospital management, 108  
 Hydrocephalus, 17–22  
 Hyperexcitability, 52, 55  
 Hyperexcitation, 52  
 Hyperkinesias, 41  
 Hyperkinetic posttraumatic movement disorders  
   akathisia, 45  
   ballism, 44–45  
   chorea, 44–45  
   myoclonus, 44  
   psychogenic movement disorders, 45  
   tics, 44  
 Hypocapnia, 11  
 Hyponatremia, 53

## I

Immobility  
   aerospace science, 25  
   atelectasis, 29  
   bed rest, 25  
   bone density, 26–27  
   cardiovascular adaptation, 29  
   cardiovascular and pulmonary systems, 28  
   cognitive and behavioral changes, 28  
   complications, 25  
   contracture deformities, 27  
   deep vein thrombosis, 29  
   digestive and excretory systems, 29–30  
   early mobilization, 30  
   endocrine, 29  
   heterotopic ossification, 27  
   intensive care unit survivors, 25  
   medical treatment, 25  
   metabolic disorders, 29  
   multidisciplinary approach, 25  
   muscle atrophy and fatigue, 26  
   musculoskeletal, nervous and integumentary  
     systems, 26–28  
   neuronal involvement, 28  
   orthosis, 26  
   pain, 28  
   pneumonia, 29  
   postural hypotension, 29  
   pressure ulcers, 28  
   vital function, 25  
   World Health Organization, 25  
 Immobility complications, 25  
 Immobilization, 25–30  
 Independent component analysis (ICA), 100  
 Inspiratory muscle training (IMT), 10  
 Intensify ischemia, 11  
 Interhemispheric hygroma, 21  
 Interleukin-6 (IL-6), 4  
 Interleukin-1 $\beta$  (IL-1 $\beta$ ), 4  
 Interleukin-10 (IL-10), 5  
 Internal readaptation programs, 118  
 The International League Against Epilepsy (ILAE), 49  
 Intracranial pressure (ICP), 11  
 Isotropic diffusion, 97

## K

Kickboxing, 79, 80, 82, 83  
 Knockout (KO)  
   concussion, 80

## L

Language disorders, 107  
 Levetiracetam, 53  
 Long-term learning memory, 111

## M

Magnetic resonance imaging (MRI), 96  
   conventional, 101  
   fMRI, 100  
   perfusion MRI, 100  
 Magnetic resonance spectroscopy (MRS)  
   proton, 100, 101  
 Major depressive disorder (MDD), 91  
 Manually assisted treadmill training  
   (MATT), 37  
 Matrix metalloproteinases (MMPs), 17  
 Maximal inspiratory pressure (MIP), 10  
 Mayo classification, 50  
 Mean diffusivity (MD) reflects, 98  
 Mean transit time (MTT), 100  
 Mechanical ventilation (MV), 9–12  
 Medication approach, 111  
 Memory  
   and attention deficits, 107  
   long-term, 107  
   rehabilitation, 110  
   speech therapists, 111  
   storage, 111  
   type of injury, 111  
   working, 107  
 Mental health diseases, 117  
 Mesial TLE (MTLE), 53  
 Metabolic disorders, 29  
 Methylenetetrahydrofolate reductase  
   (MTHFR), 51  
 Microglia, 3  
 Microthrombi, 18  
 Mild TBI (mTBI)  
   acute EEG, 63  
   causes, 71  
   chronic EEG, 63  
   classification, 69  
   and concussion, 70, 71  
   definition, 69  
   disability, 71  
   EEG abnormalities, 63  
   emotional and physical symptoms, 69  
   focal and diffuse brain injury, 69–70  
   management, 72  
   mortality, 70–71  
   PCS, 71, 72  
   primary and secondary injuries, 70  
   psychological and physical symptoms, 73  
   rehabilitation, 72–73  
   subacute EEG, 63  
 Mild traumatic brain injury, 87–88  
 Mixed-martial arts (MMA), 79, 82, 83  
 Molecular neuroimaging, 101–103  
 Mortality, 70–71  
 mTOR complex 1 (mTORC1), 51  
 Multiplanar T1-weighted images, 96  
 Muscle atrophy, 26  
 Musculoskeletal disabilities, 115  
 Musculoskeletal disturbances, 118  
 Myoclonus, 44

**N**

N-acetyl aspartate (NAA), 100, 101  
 National Football League (NFL), 88  
   chronic traumatic repetitive injury, 77  
   concussion protocol, 76  
   football activities, 77  
   game-replay review, 77  
   history, 75  
   league's head, neck and spine committee, 77  
   light aerobic exercise, 77  
   local and national USA policies, 75–76  
   policies and rules, 76  
   rest and recovery, 77  
   training, 77  
   UNC, 77  
 Nerve growth factor (NGF), 3  
 Neural progenitor cells (NPCs), 2, 3  
 Neural progenitor stem cells (NPSCs), 1, 2  
 Neural stem cells, 1, 2, 5  
 Neuroblasts, 1, 3, 4  
 Neurogenesis, 1–6  
 Neuromodulation, 35  
 Neuropsychological rehabilitation  
   cognitive impairments, 107–108  
   cognitive rehabilitation, 107  
   manage post-TBI patients (*see* Post-TBI patients)  
   Rancho Los Amigos scale (*see* Rancho Los Amigos scale)  
 Neurotrophins (NTs), 3  
 Nonconvulsive status epilepticus (NCSE), 50, 53  
 Noninvasive ventilation (NIV), 12  
 Notch signaling pathway, 2  
 Nuclear factor kappa B (NF- $\kappa$ B), 4

**O**

Obsessive-compulsive disorder (OCD), 89  
 Occupational therapy, 117, 119  
 Osteomuscular, 117  
 Oxyhemoglobin behaves, 100

**P**

Pain, 28  
 Parkinson's disease (PD), 42, 43, 87, 89  
 Parkinsonism, 43, 44  
 Penetrating ballistic-like brain injury (PBBi) model, 54  
 Perampanel, 54  
 Perfusion MRI, 100  
 Persistent post-concussion syndrome (PPCS), 72  
 PET-CT technology, 101  
 PET-MR, 101  
 Phenytoin, 53, 56  
 Pilot professional/vocational rehabilitation program (PPVRP)  
   adequacy, 119–120  
   anamnesis, 118  
   ARCO, 117  
   characterization, 118  
   cognitive rehabilitation outpatient clinic, 117  
   difficulties found and outcome of cases attended, 119  
   implementation and development, 118–119  
   LIIST, 117  
   limitations, 119–120  
   occupational therapy, 117  
   physical, 117  
   potentialities, 119–120

  process of education and initial proposal, 117–118  
   return to work  
     acute stage, 114  
     brain injuries, 115  
     capacity, 115  
     characteristics, 116  
     CVA, 116  
     description, 116  
     emotional, and social limitations, 115  
     facilitators, 116  
     guidelines, 115  
     musculoskeletal disabilities, 115  
     neurological program, 115  
     organizational characteristics, 115  
     outpatient stage, 115  
     post-TBI rehabilitation, 114  
     probability, 115  
     professional evaluation, 115  
     subacute stage, 115  
     type of attention, 115  
 Pneumonia, 29  
 Policy, 76  
 Positive end-expiratory pressure (PEEP), 10, 11  
 Positron-emission tomography (PET), 101–103  
 Post-concussion syndrome (PCS), 37, 71, 72  
   and CTE, 88  
   diagnosis, 88  
   neuropsychiatric symptoms, 89  
   unemployment rate, 91  
 Post-TBI patients  
   cognitive function evaluation, 110  
   cognitive rehabilitation (*see* Cognitive rehabilitation)  
 Post-traumatic amnesia (PTA), 62, 69  
 Post-traumatic epilepsy (PTE)  
   clinical/natural history, 52–53  
   EEG, 64  
   epidemiology, 51  
   epileptic seizure, 49  
   experimental interventions, 54–55  
   family history/genetic associations, 51  
   investigations, 53  
   management, 53–54  
   pathophysiology, 51–52  
   phenytoin, 56  
   post-traumatic animal models, 55  
   risk factors, 50  
   surgical intervention, 55  
   TBI, 49  
 Post-traumatic hydrocephalus (PTH)  
   blood products, 18  
   clinical manifestations, 19  
   clinical recovery, 17  
   Dandy's report, 17  
   decompressive craniectomy, 21–23  
   differential diagnosis, 19–20  
   direct trauma and apoptosis, 18  
   ependymal cilia, 17  
   hydrocephalus, 21–23  
   MMPs, 17  
   neuropsychological, 19  
   NPH, 19  
   outcome, 21  
   pathogenesis, 17–18  
   pathology, 18  
   rehabilitation, 21

- rehabilitation setting, 17
  - risk factors, 18
  - skull CT, 18
  - TBI, 17
  - traumatic hygroma, 21
  - treatment, 20
  - Post-traumatic seizures (PTS), 64
    - epidemiology, 50–51
    - risk factors, 50
  - Post-traumatic stress disorder (PTSD), 108
  - Postural hypotension, 29
  - Pressure ulcers, 25, 26, 28
  - Proton magnetic resonance spectroscopy (<sup>1</sup>H-MRS), 100, 101
  - Psychogenic non-epileptic seizures (PNES), 53, 54
  - Pulmonary embolism, 29
  - Punch-drunk syndrome, 88
- Q**
- Quantitative EEG (qEEG), 61–63, 65
- R**
- Radial diffusivities (RD), 98
  - Rancho Los Amigos scale
    - automatic-appropriate, 109
    - confused-agitated, 108–109
    - confused-appropriate, 109
    - confused-inappropriate, non-agitated, 109
    - generalized response, 108
    - localized response, 108
    - purposeful-appropriate, 109–110
  - Range of motion (ROM), 37
  - Regional cerebral blood flow (rCBF), 101, 102
  - Region-of-interest (ROI)s, 98
  - Rehabilitation
    - mTBI, 72–73
    - transdisciplinary, 73
  - Rehabilitation program
    - PPVRP (*see* Pilot professional/vocational rehabilitation program (PPVRP))
  - Reorganization, 5–6
  - Repeated mild TBIs (rTBIs), 79
  - Respiratory rehabilitation
    - acute and subacute phases, 9
    - intracranial hypertension, 9
    - mechanical ventilation, 9–12
    - physiotherapeutic interventions, 9
    - spontaneous ventilation, 12–15
    - TBI patients, 15
  - Responsive neurostimulation (RNS), 55
  - Resting-state fMRI (RS-fMRI), 100
  - Return to work
    - and PPVRP, 114–116
  - Robotic- assisted gait training (RAGT), 37
  - Robotic body weight-supported treadmill training (RBWSTT), 37
  - Robotic rehabilitation
    - advantages, 38
    - behavioral symptoms, 35
    - BIAA, 35
    - cognitive dysfunction, TBI, 35
    - conventional therapy, 38
    - disadvantage, 38
    - external stimulation, 35
    - neurologic disabilities, 35
    - neuromodulation, 38
    - robotic therapy, 36–38
  - Robotic-assisted gait training (RAGT), 37
  - Robotic-assisted treadmill training (RATT), 37
- S**
- Selective attention training, 110
  - Sensory evoked potentials (SEPs), 37
  - Short-lived radioisotopes, 101
  - Single photon emission computed tomography (SPECT), 101–103
  - Slit guide, 5
  - Smoothed (Smo) inhibition, 2
  - Social assistance, 116
  - Social inclusion
    - and work, 113–114
  - Social security, 116, 117
  - Solute carrier family 1 member 1 (SLC1A1), 51
  - Sonic Hedgehog signaling pathway (Shh), 2–3
  - Speech and language therapist, 21
  - Spinal cord injury, 36
  - Spontaneous breathing trial (SBT), 9
  - Spontaneous recovery, 1, 5, 6
  - Standardized low-resolution brain electromagnetic tomography (sLORETA), 64
  - Status epilepticus (SE), 50
  - Stroop test, 101
  - Subarachnoid and intraventricular hemorrhages, 95
  - Subarachnoid capillaries, 17
  - Subarachnoid hemorrhage (SAH), 18, 19, 21, 22
  - Subconcussion, 80
  - Subgranular zone (SGZ), 1, 3
  - Subventricular zone (SVZ), 1, 3, 5
  - Susceptibility-weighted imaging (SWI), 96, 97
  - Sustained attention training, 110
  - Synaptogenesis, 6
- T**
- T2\*-weighted gradient-recalled echo (T2\*-GRE), 96, 97
  - Task-based fMRI, 100
  - Tics, 44
  - Tochopherol, 54
  - Tracheostomy, 12
  - Tract-based spatial statistics (TBSS), 98, 99
  - Tractography, 98
  - Training strategies
    - outside the outpatient unit, 111
  - Transdisciplinary teams, 69
  - Transforming growth factor beta (TGF-β), 5
  - Traumatic brain injury (TBI)
    - acute and chronic stages, 64
    - acute phase, 11
    - Annegers' classification, 50
    - anxiety and depression, 61
    - APOEε4 allele, 51
    - ARDS, 11
    - astrocytes and microglia activation, 1
    - cerebral function, 62
    - critical stage, 27
    - EEG, 53, 62–63
    - Eph, 3
    - epileptic events, 49
    - eucapnia, 11
    - functional neuroimage (*see* Functional neuroimage)

- Traumatic brain injury (TBI) (*cont.*)  
functional plasticity, 6  
gastrointestinal tract changes, 30  
history, 96  
interleukin IL-1 $\beta$ , 4  
Interleukin-6 (IL-6), 4  
Interleukin-10 (IL-10), 5  
Mayo classification, 50  
mTBI, 63  
and neurogenesis, 1–3  
neurologic complication, 49  
neuropsychological rehabilitation (*see* Neuropsychological rehabilitation)  
non-epileptic patients, 52  
Notch signaling pathway, 2  
NTs, 3  
paresis/paralysis, 27  
PSG, 65  
and PTE, 52  
REM sleep, 65  
reorganization of connectivity, 5–6  
respiratory rehabilitation, 9  
seizure prophylaxis, 53  
severity, 50, 51  
Shh, 2–3  
sLORETA, 64  
social inclusion, and work, 113–114  
spontaneous recovery, 6  
synaptogenesis and plasticity, 1  
temporal phases, 51  
TGF- $\beta$ , 5  
TNF $\alpha$ , 4–5  
ventilatory management, 11  
Wnt signaling pathway, 2  
Traumatic subdural hygromas (TSH), 21  
Tremor, 41  
Tumor necrosis factor-alpha (TNF- $\alpha$ ), 4–5, 54  
Tyrosine kinases (Trk), 3  
Tyrosine receptor kinase A (TrkA), 3  
Tyrosine receptor kinase C (TrkC), 3
- V**  
Vagal nerve stimulation (VNS), 55  
Ventilator-associated pneumonia (VAP), 9  
Ventriculomegaly, 17, 19, 20  
Ventriculoperitoneal shunt (VPS), 19, 20, 22  
Vocational rehabilitation  
PPVRP (*see* Pilot professional/vocational rehabilitation program (PPVRP))  
Voxel-wise approaches, 98
- W**  
Wnt signaling pathway, 2  
Work and social inclusion, 113–114
- Z**  
Zackery Lystedt Law, 75