

# Index

- ab initio* modeling, 296, 297
- absorption corrections for x-radiation, 19
- activation barrier, 297, 304, 316
- activation volume of RDX, 55, 59
- ammonium nitrate (AN), 99
- ammonium perchlorate (AP), 110, 111, 117, 123
- diffraction spectra, 110
  - static compression, 111
- amorphous nitrogen, 92
- anisotropic sensitivity, 242
- anthracene, 228, 231–234, 247
- Astin, 4
- azide, 22–25, 75, 76
- band structure calculations, 296, 320
- basis set, 304, 317
- Bassett, W. A., 104
- benzene, 19–21
- Birch-Murnaghan EOS formalism, 104, 105
- Block, S., 15, 17, 25
- Bridgman opposed anvils, 7, 8, 34
- Bridgman, P. W., 5, 7, 23
- bromine, 19, 21
- bulk EM representation, 261
- bulk modulus, 102, 104–106, 109–111, 121
- Bunting, E. N., 2, 3, 5, 6, 9, 14
- Cady, H., 104, 108
- calibration of ruby R-line pressure shift, 25
- precise, 25
  - preliminary, 25
- carbon disulfide, 21
- carbon tetrachloride, 20, 21
- Cassegrain optics for FTIR, 52
- charge trapping, 316
- chemical decomposition, 221, 223
- chemical reaction, 292, 294–296, 311, 312, 316, 319–321
- CL-20, 103, 111, 112, 117, 118, 120, 123, 216
- decomposition by laser initiation, 216
  - products of decomposition, 216
  - static compression, 112
- COMPASS force field, 278, 279
- compressibilities, 23–25
- Compressibility of Explosive Liquids, 64
- NM, 67
- CONO formation, 317, 318
- crystal structure, 235, 243
- crystallography of RDX phases, 46, 47
- CsF transition, 18
- cubic gauche structure, 76, 77, 81, 82, 89, 94
- Debye temperature, 260
- decomposition kinetics, 206, 209
- decomposition of AP, 210
- decomposition rate constants, 55, 58, 59
- decomposition reaction, 291, 294, 316
- defect, 297, 304–309, 311, 315, 320
- deformation, 300, 301, 303, 308, 313–315
- density functional theory, 296, 297, 304, 317
- detonation initiation, 294–296, 316
- deviatoric stress, 228, 230, 231
- di-vacancy (vacancy dimmer), 297
- diamond, 76, 79–86, 89–94
- diamond anvil cell (DAC), 78, 79, 82, 92, 220, 229, 230, 247
- Bassett, 41
  - dynamic diamond cell, 68
  - Holzappel, 41
  - Mao-Bell, 39
  - Merrill-Bassett, 42
  - NIST, 1–3, 12, 33, 38, 39, 41, 45, 52, 69

- dilatometry, 176  
 dynamic compression, 175  
 Hugoniot, 176, 177  
 shock compression, 177
- dimer, 233, 234
- dislocation, 222, 229, 233, 234, 241, 257, 294, 297, 299–303, 311, 313, 315, 316, 319, 320
- dissociation, 291, 292, 294, 296–298, 303, 304, 308, 310–312, 314, 315, 317–319, 321
- dynamic compression, 221, 223
- elastic properties of TATB, 277
- electronic properties, 297, 312, 319
- electronic structure, 228, 231–233, 246, 296–299, 301, 305, 310, 312, 313, 315, 317, 319, 320
- energetic barrier, 292, 299
- energetic material (EM), 128, 132, 134, 219–221, 223, 225, 226, 234, 235, 241, 247, 255–262, 264–269, 278–281, 291, 292, 294–297, 302, 303, 307, 311, 316, 319
- energy transfer, 221
- equation of state (EOS), 98, 100–111, 113, 115, 120, 121, 123, 124, 255, 256, 258, 261, 269, 270, 272
- Estane 5703, 128, 131, 132, 156, 160, 163, 171, 172, 180, 187, 189
- Estane<sup>TM</sup> force field, 275
- excimer, 233
- excitations, 291, 295–297, 299, 301, 302, 311–316, 318–321
- excited state, 292, 311, 313–316, 320
- exciton, 223, 295
- excitonic mechanism, 315, 320
- experimental kinetic measurement methods, 203  
 combustion front propagation (CFPR), 209  
 single-shot laser initiation, 211  
 time-resolved absorbance, 210  
 time-to-explosion, 208
- explosives, 75, 76, 98, 100, 103–105, 108–110, 112, 114, 120, 127–130, 132, 133, 153, 161, 163, 178, 180, 181, 184, 190, 191, 193  
 high, 99, 100, 103, 112  
 low, 99
- first principles, 296, 319
- fixed-point pressure scale, 39, 67
- fluorescence, 225, 229, 231–233, 246
- fluorescing materials, 25
- force fields, 258, 262, 266–270, 272–281
- Forman, R., 25
- formulations, 129, 131, 132, 134, 152, 153, 160, 176  
 binders, 129, 131, 132, 134, 153, 160  
 equation of state, 134, 152, 176
- FOX-7 (diamino-dinitroethylene, C<sub>2</sub>H<sub>4</sub>N<sub>4</sub>O<sub>4</sub>), 296, 303–312, 314, 316–321
- freezing pressures of liquids, 23, 25, 31
- FTIR, 2, 46, 48, 52, 53, 56, 69
- gas gun, 181, 223  
 spectroscopy, 181
- Gasket Technique, 14, 15
- General Services Administration, 9
- glass transition pressure, 31
- Helmholz free energy, 101, 102
- high energy density material, 75, 77
- high explosives, 220, 291, 296, 312, 315, 320
- high pressure, 76, 77, 85, 87, 88, 90–93, 220, 229, 231, 232, 234–236, 241, 243, 244, 247
- High Pressure Diamond Optics, 14
- high pressure phases, 98, 112, 114, 116
- hindered shear, 242, 245
- HMX, 99, 103, 105–107, 115, 121, 123, 209, 212–214  
 absorbance change, 212  
 advanced interface model, 213  
 by time-resolved-absorbance, 210  
 chain-type model, 213  
 decomposition mechanism, 214  
 EOS parameters, 123  
 high pressure polymorphs, 105  
 pressure/volume data, 104  
 rate determining step, 214
- HNIW, 103, 111, 117  
 compression, 111  
 high pressure polymorphs, 111, 117  
 phase diagram, 118  
 XRD patterns, 121
- HONO formation, 317
- hot spots, 203, 205, 207, 208, 257
- Hot-Molecular Zone (HMZ), 203
- Hugoniot, 104, 105, 110, 268
- hydrogen transfer, 294, 308
- hydrostatic limits, 28
- hydrostatic pressures, 15, 22, 28, 31, 69
- hydrothermal diamond cell (HDAC), 104
- ice VI, 15, 16, 19, 68
- inter-molecular interactions, 293, 294, 299, 305, 307

- interactive potentials, 259, 264–267, 269–272, 277
- intra-molecular interactions, 299, 318
- invention, 1–3, 69
- ionic reactions, 242, 245
- IR spectra of RDX phases, 48, 55
- IR spectroscopy, 69, 110, 113
- ir spectroscopy, 7, 17
- isomerization, 294, 304, 308, 317, 318
- isothermal EOS parameters, 108
- isotherms, 106, 122, 124
- Kel-F 800, 128, 129, 132, 133, 153, 154, 156, 158, 159, 163, 166, 171, 174, 184–187, 189
- key MD simulations, 266
- kinetic models, 204, 206, 212, 213
- advancing interface, 206, 212, 213
- chain-type nucleation, 207, 212, 213
- Lee-Tarver model, 207
- sigmoid, 204, 206, 212
- La transition, 18
- laser heating, 76, 78, 82, 83, 85–87, 89, 94
- laser spectroscopy, 225
- Lee-Tarver model, 210
- Lippincott, E. R., 2, 3, 5, 6, 14
- McMurdie, H. F., 4, 5, 9, 17
- MD simulations, 255, 256, 259, 261, 262, 264–274, 277–280
- mechanical anisotropy, 241, 242
- Menikoff, R., 257
- Merrill-Bassett cell, 209
- mesoscale, 256, 258, 275
- metastable polymeric nitrogen, 94
- methanol:ethanol:water mixture, 30, 31, 44
- methods of MD, 258
- Hamiltonian, 259
- Lagrangian, 259
- Newtonian, 259
- molecular conformers, 242, 243
- molecular crystals, 219–221, 228, 229, 231, 234, 236, 247
- molecular dissociation, 308
- molecular dynamics (MD), 254, 258, 273, 276, 280, 314
- molecular orientation, 303
- molecular structure, 238, 241, 243, 244
- Nd:YAG laser, 214, 216
- nitro-nitrite isomerization, 294, 304
- nitrogen, 75–94
- nitroglycerine, 99
- Nobel Prize, 5
- nonhydrostatic pressures, 22, 29, 53
- nonhydrostaticity, 227–231
- nonmolecular nitrogen, 76–78, 89, 92
- nonreactive force fields, 273
- nonreactive models, 269
- Olinger, B., 104–109, 116
- optical absorption, 300, 313, 315
- optical fluorescence system, 32
- optical polarizing microscopy, 2, 24, 43
- optical properties, 297
- optical spectroscopy, 225
- orientational defects, 303
- periodic model, 296
- PETN, 109, 110, 116, 123, 239, 241–247
- static compression, 110, 117
- PETN (pentaerythritol-tetranitrate,  $C_5H_8N_4O_{12}$ ), 296, 315
- phase diagrams, 45, 100, 115, 117, 118, 120, 123
- NM, 50
- RDX, 45
- phase transition, 226, 235–238, 240
- photoluminescence, 245, 246
- Picatinny Arsenal, 22
- Piermarini, G. J., 4, 213
- plane waves, 304
- polymer materials, 133, 142
- polymeric nitrogen, 76–79, 82, 83, 86, 87, 91, 92, 94
- polymers, 127–129, 132, 133, 135, 142, 143, 146–150, 152–155, 157, 158, 161–163, 166–168, 170–175, 177, 178, 180, 182, 184, 186, 188, 190, 191, 193–195
- polymorphism, 234–236, 247
- polynitrogen, 75
- polytetrafluoroethylene, 131
- potassium nitrate transition, 21
- potential energy surface, 258, 260, 267, 277
- powder x-ray diffraction, 48
- pressure measurement, 16, 24, 28, 32, 36, 43, 45, 65, 69
- propellant, 77
- pseudopotentials, 304
- Raman, 163, 167
- high Pressure, 163, 167
- Raman spectroscopy, 114, 115, 225, 226, 236, 240, 243, 247
- RDX, 99, 103, 105–107, 114, 123, 235–240, 247
- high pressure polymorphs, 105

- RDX (cyclotrimethylene-trinitramine,  $C_3H_6N_6O_6$ ), 296–306, 312–315, 320
- RDX decomposition, 51, 54, 55, 57, 58, 214
- reaction energy, 291, 311, 313, 317, 318, 320
- reaction initiation of solid HEs, 204
- coalescence, 206
  - induction period, 205
    - deceleration, 205
    - growth process, 205
    - ingestion, 206
    - nucleation, 205
  - reaction rates, 203, 206, 207, 212, 213, 215, 217
  - reactive model (ReaxFF), 265, 268, 269, 280
  - reversed molecule, 305, 306
  - ruby, 229, 230, 232
  - Ruby fluorescence, 2, 24, 28, 30, 33, 36, 45, 46, 69
- shear strain, 297, 301, 308–311, 314, 319, 320
- shear stress, 228, 229
- shock compression, 219–221, 225, 228, 231–233, 235, 238, 239, 241, 244, 245, 247
- shock Hugoniot, 255, 256, 259, 267, 273
- shock initiation, 221, 238, 241
- shock wave, 219–221, 223–230, 235, 238, 241, 245, 247, 291, 296, 299, 311–313, 320
- shock-induced decomposition, 223, 245
- single crystal x-ray diffraction, 14–17, 20, 21, 23, 32, 42
- single crystals, 2, 17, 21, 31, 45, 47, 61, 65
- singly bonded nitrogen, 75
- slip planes, 245
- Smithsonian Institution, 7
- sodium perchlorate, 99
- specific volume of NM, 67
- static compression, 220, 221, 227–229, 231, 233, 235, 236, 238, 239, 243, 245
- static high pressure, 203, 204, 209, 211, 214, 216, 217, 220, 229, 231, 235, 236, 241, 243, 244, 247
- static high pressure MD simulations, 260
- Stokes Falling Ball method, 30, 69
- strain energy, 102
- stress, 219–221, 224, 226–233, 238, 240, 241, 243, 245–247, 301–303, 312, 319, 321
- structural defects, 229
- superhard, 76, 93, 94
- superpressed state of  $H_2O$  and  $D_2O$ , 68, 69
- surface, 291–293, 297–299, 302, 311–314, 319
- Sylgard 184, 163, 167, 171, 172, 175, 189
- symmetry-adapted perturbation theory (SAPT), 277, 278
- synchrotron, 87
- TATB, 99, 108, 109, 115, 121, 123, 215, 216
- absorbance, 215, 216
  - Compression, 109
  - diffraction spectra, 116
  - high pressure polymorphs, 121
  - laser-initiated decomposition, 215, 216
  - photolysis, 216
- TATB (triamino-trinitrobenzine,  $C_6H_6N_6O_6$ ), 293, 294, 296, 303, 304, 306–311, 316
- thermal decomposition rates, 52, 55, 61, 63
- THV 500, 156, 158, 159, 162, 163, 166, 182, 185, 186
- time resolved spectroscopy, 225, 226, 240
- TNT, 293, 294, 315
- trinitrotoluene (TNT), 99, 124
- type I diamond, 10
- Type II diamond, 8, 9
- ultrafast spectroscopy, 295, 321
- ungasketed technique, 31, 34
- uniaxial strain, 225, 228, 229, 238, 241, 243
- University of Maryland, 5, 6
- vacancy, 297, 298, 315, 319
- Van Valkenburg, A., 2–5, 14
- vibrational structure, 231
- vibron, 77, 81, 84–86, 90
- viscosity of liquids, 30
- volume thermal expansion, 107
- Wachtman, J. B., 25
- Weir, C. E., 2–4
- x-ray diffraction, 78–83, 85, 87–90
- x-ray precession camera, 16, 18, 20, 21, 23
- zinc sulfide transition, 44