

Index

A

- Activin, and Smad, 130
- Aging of the immune system, 162–163
- Amphibian spleen, 2
 - and melano-macrophage centers (MMC), 2
- Angiopoietins, 54, 55
- Antigen transport and reticula in old mice, 167–168
- Avian spleen, 123

B

- Bapx1 transcription factor, 128–130
- B cell follicles, 59, 60, 63–67, 69
- B-1 cells, 132
- Blood vasculature, 66, 68

C

- CCL19, 42, 43, 81, 86, 144, 145
- CCL21, 42, 43, 79, 81–83, 86, 87, 89, 144, 145, 151–153
- CCR7, 42, 43, 81–83, 86
- CD34, 85, 86
- CD30L, 149–153
- CD4 memory, 148–149, 151–152
- Chemokine receptor
 - CCR6, 110, 112, 113
 - CCR7, 113
 - CCR9, 113
 - CXCR5, 112, 113, 115
- Chemokines, 39–43, 144–146, 150
 - CCL19, 110, 112
 - CCL20, 110, 113
 - CCL25, 113
 - CXCL13, 110, 112, 113
- Commensal bacteria, 109, 110
- Conduits, 68, 69, 125, 127, 133, 143–146

- Core 2 branching enzyme (Core 2 GlcNAcT), 85–86
- Cryptopatches (CP), 107–116
- CXCL12, 83
- CXCL13, 42, 43, 80, 81, 83, 89, 144–147, 150, 151, 153
- CXCR4, 81, 83
- CXCR5, 41–43

D

- Deafferentiation, 87, 88
- Differences in spleen structure, 124
- Dominant hemimelia (Dh), 128

F

- Fibroblastic reticular cells (FRCs), 77, 79, 85, 88
- Fibroreticular stromal cells (FSC), 143–144, 146, 147, 150, 152, 153
- Follicle-associated epithelium (FAE), 98, 108, 109, 114
- Follicular dendritic cells (FDCs), 60, 63, 65, 66, 69, 103
 - in aging, 165–167
 - relationship to other dendritic cells, 163
- Fucosyltransferase VII, 85

G

- Germfree mice, 109, 110
- Germinal centers (GCs), 59, 60, 66, 99
- GlcNAc6ST-2, 85, 87
- GlyCAM-1, 85–87

H

- HEVin, 86
- High endothelial venules (HEVs), 76–89
- High endothelial vessels (HEVs), 65, 67–69
- HOX11 transcription factor, 121, 128, 129

I

- ICAM-1, 81, 82, 84
- ICAM-2, 82
- Id2*, 100
- Id2 transcription factor, 10, 11
- IgA, 99, 103
- Ikaros transcription factor, 9–11
- IL-7 and Peyer's patches, 40, 41
- IL-7 receptor, 39–42
- Immunoglobulin A (IgA), 109, 110, 113–115
- Inflammation and LEC cells, 56
- Inflammatory macrophages, 55–56
- Integrin, 112
- Interferon (IFN) β , 18
- Interleukin 7 (IL-7), 41
 - IL-7 and lymphopoiesis, 39–40
- Intraepithelial lymphocytes (IEL), 108, 113, 114
- Isolated lymphoid follicles (ILFs), 29, 33, 34, 107–116

L

- LFA-1, 81–83
- LIGHT, 17, 18
- L-selectin, 81–86
- LT α , 149–151
- LT β , 149–151
- LT β R-Ig decoy receptor, 17, 18
- LT receptors, 15, 19, 20
- Lymphatic, 61, 63, 65–69
- Lymphatic capillaries, 51, 55
- Lymphatic circulation, 51
- Lymphatic endothelial cells (LECs), 49–56
 - and Kaposi sarcoma, 50
- Lymphatic vasculature, 61, 66
- Lymph node hierarchy, 17
- Lymphoid tissue inducer (LTi) cells, 60, 76, 80, 89, 99–102, 108, 110–114, 116, 149–150
 - discovery, 8
 - phenotype, 8
- Lymphoid tissue initiator (LTin) cells, 99–102
- Lymphoid tissue organizer cells (LTo)
 - origin, 11–12
 - phenotype, 9
- Lymphotoxin (LT), 15–21, 111, 113, 114
- Lympho-toxin beta receptor (LT β R), 26–32, 34
- Lymphotoxin β (LT β)/LT β receptor, 76
- LYVE-1 antigen, 54

M

- Marginal reticular cells (MRC), 9, 12, 126
- Marginal sinus (MS), 125, 126, 130, 134–136, 143, 144, 146, 147

- Marginal zone (MZ), 123, 124, 133, 143, 146–148, 153
 - marginal zone macrophages, 125, 135
- Marginal zone B (MzB) cells, 135
- M cells, 98, 99
- MECA 79, 82, 85, 86
- MECA 367, 82
- Mesenchymal LTo cells, 76
- Mesenchymal scaffolding, 126, 132–136
- Microflora, 109, 111
- Molecular model of FDC–B cell interactions, 163–165
- Mucosal addressin cell adhesion molecule (MAdCAM)-1, 80–83, 85, 87, 88

N

- Nfkb1*, 31
- Nfkb2*, 27, 28, 30, 31
- NF- κ B-inducing kinase (NIK), 27, 28, 30–31, 34
- Nkx2-3 transcription factor, 130
- Nkx2-5 transcription factor, 129
- Nkx3.2 transcription factor, 128
- Nuclear Factor kappa B (NF- κ B), 25–34

O

- Old B and T cells, 168–170
- Oplop* mice, 125
- OX40L, 149–153

P

- Paracortex (T cell area), 77
- Pbx1 transcription factor, 129
- Periarteriolar lymphoid sheath (PALS), 123–127, 133
- Pericytes, 78, 79, 84
- Peyer's patches (PP), 107
- PNAd, 80–89
- Podoplanin, 144, 145, 152, 153
- Prox1 transcription factor, 54, 55

R

- RANK, 26, 27, 29, 31–33
- Red pulp, 122–127, 130–135
- RelA, 26, 27, 31–33
- RelB, 26–28, 30–34
- RET, 100, 101
- Retinoic acid related orphan receptor (ROR γ t), 61, 62, 112, 114
- Reverse signaling and TNF, 20
- ROR, 10
 - ROR γ t, 10, 11
- Rory*, 100

S

Sonic hedgehog (Shh), 128, 130
and smoothened, 129
Splanchnic mesodermal plate (SMP), 127
Spleen, neonatal, 131
Spleen ontogeny, 121–124
Splenic ellipsoid, 123
Splenic embryonic vessels
in humans, 131
in mouse, 134
Splenic marginal zone, 147
Splenic phylogeny, 148
Streptococcus pneumoniae, 69
Stromal cells, 60, 63–67, 101–103
Stromal organizer cells, 111, 112
Subcapsular sinus (SCS) macrophages, 60
Szakal, A.K., 161–170

T

Tal-1/Scl transcription factor, 52, 53
Taub, D.D., 162
Thymic stromal lymphopoietin (TSLP),
40, 41

Tlx1 transcription factor, 121, 128, 129
TNF α , 26, 32
TNF receptors
ligand binding, 19
types, 16
Toll like receptor (TLR), 109, 110
TRAF, 19–21
association with TNF and LT receptors, 20
TRANCE
TRANCE and LT β R, 19
TRANCE and LTi, 19
Tumor necrosis factor (TNF), 15–21

V

Vascular cell adhesion molecule-1 (VCAM-1),
83, 84
Vascular endothelial growth factor (VEGF),
52–56
receptors, 53
VEGF-A, 88, 89

W

White pulp, 121–127, 130–135