

# Index

## A

Adiabatic connection method (ACM), 50  
AgGaGe<sub>3</sub>Se<sub>8</sub>, 15  
AgGaGe<sub>n</sub>S(e)<sub>2(n+1)</sub>, 16  
AgGaS<sub>2</sub>, 9, 11, 46, 122  
AgGaSe<sub>2</sub>, 9, 47, 122  
Antimony, 121  
Asymmetric units, 121  
AZrPSe<sub>6</sub>, 31

## B

BaGa<sub>4</sub>S<sub>7</sub>, 14, 121  
Ba<sub>2</sub>In<sub>2</sub>S<sub>5</sub>, 130  
Ba<sub>12</sub>In<sub>4</sub>S<sub>19</sub>, 131  
BaMgF<sub>4</sub>, 24  
Becke's parameter-fitting hybrid methods, 50  
β-A<sub>2</sub>Hg<sub>3</sub>M<sub>2</sub>S<sub>8</sub>, 19  
Borates, 1  
Borophosphates, 3  
Bridgman–Stockbarger (BS) growth/  
technique, 3, 121, 132

## C

CdGeAs<sub>2</sub>, 10  
CdSiP<sub>2</sub>, 6  
Chalcogenides, 1, 4, 6, 121  
multinary, 121  
Chalcogenoborate, 34  
Chalcohalides, 33  
Chalcopnictates, 26  
Chalcopyrites, 4, 8  
Charge transfers, 45  
Conductor-like screening model (COSMO), 62

cPW1PW91, 51  
Cr(CO)<sub>3</sub>Bz, 53  
Crystal growth, 81  
CsAg<sub>2</sub>TeS<sub>6</sub>, 26  
CsCdBr<sub>3</sub>, 25  
CsGeCl<sub>3</sub>, 25  
CsHg<sub>3</sub>I<sub>8</sub>, 25  
Cs<sub>5</sub>P<sub>5</sub>Se<sub>12</sub>, 30

## D

Density functional (DF) calculations, 45, 50  
Difference frequency generation (DFG), 6  
Different-bond-type hybrid compounds, 26

## E

Er<sup>3+</sup>:SrWO<sub>4</sub>, 101  
Eu<sub>2</sub>Ga<sub>2</sub>GeS<sub>7</sub>, 130

## F

FeCp<sub>2</sub>COMe, 53

## G

GaSe, 6

## H

Halides, 1  
(Hg<sub>8</sub>As<sub>4</sub>)(Bi<sub>3</sub>Cl<sub>13</sub>), 33  
HgGa<sub>2</sub>S<sub>4</sub>, 14  
(Hg<sub>6</sub>P<sub>3</sub>)(In<sub>2</sub>C<sub>10</sub>), 33

Hyperpolarizability, 55  
Hyper-Rayleigh scattering (HRS), 67

**I**

Infrared second-order NLO materials, 45  
Intraligand charge transfers (ILCT), 49  
IR absorption, 45

**K**

KGd(WO<sub>4</sub>)<sub>2</sub> (KGW), 81  
(K3I)[SmB12(GaS4)3] 35  
K<sub>2</sub>Ln<sub>2</sub>Sb<sub>2</sub>S<sub>9</sub>, 123  
K<sub>2</sub>P<sub>2</sub>Se<sub>6</sub>, 29  
Kohn–Sham density functional theory, 51  
KZrPSe<sub>6</sub>, 31

**L**

LaCa<sub>2</sub>GeS<sub>4</sub>Cl<sub>3</sub>, 36  
La<sub>3</sub>CuGeSe<sub>7</sub>, 21  
La<sub>2</sub>Ga<sub>2</sub>GeS<sub>8</sub>, 130  
La<sub>4</sub>FeSb<sub>2</sub>S<sub>10</sub>, 123  
La<sub>4</sub>GaSbS<sub>9</sub>, 125  
Large crystals, 1  
Laser-induced damage threshold (LIDT), 9  
LiB<sub>3</sub>O<sub>5</sub>, 116  
Li<sub>2</sub>CdGeS<sub>4</sub>, 19  
Li<sub>2</sub>Ga<sub>2</sub>GeS<sub>6</sub>, 18, 122  
Ligand-independent mechanism, 50  
Ligand-to-ligand charge transfers (LLCT), 49  
LiMX<sub>2</sub>, 12  
Ln<sub>2</sub>Mn<sub>3</sub>Sb<sub>4</sub>S<sub>12</sub>, 123  
Ln<sub>4</sub>InSbS<sub>9</sub>, 125

**M**

Main group elements, 121  
Metal  
    chalcogenides, diamond-like, 16  
    halides, binary/ternary, 21  
    pnictidohalides, 4, 33  
Metal–metal interaction CT (MMCT), 49, 61  
Mid-infrared, 1  
MLCT/LMCT, 49  
MMCT. *See* Metal–metal interaction CT (MMCT)  
[MoCu<sub>2</sub>S<sub>4</sub>(SPh)<sub>2</sub>]<sub>2</sub><sup>−</sup>, 61  
[Mo<sub>2</sub>CuS<sub>4</sub>]<sup>1−</sup>(*edt*)<sub>2</sub>(PPh<sub>3</sub>), 61  
[MS<sub>4</sub>Cu<sub>4</sub>X<sub>2</sub>Py<sub>6</sub>], 67

**N**

NaSb<sub>3</sub>F<sub>10</sub>, 26, 47  
Na<sub>0.5</sub>Pb<sub>1.75</sub>GeS<sub>4</sub>, 20  
Na<sub>9</sub>Gd<sub>5</sub>Sb<sub>8</sub>S<sub>26</sub>, 123  
Nd:YAG–SrVO<sub>4</sub>/LBO yellow laser, 116  
Nd:YAG–SrWO<sub>4</sub>, 111  
    KTP yellow laser 113  
Nd:YVO<sub>4</sub>–SrWO<sub>4</sub>, 106  
Nd<sup>3+</sup>:KGd(WO<sub>4</sub>)<sub>2</sub>, 83  
Nd<sup>3+</sup>:SrWO<sub>4</sub>, 89  
Niobates, 3  
Noncentrosymmetry (NCS), 3, 122  
Nonlinear optical (NLO) crystals, 1, 45

**P**

Pb<sub>2</sub>GeSe<sub>4</sub>, 15  
Pentanuclear clusters, 67  
Perdew–Wang gradient-corrected correlation functional, 51  
Phosphates, 3  
Pnictides, 1  
Pnictidohalides, 4  
Potassium dihydrogen phosphate (KDP), 46

**Q**

Quasi-phase matching (QPM), 3

**R**

Raman, 81  
RbAg<sub>2</sub>TeS<sub>6</sub>, 26  
RbCdI<sub>3</sub>·H<sub>2</sub>O, 25  
Re<sub>2</sub>(allyl)<sub>4</sub>, 54  
Re clusters, dinuclear, second-order NLO, 54  
Re<sup>3+</sup>:KGd(WO<sub>4</sub>)<sub>2</sub>, 83  
Re–Re bonding, 56  
Re<sub>2</sub>(μ-S)<sub>2</sub>O<sub>2</sub>(CH<sub>2</sub>CMe<sub>3</sub>)<sub>4</sub>, 54

**S**

SbQ<sub>x</sub>, 124  
[Sb<sub>7</sub>S<sub>8</sub>Br<sub>2</sub>](AlCl<sub>4</sub>)<sub>3</sub>, 36  
Scheelites, 89  
Second-harmonic generation (SHG), 3  
Self-frequency conversion, 81  
Semiconductor compounds, 4  
Sm<sub>4</sub>GaSbS<sub>9</sub>, 125  
SrWO<sub>4</sub> crystals, 81, 89  
Stimulated Raman scattering (SRS), 81

**T**

Thioborates, 26

Thiotellurites, 26

 $\text{Tl}_4\text{HgI}_6$ , 24 $\text{Tl}_3\text{PbBr}_5$ , 22 $\text{Tl}_3\text{PbCl}_5$ , 22 $\text{Tm}^{3+}:\text{KGd}(\text{WO}_4)_2$ , 89 $\text{Tm}^{3+}:\text{SrWO}_4$ , 101

Transition-metal compounds, 45

Trinuclear anionic clusters, 61

**W** $\text{W}(\text{CO})_5\text{PyCHO}$ , 53**Z**

Zinc-blende compounds, 8

 $\text{ZnGeP}_2$ , 9, 47, 122 $\text{ZnSiP}_2$ , 8 $\text{ZnY}_6\text{Si}_2\text{S}_{14}$ , 20