

Index

Note: Page numbers followed by “f” and “t” indicate figures and tables respectively

0–9

- 1,4-benzoquinone, 74, 78, 79, 82*t*, 91, 93, 95, 98, 99
 - dependence of induction period on concentration of, 90*f*
 - under variation of
 - FMN concentration, 80*f*
 - luciferase concentration, 79*f*
 - order of bioassay components, 81*f*
 - time of reagent incubation, 82*f*
- 17 α -ethynylestradiol (EE2) testing, 137
- 2-hydroxy-3',4'-dichlorobiphenyl, 127
- 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD), 130, 133, 134, 135, 138
- 2,4'-dichlorodiphenyldichloroethylene (2,4'-DDE), 136
- (2S,4S)-2-methyl-2,3,3,4-tetrahydroxytetrahydrofuran borate, 49
- 3-hydroxybutanoyl-HSL, 48–49
- 3-oxo-hexanoyl-homoserine lactone, 49
- 3':5'-cyclic AMP (cAMP), 49
- 4a-hydroxy-FMNH product, 51
- 4n-nonylphenol, 136
- (4S)-4,5-dihydro-2-(6-hydroxy-2-benzothiazolyl)-4-thiazolecarboxylic acid, 15
- 6,7-dimethyl-8-ribityllumazine, 46
- Alcohol dehydrogenase (ADH), 87, 90, 95, 96, 98, 99, 101*t*
 - activity measurement, scheme of, 89*f*
 - activity under different concentration of deltamethrin, 102*f*
 - biomass of algae and, 94*f*
 - in coupling of enzymatic reactions, 88*f*
 - in pond water study, 93*f*
- Aliivibrio*, 23, 39, 40*t*, 41, 47, 51
 - regulatory genes, 46
- Alkane aliphatic hydrocarbons, 122–124
 - in seawater, 123
- Alkanes, 122, 123, 124, 126, 139
- Alpha-emitting radionuclides, 103
- Antigen–antibody complex (Ag–Ab), 177, 192
- Aposematism (warning display), 4, 5, 7–8
 - in Lampyridae, 7
 - luminescence
 - in centipedes, 8
 - in millipedes, 8, 8*f*
 - in marine organisms, 8–9
 - in mushrooms, 8
 - in poisonous butterflies, 8
- Aryl hydrocarbon receptor (AhR), 130
- Autoinducers, 47, 48–49, 163
- Automated capillary electrophoresis, 184

A

- Abortive infection, 159
- Acetate kinase, 180
- Acinetobacter baylyi* ADP1, 123, 124
- Acinetobacter* bioreporter
 - DF4/PUTK2, 128
 - DF4-8, 128
- Acyl-homoserine lactone (acyl-HSL) synthase, 46
- AhR nuclear translocator (ARNT) gene/protein, 130, 134, 135

B

- Bacterial bioluminescent assays, detection of organic compounds using, 114–115
- benzene, toluene, ethylbenzene, and xylene (BTEX), 115–120. *See also* BTEX bioreporters
- alkane aliphatic hydrocarbons, 122–124
- biphenyls, 125–127
- chlorinated aliphatic hydrocarbons, 124–125
- phenol and derivatives, 127–128

- Bacterial bioluminescent assays, detection of organic compounds using (*cont.*)
 polychlorinated biphenyls, 125–127
 polycyclic aromatic hydrocarbons, 120–122. *See also* PAH bioreporters
- Bacterial luciferase, 42, 157
 bioluminescent system, 70
 shortcomings of, 70
 enzyme-substrate interactions, 69
 luciferase system, 70
 NAD(P)H:FMN-oxidoreductase system, 70
- Bacterial luminescence
 biochemistry of, 42–43
lux genes (*see lux* genes)
 origin and function of, 51–53
 fatty-acid reductase genes, 52
 flavin cofactor, 51
 growth dependence actors, 52
 halotolerant flavodoxin, 53
 species of, 39–42
 ecological incidence and diversity, 42
 and ecological sources of, 40–41*t*
 not-yet cultured species, 41–42
- Bacterial *lux* genes. *See lux* genes
- Bacteriophages, 156, 157, 159, 160, 161–162, 164, 165, 166
 bioluminescent, 167
- Batesian mimicry, 5, 8
- BEAMing technology, 189
- Beta-emitting radionuclides, 103
- Beta-galactosidase, 157
- Bioluminescence
 application viability of, 69
 assays, reagents for (*see* Bioluminescent assays)
 by bacteria, 38
 colonies of *P. mandapamensis*, 39*f*
 luciferase, 42
 origin and function of (*see under* Bacterial luminescence)
 oxygen-dependence of, 38
 species of (*see under* Bacterial luminescence)
 cDNA coding, 176
 kinetics parameters, 74
 as reporter system, 158
- Bioluminescence, function of
 camouflage, 12–13
 functionless bioluminescence, 14
 intraspecific communication, 12
 light organ lure, 10
 attraction of spore-dispersing organisms, 11
 in prey attraction, 10–12
 searchlight, 13
 night-vision illumination, 13
 self-defense, 4–10
 aposematism, 4, 7–8
 Batesian mimicry, 5
 photophores, 5, 6
- Bioluminescent assays
 application areas, 75–76
 enzymatic bioassays. *See* Enzymatic bioluminescent bioassays
 enzyme incubation, 81–83
 existing methods, disadvantages of, 75
 immobilized enzymes, 76
 varying the composition of, 78–80
 of infectious agents, 185–186
 NADH:FMN-oxidoreductase-luciferase (*see* NADH:FMN-oxidoreductase-luciferase)
 in oncology diagnostics, 187–189
 reaction mixture, varying order of components, 80–81
- Bioluminescent binding assay, in homogeneous format, 192–193, 193*f*
- Bioluminescent enzyme system technology (BEST), 70
- Bioluminescent organisms
 deep-sea benthos, 26–27
 defensive reactions, 4–5
 diversity of, 5*f*
 marine taxa, 19
 coastal water, 21–22
 intertidal zone, 19–20
 in midwater, 23–26
 occurrence, 4
 prey attraction, hypotheses, 10
 terrestrial taxa, 14
 in air, 16–17
 in freshwater, rarity, 17–19
 on land, 14–16
 in underground, 17
- Bioluminescent plaques, 160
- Bioluminometric SNP genotyping, 191*f*
- Biotin-dUTP, 191
- Biotin–streptavidin conjugation, 184
 and avidin system, 180*f*
- Biphenyls, 125–127
 gene cluster (*bph*), 125, 126
- Bisphenol-A (BPA), 139
- Blooming water, bioluminescent control for, 95–99
 based on coupled enzyme system, 96*f*
 contamination of MES, 98

growth of microalgae biomass and, [97f](#)
 sensitivity of assays to contamination, [99](#)
 BLYAS, [138](#), [139](#)
 BLYES, [138](#), [139](#)
 BRET (bioluminescence resonance energy transfer) phenomenon, [192](#), [193f](#)
 BSA (bovine serum albumin), [29](#), [78](#)
 BTEX (benzene, toluene, ethylbenzene, and xylene), [115–120](#). *See also* BTEX bioreporters
 BTEX bioreporters, [115–117](#)
 firefly luciferase gene (*luc*), [118](#)
 isopropylbenzene (*ipb*) pathway, [119](#)
 for organic contaminants and representative environmental applications, [116–117t](#)
 P. putida TVA8 BTEX bioreporter, [119](#)
 T-2 toluene degradation (*tod*) pathway, [119](#)
 in toluene benzene utilization (*tbu*) pathway, [118](#)
 and TOL-encoded pathway, [118](#)
C
 Calanoids, [23](#), [25](#)
 Ca²⁺-regulated photoproteins, [182](#), [183](#), [184](#), [194](#)
 Camouflage, [12–13](#)
 Camp receptor protein (CRP), [49](#)
celB glycosidase, [164](#)
 Chemical-activated luciferase expression (CALUX) assay, [134](#), [135](#)
 H4IIE-luc cell line-based, [134](#), [135](#)
 variation in results, [135](#)
 Chemically synthesized substrates, [177](#)
 Chlorinated aliphatic hydrocarbons, [124–125](#)
 1,1,1 trichloroethane (TCA), [124](#)
 dichloromethane (DCM), [125](#)
 trichloroethylene (TCE), [124](#)
 Coastal ostracod (*Vargula hilgendorfii*), [7f](#)
 Coelenterazine, [4](#), [21](#), [23](#), [24–25](#), [180](#), [181](#), [182](#)
 biosynthesized by copepods, [26f](#)
 Competition-for-substrate method, [90](#)
 Competitive immunoassay, [177](#), [178f](#)
 Counter-illumination, [4](#), [12](#)
 hypothesis, [13](#)
 in marine mesopelagic dwellers, [13](#)
 CRISPR systems, [159](#)
 Cypridinid luciferin, [21–22](#), [22f](#)
 Cytochrome P450 gene CYP2D6 (CYP2D6*3), [191](#)
 Cytomegalovirus (CMV), assay of, [186](#)

D

Deep-sea benthos, [26–27](#)
 Dichlorodiphenyltrichloroethane (DDT), [113](#)
 Digoxigenin–antidigoxigenin conjugation, [184](#), [186](#), [189](#), [191](#)
 Digoxigenin-dUTP, [191](#)
 Dioxin-responsive elements (DREs), [130](#)
 bioluminescent reporter, [133](#)
 Direct chemical conjugation, [184](#)
 Direct reaction coupling, [90](#)
 DNA barcoding, [28](#)
 DNA hybridization assays, [186](#), [192](#)
 for *Plasmodium falciparum* (malaria), [186](#)
 DNA hybridization technologies, [157](#)
 DNA/RNA assays, [183](#)
 DTT (dithiothreitol), [78](#), [80f](#)
 Dual analyte single-well bioluminescence assay, [190](#), [190f](#)

E

Ecological monitoring, [70](#), [71](#), [76](#), [91](#), [105](#)
 EGFP (enhanced green fluorescent protein)-based fluorescent reporter strain, [124](#)
 ELISA (enzyme-linked immunosorbent assay), [192](#)
 Endocrine-disrupting chemicals (EDCs), [136](#)
 eukaryotic cell-based bioluminescent assays for, [131–132t](#)
 Endolysins, [156](#), [166](#)
Enterobacteriaceae, [39](#)
 Environmental toxicity, [86](#)
 Enzymatic bioluminescent bioassays. *See also* Bioluminescent assays
 designing new assays, [87–91](#)
 coupling enzymes, example of, [88f](#)
 enzyme interaction, [90](#)
 light emission process, [88](#)
 enzyme activity measurement
 NADH-dependent dehydrogenases, [89](#)
 proteolytic enzymes, [88f](#), [89](#)
 for pesticides, [99–103](#)
 influence of toxic substance on, [101t](#)
 pond water analysis, parameters of, [93f](#)
 principles of, [71–73](#)
 biotesting diagram, [73f](#)
 coupled enzyme system (*see* NADH:FMN-oxidoreductase-luciferase)
 exogenous compounds on, [72](#)
 luciferase biotesting, [72](#), [72f](#)
 modified diagram, [73f](#)
 single system of, [86–87](#)

for toxicological bioassay, 91–95, 99–103

Enzymes, 179

- auxiliary label enzyme, 180
- immobilized, 75, 76, 86
- incubating in test solution, 81–83
- of luminous bacteria, 69, 70, 72
- proteolytic, 89, 100
- test system, 87

Enzymolum, 76, 78*f*, 104

- design of, 78
- in toxicological bioassay of air, 85–86
- comparative characteristics of, 86*t*

Escherichia coli cells, 162, 167, 179

- DH5 α , 118
- fre*-like gene, 43
- serotype O157:H7, 163

Estrogen response elements (EREs), 136

Eukaryotic cell-based bioluminescent assays

- dioxin-like activities, 130*t*
- endocrine-disrupting activities, 131–132*t*

F

Felix-O1 reporter, 162, 163

Fireflies, 5, 8, 11, 16, 27, 179

Firefly luciferase, 75, 114, 118, 128, 142, 157, 179, 182, 189

- luc* genes, 118, 158, 165

Firefly luciferin, 4

Firefly mimics, 8

Firefly squid (*Watasenia scintillans*), 5, 6*f*, 21

Fishing-lines, 10

Flavin mononucleotide (FMN), 70, 79, 81, 81*f*, 98

- reduced FMN (FMNH₂), 42, 43, 51, 88, 88*f*, 158, 161

Fluoromycobacteriophages, 165

Fungus gnats (*Arachnocampa luminosa*), 10, 11, 11*f*, 15, 16

G

Gas chromatography coupled with mass spectrometry (GC/MS), 114

Glow-worms, 16

Gram-negative pathogens, 39

Gram-positive pathogens, 163

Green fluorescent protein (GFP), 157, 163, 165, 193

Groundwater monitoring

- diesel-oil-contaminated, 122
- TCE and 1,1,1 trichloroethane (TCA) contaminated, 124

H

Harvey, Newton, 27

HELN α bioreporters, 138

HELN β bioreporters, 138

High-performance liquid chromatography (HPLC), 113–114

Homogeneous immunoassay, 177, 192

Human HepG2-derived HG2L1.1c3 cell line, 133, 134

Human immunodeficiency virus

- Type 1 (HIV-1), 186

Human mannose-binding lectin (*MBL2*) gene, 191

Humanized bacterial luciferase, 142

Hydroxylated PCBs (OH-PCBs), 126

I

Ice nucleation protein, 157

Immobilization

- enzymes in bioluminescent assays
 - coupled enzyme system, 76
 - effective methods of, 76, 77*t*
- reagents
 - in benzoquinone, 80*f*, 82*f*
 - in CuSO₄, 80*f*, 81, 82*f*
 - enzyme-stabilizing additives in, 78
 - FMN in, 79, 80*f*
 - variation of luciferase, 79*f*
 - variation of NADH content, 79*f*, 80
 - varying the composition of, 78–79

Immunoassay, 177–179

- homogeneous, 177, 192
- luciferase as labels, 179–183

Immunocomplex detection system, 178

In vitro binding assays, 177

Information on 3D structure, 177

Integrated bioluminescent methods, 104. *See also* NADH:FMN-oxidoreductase-luciferase

Isopropylbenzene (*ipb*)

- pathway, 119, 120

J

JAK2 (Janus kinase 2) V617F somatic point mutation, 188, 189

Japanese firefly (*Luciola lateralis*), 18*f*

Japanese fireworm (*Odontosyllis*), 9, 10*f*

K

Krebs cycle, 115

L

- Laser-induced fluorescence, 184
- Light emission reaction
 - catalyzed by bacterial luciferase, 158
 - catalyzed by firefly luciferase, 158
- Light production. *See* Bioluminescence
- Light-emitting proteins, 176
- Light-producing reaction, 177
- Listeria*, 156, 161, 164
- Listeria* reporter phage A511::*luxAB*, 160, 161
- Listeriosis, 164
- Littoral zone, 19
 - luminous earthworms in, 20
- Locked nucleic acid (LNA), 188, 189
- Luciferase
 - biotesting, 72–73
 - scheme, 72f
 - biotin-streptavidin/avidin system, 180f
 - as immunoassay labels, 179–183
 - bacterial luciferases, 179
 - biotinylated luciferase, 180
 - firefly luciferases, 179
 - immunoprecipitation system (*see* Luciferase immunoprecipitation system (LIPS))
 - luciferase index (LI), 73
 - single-chain polypeptides, 181
 - homologous glycosylated, 182
- Luciferase immunoprecipitation system (LIPS), 181, 181f
- Luciferin, 4, 15, 21, 28, 118, 138, 179, 189
 - cypridinid luciferin, 21–22, 22f
 - substrate luciferin, 158
- Luminescent secretion clouds, 7
- Luminous species
 - brittle star (*Ophiopsila*), 9f
 - coastal fish (*Acropoma japonicum*), 24f
 - earthworm (*Microscolex phosphoreus*), 18f
 - freshwater snail (*Latia neritoides*), 19f
 - marine snail (*Angiola zonata*), 20f
 - millipede (*Paraspirobolus lucifugus*), 8, 8f
 - mushroom (*Mycena chlorophos*), 8, 9f
- lux* genes, 39, 41, 43–47
 - gene content and gene order of, 44f
 - luxCDABEG*, 39, 43
 - luxF*, 43
 - lux-rib* operons, 44, 45
 - genus-specific differences, 45
 - nonfluorescent flavoprotein coding, 43
- lux* operon expression, regulation of, 47–51
 - population density-responsive regulation, 47
- luxR_{Vh}* (*lux* operon in *V. harveyi*), 47

M

- Mann-Whitney U test, 83
 - MCF-7 cell line
 - β -Globin promoter, 136
 - derived MELN reporter cell line, 136
 - derived MVLN reporter cell line, 136
 - MDA-kb2 cell line, 137
 - bioreporter assay, 140
 - MELN assay, 136, 138, 140, 141
 - reporter cell line, 136, 138
 - Mercaptoethanol, 78, 80f
 - Microecosystem (MES) components, 91–95
 - experiments, 93
 - biomass of algae, 94f
 - dynamics of bioassay parameters for, 94f
 - Midwater hatchetfish (*Sternoptyx pseudobscura*), 13f
 - Misdirection (or distraction/obstruction of sight), 4
 - Mouse mammary tumor virus (MMTV) promoter, 137
 - Multianalytical bioluminescent assay, 189–192
 - Mushrooms, 8, 16
 - attraction of spore-dispersing organisms, 11
 - Mycobacterium* reporter phages, 162, 165, 166
 - Mycobacterium tuberculosis* (MTB) infection, 185
- N**
- N3-(oxohexanoyl-) L-homoserine lactone (OHHL), 163
 - NADH:FMN-oxidoreductase-luciferase, 70, 71–72, 76
 - advantages of, 75
 - in Lake Shira water analysis, 74
 - in pond water MES, 91–95
 - in single system of enzymatic assays, 87
 - Natural water biotesting
 - reference water, Lake Baikal, 84
 - samples
 - characteristics, 83, 84f
 - light emission intensity dependency on, 85f
 - toxicity bioassay of, 84f
 - toxicological characteristics of, 84f
 - using immobilized coupled enzyme system, 83–85
 - using soluble enzyme system, 83–85
 - Nucleic acid hybridization assay, 185f

- based on bioluminescent reporters, 183–185
 - with recombinant aequorin as reporter, 185*f*
- Nucleic acid sequence-based amplification (NASBA) assay, 186
 - for *Chlamidia pneumonia* RNA, 186
- O**
- Off-the-shelf digital cameras, 157
- Organic toxicant-induced health risks, 129
 - using eukaryotic cell-based bioluminescent assays
 - advantage of, 140
 - bioassays for dioxin and dioxin-like compounds, 130, 130*r*, 133–135
 - for endocrine-disrupting chemicals, 131–132*t*, 136
 - for environmental applications, 139–141
 - for hormonally active chemicals, 135–139
- Organophosphorous pesticides, 99, 100, 102
- P**
- PAH bioreporters, 120–122
 - environmental naphthalene detection, 121
 - P. fluorescens* HK44, 120, 121
 - phenanthrene degradative genes, 121
- Pepsinogen I assay, 189
- Pepsinogen II assay, 189
- Pesticides, toxicological bioassay of, 99–103
 - characteristics of toxic substances, 100*r*
 - inhibition of trypsin activity, 100
 - sensitivity of coupled enzyme, 99
 - triple enzyme systems, 99, 102*f*
- Phengodid beetle (*Rhagophthalmus ohbai*), 5, 6*f*, 7
- Phenol and derivatives, 127–128
- Photobacterium*, 39, 40*r*, 51, 52
 - colonies of *P. mandapamensis*, 39*f*
- Photorhabdus*, 39, 41*t*
- Polaroid films, 157
- Pollutants, 71, 74
 - anthropogenic pollutants studies, 89
 - on bioluminescence of soluble enzyme system, 82*t*
 - on immobilized coupled enzyme system, 82*t*
 - in Lake Shira water, 74
 - maximum permissible concentration (MPC), 76
 - toxicological bioassay with increased sensitivity to, 78–79
 - in Yenissei River water, 74
- Polychlorinated biphenyls (PCBs), 125–127, 130
- Polychlorinated dibenzofurans (PCDFs), 130
- Polychlorinated dibenzo-*p*-dioxins (PCDDs), 130
- Polycyclic aromatic hydrocarbon (PAHs), 120–122, 130. *See also* PAH bioreporters
- Polymerase chain reaction (PCR), 183, 184
 - PCR-based assays, 157
- Pond ecosystem, 91–95
 - bioluminescent bioassays for, 93*f*
 - biomass of algae, 94*f*
 - dynamics of, 92*f*
- Portable Laboratory for Toxicity Detection (PLTD), 104
- Primer extension (PEXT) reactions, 188, 191
- pRLuc42R bioreporter, 128
- Prophage, 156
- Prostatic acid phosphatase (PAP) with PSA, 189
- Proteolytic interactions, 90
- Protobioluminescence, 51
- Prototype biosensors, 105
- PSA with alpha-fetoprotein (AFP), 189
- Pseudomonas fluorescens* OS8(pDNdmpRlux), 127
- Pseudomonas oleovorans* alk operon genes, 122, 123
- Pseudomonas* strain, CF600, 127
- pTOLLUX plasmid, 115
- Pyrethroid, 99, 100
- Pyruvate phosphate dikinase, 180
- Q**
- Quantitative realtime PCR assay, 184
- Quorum sensing, 45, 47
 - in *A. fischeri*, 47
 - 3':5'-cyclic AMP (cAMP), 49
 - control of luminescence in, 50*f*
 - octanoyl-HSL (AI-2_{AF}), 50–51
 - regulatory factors in, 49–50
 - in *Vibrio campbellii* (*V. harveyi*), 47, 48
 - phosphorylation/dephosphorylation cascade, 49*f*
- R**
- Realtime PCR, 184, 189
- Receptor mutation, 159

Recombinant proteins, 176–177
 Reduced flavin mononucleotide (FMNH₂), 42, 43
 NADH:FMN oxidoreductase, 43
Renilla luciferases (RLuc), 180
 bioluminescence-based hybridization assay, 188f
 for miR21 in breast cancer cells, 187
 Reporter bacteriophages, 157
 advantage of, 161
 antibiotic resistance, 166
 in combination with other technologies, 166–167
 construction and use, 159–162
 drawbacks of, 161–162
 on foodborne pathogens, 167
 specific bioluminescent phages, 162–166
 on biosafety-relevant human pathogens, 165
 on clinically relevant pathogens, 165
 plant pathogenic bacteria, 166
 Restriction-modification systems, 159
 Reverse transcriptase-polymerase chain reaction (RT-PCR) assay, 184
 for *Mycobacterium tuberculosis* (MTB) infection, 185
 for prostate-specific antigen (PSA), 187
 (RS)-3-(3,5-dichlorophenyl)-5-methyl-5-vinylloxazolidine-2,4-dione (vinclozoline), 137
 Russian Federation standards, 83

S
 (S)-3-hydroxytridecan-4-one (cholerae autoinducer, CAI-1), 49
 Sakyo Kanda, 27
Salmonella, 162, 167
Salmonella reporter phage Felix O1, 161
 Sandwich-type immunoassay, 178f
Shewanella, 39, 40t, 41, 42, 46
Shewanellaceae, 39
Shigella, 167
 Single-tube luminometers, 161
 Split-protein reassembly technology, 192
 Spore-feeding fungus gnat (*Keroplatus nipponicus*), 11, 11f
 Startling (or deterring), 4
 click beetles, 6
 phengodid, 6
 Symbiotic luminescence, 23–24

T
 T-47D ER-CALUX, 137
 Terrestrial luminous organisms, 14
 Thyrotropin (TSH) assay, 183
 third-generation assay, 183
 Toll-like receptor 4 (TLR4) gene, 191
 Toluene benzene utilization (*tbu*) pathway, 118
 Toluene degradation (*tod*) pathway, 119
 Toxicity coefficient (TC), 73, 74
 and luciferase index (LI), 73, 74
 Toxicological bioassay
 pond ecosystem, 91–95
 signal system of enzymatic assays for pesticides, 99–103
 Trypsin, 69, 87, 90, 91, 95, 100, 102
 in microalgae biomass, 97f, 98
 in pond water analysis, 93f
 scheme of activity, 89f
 triple enzyme systems, 96, 99

U
 U2-OS cell line, 138

V
 Vertebrate endocrine system, 135
Vibrio, 39, 40t, 51
Vibrio harveyi luxAB genes, 122
Vibrionaceae, 38, 39

W
 Waste water biotesting
 reference water, Lake Baikal, 84
 samples
 characteristics, 83, 84t
 light emission intensity dependency on, 85f
 toxicity bioassay of, 84t
 toxicological characteristics of, 84t
 using immobilized coupled enzyme system, 83–85
 using soluble enzyme system, 83–85

X
 Xenobiotics, 71
 contamination of MES, 98
Xenopus Vitellogenin A2 gene, 136

Y

Yata Haneda, [27](#)

Yeast-based assays, [138](#), [139](#)

hAR/ARE-*luc* bioassay, [139](#)

Yellow fluorescent protein (YFP), [46](#), [192](#)

energy transfer (EYFP), [192](#)

Yenisei River water analysis, [74](#)

Yersinia pestis, [166](#)

Z

Zebrafish assay, [134](#), [137](#)