

# Index

## A

- Acanthamoeba* keratitis (AK), 194–195, 200
- Acanthamoeba* species, FLA
  - diagnosis, 198–199
  - genotypes, 196
  - history of, 194
  - keratitis, 194–195
  - life cycle, 194
  - in Malaysia and Thailand, 195
  - morphological characteristics, 190–192
  - in Singapore, 195
- Acute dermatolymphangioadenitis (ADLA), 43
  - candidiasis, 44, 45
  - prevention of, 47–48
  - treatment of, 47
- Acute filarial lymphangitis, 44
- Albendazole
  - B. malayi*, 46–47
  - STH infections, 218
- Amoebic liver abscess (ALA), *E. histolytica*
  - clinical symptoms and signs, 118
  - in Malaysia, 112, 113
  - treatment of, 123–124
  - in Vietnam, 111
- Ancylostoma*, STH infections, 208–209
- Anopheles*
  - An. baimaii*, 63
  - An. cracens*, 18
  - An. dirus*, 18, 62
  - An. epiroticus*, 62
  - An. hackeri*, 18
  - An. latens*, 18
  - An. maculatus*, 62
  - An. minimus*, 62
  - An. sundaicus* complex, 66
- Antigen card test, 41
- Anti-toxoplasma therapy, 167–168

- Apical membrane antigen 1 (AMA-1), 24
- Apoptosis, *Blastocystis* sp., 137–138
- Ascaris lumbricoides*, STH infections, 207
- ASEAN Economic Community (AEC), 2
- Association of Southeast Asian Nations (ASEAN), 1

## B

- Balamuthia mandrillaris*, FLA
  - description of, 197–198
  - diagnosis of, 198–199
  - morphological characteristics, 190–191, 193
- Binax NOW<sup>®</sup>, 41
- Blastocystis* sp.
  - amoeboid forms, 134–135
  - apoptosis, 137–138
  - cryopreservation, 146
  - cyst-like structures, 135–136
  - diagnosis of, 138–139
  - drug resistance, 141–142
  - fecal surveys, 139–140
  - genetic diversity, 142–143, 147
  - genome sequence of, 147, 148
  - granular forms, 133–134
  - life cycle, 136–137
  - molecular studies, 132
  - pathogenesis, 140–141
  - reclassification, 131, 132
  - reproduction modes, 137
  - subtype 3, 143–144
  - transmission, 147
    - human-to-human, 144
    - risk factors, 146
    - waterborne, 145–146
    - zoonotic, 144–145
  - vacuolar forms, 133

- BLF. *See* Brugian lymphatic filariasis (BLF)
- BmR1*, 40, 41
- Borneo island, *An. sundaicus*, 62
- Brugia*
- B. malayi* (*see* *Brugia malayi*)
  - B. pahangi*, 37
  - B. timori*
    - in Indonesia, 36
    - in Timor-Leste, 37
- Brugia malayi*
- brugian lymphatic filariasis
    - albendazole, 46–47
    - diethylcarbamazine, 45–46
    - ivermectin, 46
  - in Brunei Darussalam, 35
  - in Indonesia, 36
  - in Malaysia, 36
  - microfilaria of, 39, 40
  - periodic and subperiodic forms, 34
  - in Philippines, 36–37
  - in Thailand, 34
- Brugian lymphatic filariasis (BLF), 42
- acute dermatolymphangioadenitis, 43–44
  - acute filarial lymphangitis, 44
  - chronic manifestations, 44–45
  - pathogenesis, 43
  - treatment of
    - acute dermatolymphangioadenitis, 47–48
    - albendazole, 46–47
    - diethylcarbamazine, 45–46
    - ivermectin, 46
    - lymphoedema, 48
- Brunei Darussalam
- cryptosporidiosis, 87
  - filarial worms, 35
  - STH infections, 214–215
- C**
- Cambodia
- An. dirus*, 62
  - cryptosporidiosis, 86–87
  - E. histolytica*, 105–106
  - filarial worms, 35–36
  - Giardia*, 91, 93
  - P. knowlesi*, 16
  - schistosomiasis, 241
  - STH infections, 213
  - taeniasis, 235
- Cestode infections
- in Cambodia, 235
  - cysticercosis, 234–235
  - in Indonesia, 235–236
  - in Lao PDR, 236–237
  - in Malaysia, 237–238
  - in Myanmar, 238
  - in Philippines, 238
  - in Singapore, 238
  - taeniasis, 234–235
  - in Thailand, 238–239
  - in Vietnam, 239
- Clonorchiasis, 244
- Clonorchis sinensis*, 244–245
- Community-led total sanitation (CLTS), 219
- Cryopreservation, *Blastocystis* sp., 146
- Cryptosporidiosis
- in animals, 83–85
  - environmental contamination, 85–87
  - HIV/AIDS individuals
    - in Indonesia and Cambodia, 81
    - in Malaysia, 81
    - in Thailand, 80
    - in Vietnam, 81
  - non-HIV/AIDS individuals, 81–83
- Cryptosporidium*
- biology and life cycle, 78
  - clinical manifestation, 78–79
  - cryptosporidiosis (*see* Cryptosporidiosis)
  - diagnosis of, 79–80
  - infection of, 79
  - pathology of, 78–79
  - taxonomy of, 77–78
  - transmission of, 79
  - treatment of, 80
- Cysticercosis. *See* Taeniasis
- D**
- Diethylcarbamazine (DEC), 39, 45–46
- Dimethylsulfoxide (DMSO), 146
- Dirofilaria*
- D. immitis*, 37
  - D. repens*, 37
- Disability-adjusted life year (DALY)
- metric measurement, 206
- E**
- Elephantiasis, 44–45
- Entamoeba histolytica/dispar*
- axenic cultivation, 114–115
  - clinical symptoms and signs
    - amoebic liver abscess, 118
    - extraintestinal amoebiasis, 119
    - intestinal amoebiasis, 118

- diagnosis of
  - antibody detection tests, 120–121
  - antigen detection tests, 121
  - culture and isoenzyme analysis, 120
  - loop-mediated isothermal amplification, 123
  - microscopy, 119–120
  - PCR-based techniques, 122–123
- E. coli*, 124, 125
- E. hartmanni*, 124, 125
- E. moshkovskii*, 124, 125
- epidemiology of
  - in Cambodia, 105–106
  - in Indonesia, 109–110
  - in Laos People's Democratic Republic, 106
  - in Malaysia, 111–113
  - in Myanmar, 106–107
  - in Philippines, 108–109
  - in Singapore, 105
  - in Thailand, 107–108
  - in Vietnam, 110–111
- E. polecki*, 125
- history of, 104–105
- immunity, 117
- life cycle of, 115–116
- morphology, 113–114
- pathogenesis, 116–117
- prevalence of, 104
- prevention and control measures, 124
- transmission of, 113
- treatment
  - amoebic liver abscess, 123–124
  - intestinal amoebiasis, 123
- Enzyme-linked immunosorbent assay (ELISA)
  - E. histolytica*, 120–121
  - STH infections, 212
- F**
- Fasciola* spp., 246
- Fatal syndrome, 190
- Filarial worms
  - BLF
    - acute manifestations, 43–44
    - chronic manifestations, 44–45
    - pathogenesis, 43
    - treatment, 45–48
  - B. pahangi*, 37
  - D. immitis*, 37
  - D. repens*, 37
  - health and economic impacts, 51
  - LF
    - diagnosis of, 37–42
    - epidemiology, 34
    - global programme for, 48–50
  - SEA countries
    - Brunei Darussalam, 35
    - Cambodia, 35–36
    - Indonesia, 36
    - Lao PDR, 36
    - Malaysia, 36
    - Philippines, 36–37
    - Timor-Leste, 37
- Filariasis vectors
  - biological characteristics, 67–68
  - IRS, 69
  - ITN, 69
  - larval habitats, 66–67
  - parasite infection rates, 68–69
- Fluorescein isothiocyanate (FITC), 79, 88, 198
- Focus resources on effective school health (FRESH) framework, 224
- Food-borne trematodes
  - clonorchiasis, 244
  - Clonorchis sinensis*, 244–245
  - epidemiology, 243
  - Fasciola* spp., 246
  - Opisthorchis viverrini*, 245–246
  - paragonimiasis, 244
  - Paragonimus westermani*, 246–247
  - prevalence data of, 245
  - transmission, 243–244
- Free-living amoebae (FLA)
  - Acanthamoeba* species
    - diagnosis, 198–199
    - genotypes, 196
    - history of, 194
    - keratitis, 194–195
    - life cycle, 194
    - in Malaysia and Thailand, 195
    - morphological characteristics, 190–192
    - in Singapore, 195
  - Balamuthia mandrillaris*, 197–198
    - diagnosis, 198–199
    - morphological characteristics, 190–191, 193
  - clinical manifestations and diseases
    - Acanthamoeba* keratitis, 194, 200
    - granulomatous amoebic encephalitis, 191
    - primary amoebic meningoencephalitis, 191–193
  - diagnosis
    - cerebrospinal fluid, 198
    - indirect immunofluorescence, 198

- PCR, 198, 199  
 eye keratitis, 190  
 fatal syndrome, 190  
*Naegleria* species  
 diagnosis, 198–199  
 in Indonesia and Philippines, 197  
 morphological characteristics,  
 190–191, 193  
 PAM, 196–197  
 in Thailand, 196  
 in Vietnam and Malaysia, 197
- G**  
 Geographical information systems (GIS).  
*See* Soil-transmitted helminth  
 (STH) infections  
 Geohelminths. *See* Soil-transmitted helminth  
 (STH) infections  
*Giardia*  
 biology and life cycle, 87  
 clinical manifestation, 88  
 diagnosis, 88–89  
*G. duodenalis*, 89  
 giardiasis (*see* Giardiasis)  
 pathology, 88  
 taxonomy, 87  
 transmission, 88  
 treatment, 89  
 Giardiasis  
 in animals, 92–93  
 environmental contamination, 93  
 in humans  
 Cambodia and Lao PDR, 91  
 Indonesia, 89–90  
 Malaysia, 90  
 Philippines and Vietnam, 91  
 Thailand, 90–91  
 Global Programme for the Elimination of  
 Lymphatic Filariasis (GPELF), 38  
 Granulomatous amoebic encephalitis (GAE),  
 191, 195
- H**  
 HIV/AIDS  
 cryptosporidiosis  
 in Indonesia and Cambodia, 81  
 in Malaysia, 81  
 in Thailand, 80  
 in Vietnam, 81  
 toxoplasmosis, 166–167  
 Hookworm infections, STH, 208–209  
 Human-to-human transmission, *Blastocystis*  
 sp., 144  
 Hyperinfection syndrome, 210
- I**  
 Indirect haemagglutination antibody  
 (IHA) test, 109, 110  
 Indonesia  
*An. aconitus*, 62  
 cryptosporidiosis  
 HIV/AIDS individuals, 81  
 non-HIV/AIDS individuals, 82  
 cysticercosis, 235–236  
*E. histolytica*, 109–110  
 filarial worms, 36  
*Giardia*, 89–90  
*Naegleria* species, 197  
*P. knowlesi*, 15–16  
*Sarcocystis* spp., 181–182  
 schistosomiasis, 242  
 STH infections, 212  
 taeniasis, 235–236  
 toxoplasmosis, 160  
 Indoor residual spraying (IRS), 69  
 Insecticide-treated bednets (ITN), 69  
 Intestinal sarcocystosis. *See* *Sarcocystis* spp.  
 Iodoquinol, 141  
 Irritable bowel syndrome (IBS), 140, 141  
 Ivermectin, 46
- K**  
 Katayama fever, 240  
*kir* genes, 20
- L**  
 Laos People's Democratic Republic  
 (Lao PDR)  
*An. dirus*, 62  
*E. histolytica*, 106  
 filarial worms, 36  
*Giardia*, 91  
*Sarcocystis* spp., 182  
 schistosomiasis, 241  
 STH infections, 213–214  
 toxoplasmosis, 160  
*T. saginata*, 237  
*T. solium*, 237  
 LF. *See* Lymphatic filariasis (LF)  
 Loeffler's syndrome, 207

Loop-mediated isothermal amplification  
(LAMP)

*E. histolytica*, 123

PCR, 24

toxoplasmosis, 165

Lymphatic filariasis (LF)

diagnosis of, 37

antibody detection, 39–41

DNA detection, 42

microfilaria detection, 38–39

radiological diagnosis, 42

epidemiology, 34

global programme for

breakthroughs and developments, 49

elimination and progresses made,  
49–50

impacts, 50

population risk, 49

Lymphoedema, 44–45

**M**

*Macaca fascicularis*, 8–9

Malaria vectors

*Anopheles* vectors

*An. dirus* complex, 65

*An. maculatus* (group) complex,  
65–66

*An. minimus* complex, 65

*An. sundaicus* complex, 66

biological characteristics of, 62–64

larval habitats, 61–62

types, 58–60

IRS, 69

ITN, 69

WHO, 58

Malaysia

*Acanthamoeba* species, 195

*An. maculatus*, 64

*Blastocystis* sp., 143

cryptosporidiosis

in animals, 83–84

environmental contamination, 86

in HIV/AIDS individuals, 81

in non-HIV/AIDS individuals, 82

cysticercosis, 237

*E. histolytica*, 111–113

filarial worms, 36

*Giardia*

in animals, 92

environmental contamination, 93

in humans, 90

*Naegleria* species, 197

pentastomes, 185

*P. knowlesi*, 10–13

*Sarcocystis* spp., 179–180

schistosomiasis, 242

STH infections, 210–212

*T. gondii*, 158

toxoplasmosis, 160

*Mansonina*

*Ma. bonnae*, 67

*Ma. dives*, 67

*Ma. uniformis*, 67

Mass drug administration (MDA), 34

Mebendazole, 212, 221

Metronidazole

*Blastocystis* sp., 141, 142

*Giardia*, 89

Microfilaria detection, LF, 38–39

Muscular sarcocystosis. *See Sarcocystis* spp.

Myanmar

*An. dirus*, 62

*An. minimus*, 62

cryptosporidiosis, 83

cysticercosis, 238

*E. histolytica*, 106–107

*P. knowlesi*, 15

*Sarcocystis* spp., 182

STH infections, 214

*T. gondii*, 158

**N**

*Naegleria fowleri*. *See Naegleria* species, FLA

*Naegleria indonesiensis* strain NG945, 197

*Naegleria* species, FLA

diagnosis, 198–199

in Indonesia and Philippines, 197

morphological characteristics, 190–191, 193

PAM, 196–197

in Thailand, 196

in Vietnam and Malaysia, 197

*Necator*, 208–209

*Necator americanus*-*Ancylostoma*-secreted

protein-1 (Na-ASP-2) vaccine, 222

*Necator americanus*-aspartic protease-1

(Na-APR-1) vaccine, 222

*Necator americanus*-glutathione

S-transferase-1 (Na-GST-1)

vaccine, 222

Nested PCR assay

Pmk8 and Pmk9 primers, 22

*P. vivax*, 23

rRNA gene, 23

**O**

Og4C3-ELISA, 41  
*Opisthorchis viverrini*, 245–246

**P**

PanLF Rapid test, 41  
 Paragonimiasis, 244, 247  
*Paragonimus westermani*, 246–247  
 Peninsular Malaysia, *Sarcocystis* spp.,  
 180, 181  
 Pentastomes  
   Cephalobaenida, 184  
   in Malaysia, 185  
   morphology, 184–185  
   Porocephalida, 184  
   symptomatic and asymptomatic cases, 185

## Philippines

*Blastocystis* sp., 143  
 cryptosporidiosis, 82  
 cysticercosis, 238  
*E. histolytica*, 108–109  
 filarial worms, 36–37  
*Giardia*, 91  
*Naegleria* species, 197  
*P. knowlesi*, 13–14  
*Sarcocystis* spp., 182  
 schistosomiasis, 242–243  
 STH infections, 212  
 toxoplasmosis, 160

*Plasmodium*

*P. cynomolgi*, 7  
*P. falciparum*  
   erythrocytic cycle, 7  
   LAMP method, 24  
   pre-erythrocytic phases, 6  
*P. inui*, 7  
*P. knowlesi* (see *Plasmodium knowlesi*)  
*P. malariae*  
   erythrocytic cycle, 7  
   in Malaysia, 10  
   microscopy diagnosis, 20–21  
   pre-erythrocytic phase, 6  
   in Thailand, 13  
*P. ovale*, 7  
*P. vivax*  
   description, 5–6  
   erythrocytic cycle, 7  
   nested PCR assay, 23

*Plasmodium knowlesi*

epidemiology  
   in Cambodia, 16  
   in Indonesia, 15–16

in Malaysia, 10–13  
 in Myanmar, 15  
 in Philippines, 13–14  
 in Singapore, 14  
 in Thailand, 13  
 in Vietnam, 14–15  
 history and discovery, 8–10  
*kir* genes, 20  
 laboratory diagnosis of  
   lamp PCR, 24  
   microscopy, 20–22  
   nested PCR, 22–23  
   RDTs, 25  
   real-time PCR, 23–24  
 mosquito vectors of, 17–18  
 mtDNA dataset, 19  
 natural hosts

  long-tailed and pig-tailed macaques, 16  
 mtDNA and *csp* gene, 17  
*Semnopithecus obscurus*, 17  
 transmission and parasite life cycle  
   merozoite growth, 7  
   pre-erythrocytic phases, 6  
   relapsed infection, 7  
   sporozoites, 6  
   zoonotic simian malaria, 7–8

## PlasmoNex™, 23

## Polymerase chain reaction (PCR)

*E. histolytica*, 122–123  
 free-living amoebae, 198, 199  
 toxoplasmosis, 166

Primary amoebic meningoencephalitis (PAM),  
191–193, 196–197**R**

Rapid diagnostic tests (RDTs), 25  
 Real-time PCR assay, 23–24  
 Remote sensing (RS) satellite data, 223, 224

**S**

*Sarcocystis* spp.  
 diagnosis, 182–183  
 epidemiology  
   in Indonesia, 181–182  
   in Lao PDR, 182  
   in Malaysia, 179–180  
   muscular sarcocystosis, 178, 179  
   in Myanmar, 182  
   in Peninsular Malaysia, 180, 181  
   in Philippines, 182  
   in Singapore, 181

- in Thailand, 181
  - in Vietnam, 182
- life cycle, 178–179
- Miescher's tubules, 177
- morphology, 178, 179
- S. singaporensis*, 181, 182
- S. sulawesensis*, 181, 182
- Schistosomiasis
  - in Cambodia, 241
  - in Indonesia, 242
  - in Lao PDR, 241
  - in Malaysia, 242
  - in Philippines, 242–243
  - prevalence data of, 241
  - in Singapore, 243
  - symptoms
    - intestinal, 240
    - urinary, 240
  - in Thailand, 243–247
- SEA. *See* Southeast Asia (SEA)
- Singapore
  - Acanthamoeba* species, 195
  - Blastocystis* sp., 143
  - cryptosporidiosis, 87
  - cysticercosis, 238
  - E. histolytica*, 105
  - P. knowlesi*, 14
  - Sarcocystis* spp., 181
  - schistosomiasis, 243
  - STH infections, 214–215
  - taeniasis, 238
  - toxoplasmosis, 160
- Soil-transmitted helminth (STH) infections
  - aetiology
    - Ascaris* infections, 207
    - hookworm infections, 208–209
    - Strongyloides* infections, 209–210
    - Trichuris* infections, 207–208
  - DALY measurement, 206
  - epidemiology
    - in Brunei Darussalam, 214–215
    - in Cambodia, 213
    - in Indonesia, 212
    - in Lao PDR, 213–214
    - in Malaysia, 210–212
    - in Myanmar, 214
    - in Philippines, 212
    - in Singapore, 214–215
    - in Thailand, 210
    - in Timor-Leste, 214
    - in Vietnam, 212–213
  - geographical information systems
    - advantages, 223
    - FRESH framework, 224
    - helminth control programme, 222, 223
    - public and private sector participation, 224–225
    - remote sensing satellite data, 223, 224
  - neglected tropical diseases, 206
  - prevention and control
    - anthelmintic drug administration, 216–218
    - health education, 216, 219–220
    - national governmental control programmes, 217
    - sanitation, 216, 218–219
    - school-health-based approach, 220
    - transmission, 220
  - public health and economic consequences of, 215–216
  - vaccination, 221–222
- Southeast Asia (SEA)
  - AEC, 2
  - ASEAN, 1
  - BLF
    - acute manifestations, 43–44
    - chronic manifestations, 44–45
    - global programme, 48–50
    - pathogenesis, 43
    - treatment, 45–48
  - cestode infection, 234–239
  - Cryptosporidium* (*see Cryptosporidium*)
  - E. histolytica* (*see Entamoeba histolytica/dispar*)
  - FLA (*see* Free-living amoebae (FLA))
  - Giardia* (*see Giardia*)
  - lymphatic filariasis
    - antibody detection, 39–41
    - antigen detection, 41
    - DNA detection, 42
    - microfilaria detection, 38–39
    - radiological diagnosis, 42
  - P. knowlesi* (*see Plasmodium knowlesi*)
  - trematode infection, 239–247
  - vectors (*see* Filariasis vectors)
- STH infections. *See* Soil-transmitted helminth (STH) infections
- Strongyloides stercoralis*, 209–210
- Sumba island
  - An. subpictus*, 63
  - An. sundaicus*, 63
  - An. vagus*, 63

**T***Taenia*

- T. asiatica*, 234
- T. saginata*, 234
- T. solium*, 234

## Taeniasis

- prevalence data of, 235
- T. asiatica*, 234
- T. saginata*, 234
- T. solium*, 234

## Thailand

- Acanthamoeba* species, 195
- An. dirus*, 62
- Blastocystis* sp., 143
- cryptosporidiosis
  - in animals, 83
  - environmental contamination, 85
  - in HIV/AIDS individuals, 80
  - in non-HIV/AIDS individuals, 81–82
- cysticercosis, 238
- E. histolytica*, 107–108
- food-borne (*see* Food-borne trematodes)
- Giardia*
  - in animals, 92
  - in humans, 90

*Ma. bonnae*, 68

*Ma. uniformis*, 67, 68

*Naegleria* species, 196

*P. knowlesi*, 13

*Sarcocystis* spp., 181

STH infections, 210

toxoplasmosis, 160

## Threadworm infections, 209–210

## Timor-Leste

- B. timori*, 66
- cryptosporidiosis, 87
- filarial worms, 37
- STH infections, 214

## Tinidazole, 89

Tongue worms. *See* Pentastomes*Toxoplasma gondii*

- epidemiology of, 158
- genotypic analyses, 157, 158
- life cycle of, 156–157
- observation of, 156
- routes of, 157
- tachyzoites, 157
- tissue cysts, 156, 157
- toxoplasmosis (*see* Toxoplasmosis)

## Toxoplasmic encephalitis (TE), 167.

*See also* Toxoplasmosis

## Toxoplasmosis

- clinical impacts of

anti-toxoplasma therapy, 167–168

CT scan and MRI, 167

geographical distributions, 164–165

HIV/AIDS, 166–167

IgG and IgM antibodies, 165

laboratory investigation, 165

loop-mediated isothermal  
amplification, 165

PCR, 166

in pregnant women, 165–166

RT-PCR, 167

serological diagnosis, 166, 167

toxoplasmic encephalitis, 167

outbreaks of, 159

surveillance for

in Vietnam, 160–161

in Malaysia and Lao PDR, 160

in ocular and immunosuppressed  
patients, 163, 164

in Philippines and Singapore, 160

in pregnant women and children, 161

seroprevalence of, 162

in Thailand and Indonesia, 160

waterborne, 159

zoonotic, 158–159

## Transmission modes

*Blastocystis* sp.

human-to-human, 144

risk factors, 146

waterborne, 145–146

zoonotic, 144–145

*cryptosporidium*, 79

*giardia*, 88

## Trematode infections

*Schistosoma* spp.

*S. japonicum*, 299

*S. malayensis*, 299

*S. mekongi*, 299

schistosomiasis (*see* Schistosomiasis)

*Trichinella spiralis*, 207–208

*Trichuris trichiura*, 207–208

Triton X-100<sup>®</sup>, 39

**V**

## Vietnam

*An. dirus*, 62

cryptosporidiosis, 84–85

cysticercosis, 239

*E. histolytica*, 110–111

*Giardia*

in animals, 92–93

in humans, 91



*Naegleria* species, 197  
*P. knowlesi*, 14–15  
*Sarcocystis* spp., 182  
STH infections, 212–213  
*T. gondii*, 158  
toxoplasmosis, 160–161

**W**

Waterborne toxoplasmosis, 145–146, 159  
Wb-SXP1, 40

World Health Organisation (WHO), 58  
*Wuchereria bancrofti*  
in Indonesia, 36  
in Malaysia, 36  
in Philippines, 36  
in Thailand, 37

**Z**

Zoonotic toxoplasmosis, 144–145, 158–159  
Zymodemes, 116