

## INDEX

of names and of interesting anatomical structures described in this work.

- Absinthium**, 249.  
**Aconite**, 350.  
**Aconite root**, 350.  
**Aconitum Napellus**, 258, 350.  
**Acorus Calamus**, 400.  
**Aerenchyma**, 71, 199, 401, 404.  
**Alder Buckthorn bark**, 111.  
**Althaea officinalis**, 356.  
**Alyxia stellata**, 74.  
**Amphivasal vascular bundle**, 403.  
**Amygdalae amarae**, 422.  
**Amygdalae dulces**, 418.  
**Amylum Maidis**, 62.  
**Amylum Manihot**, 61.  
**Amylum Marantae**, 62.  
**Amylum Oryzae**, 63.  
**Amylum Tritici**, 64.  
**Anacyclus Pyrethrum**, 380.  
**Anatropous ovule** 133, 141, 153, 419.  
**Angostura bark**, 85.  
**Anise**, 206.  
**Anise fruit**, 206.  
**Anisum**, 206.  
**Anthemis nobilis**, 135.  
**Arctostaphylos Uva-Ursi**, 200.  
**Arnica**, 129.  
**Arnica montana**, 129, 396.  
**Artemisia Absinthium**, 249.  
**Artemisia Cina**, 142.  
**Articulation of filament of the Compositae**, 131, 140, 301.  
**Asparagine cells**, 358, 359.  
**Aspidium Filix mas**, 404.  
**Atropa Belladonna**, 269, 363.
- Balance hairs**, 251, 253, 254, 257.  
**Barosma betulina**, 157.  
**Bast fibres, bundles of** — in sec. phloem anastomosing in a tangential plane, 88, 376, 389.  
**Bast fibres, septate** —, 378.  
**Bast fibres with very slightly thickened walls**, 104.  
**Bast parenchyma cells, conjugated** —, 128.  
**Bast parenchyma cells, radial walls of** — pitted, 119.  
**Bast parenchyma elements in tangential layers**, 94.  
**Bast parenchyma forming the chief portion of the secondary phloem, e.g.**, 357, 362, 364.  
**Bearberry bark**, 115.  
**Bearberry leaves**, 200.
- Belladonna root**, 363.  
**Bitter Almond**, 422.  
**Bitter Orange peel**, 90.  
**Black Haw**, 125.  
**Black mass in fruits of Compositae, e.g.** 133.  
**Black Mustard seed**, 445.  
**Black Pepper**, 241.  
**Bordered pits, combined** — in wood, 341, 345.  
**Bordered pits with sieve structure**, 345.  
**Brassica alba**, 443.  
**Brassica nigra**, 445.  
**Buchbean**, 197.  
**Buchu**, 157.  
**Buchu leaves**, 157.  
**Bulbus Scillae**, 65.
- Cadmium iodide**, 219.  
**Calabar beans**, 438.  
**Calamus**, 400.  
**Cambium, interfascicular** — only forming secondary parenchyma, 409.  
**Campylotropous ovule**, 438, 444, 446, 450.  
**Cannabis indica**, 278.  
**Cannabis sativa**, 278.  
**Caraway fruit**, 210.  
**Carbenia benedicta**, 291.  
**Cardamom seeds**, 423.  
**Cardamomum**, 423.  
**Carony Bark**, 85.  
**Carrageen**, 67.  
**Carum Carvi**, 210.  
**Caryophylli**, 69.  
**Caryophyllum**, 69.  
**Cascara Sagrada**, 115.  
**Casparian dots**, 200.  
**Cassava starch**, 61.  
**Cassia augustifolia**, 187.  
**Cetraria**, 333.  
**Cetraria islandica**, 333.  
**Chamomile flowers**, 135.  
**Cherry-Laurel leaves**, 177.  
**Chondrus**, 67.  
**Chondrus crispus**, 67.  
**Cinnamomum bark**, 77.  
**Cinnamomum zeylanicum**, 77.  
**Citrullus Colocynthis**, 213.  
**Citrus vulgaris Risso**, 90.  
**Claviceps purpurea**, 417.  
**Cloves**, 69.

- Cluster crystal in a crystal skin of the same chemical constitution as the wall, 77, 83, 85, 95, 129.
- Cluster crystals in aleurone grains disappearing in potash, 220, 225, 237.
- Cluster crystals perforated, 108.
- Cnicus, 291.
- Coca, 161.
- Coca leaves, 161.
- Cochlearia Armoracia, 360.
- Colchicum autumnale 427.
- Colchicum seeds, 427.
- Collecting cells, 178, 192, 263, 287, 323.
- Collenchyma cells of stems in tangential rows, 117.
- Colocynth, 213.
- Colocynthis, 213.
- Colocynth pulp, 213.
- Colourless parenchyma, cells of — in tangential rows, 117.
- Combined bordered pits in wood, 341, 345.
- Companion cells, 88, 94, 104, 109, 128, 261, 265, 271, 275, 283, 294, 297, 313, 318, 321, 354, 357, 368, 395, 413.
- Condurango bark, 81.
- Conducting tissue in the ovary, 132.
- Conii folia, 311.
- Conium, 216.
- Conium fruit, 216.
- Conium maculatum, 216, 311.
- Conjugated bast parenchyma cells, 128.
- Conjugated wood parenchyma, 337.
- Connecting frames, 215, 404.
- Connecting threads, 431, 455.
- Coriander, 221.
- Cork, storied —. See storied cork.
- Corn starch, 62.
- Coriander fruit, 221.
- Coriandrum sativum, 221.
- Cork-patches in leaves, 171.
- Cortex Alyxii, 74.
- Cortex Angosturae, 85.
- Cortex Aurantii amari, 90.
- Cortex Cinnamomi, 77.
- Cortex Condurango, 81.
- Cortex Cuspariae, 85.
- Cortex Fructus Aurantii, 90.
- Cortex Granati, 92.
- Cortex Hamamelidis, 97.
- Cortex Mezerei, 102.
- Cortex of root, inner cell layers of — showing radial rows, 93, 367, 371, 381, 448.
- Cortex Pruni virginianae, 105.
- Cortex Quillaiae, 108.
- Cortex Rhamni Frangulae, 111.
- Cortex Rhamni Purshianae, 115.
- Cortex Syzygii, 121.
- Cortex Viburni prunifolii, 125.
- Crocus sativus, 456.
- Cryptoporous stomata, 215, 331.
- Crystal, branched —, 311.
- Crystal cells, walls of — thickened, 379.
- Crystal, cluster —. See cluster crystal.
- Crystal contained in walls of chlorenchyma cells, 331.
- Crystals in cells of unripe fruits, wanting in ripe fruits, 246, 302.
- Crystals in material treated with chromic acid 311.
- Crystals in sclereids, 246.
- Crystal sand cells, 193, 194, 245, 270, 271, 272, 273, 274, 276, 363, 364, 365, 451.
- Crystal-sheaths, 379. See also Endodermis.
- Crystal, simple — in a crystal skin of the same chemical constitution as the wall, 75, 76, 83, 91, 100, 102, 108, 111, 118, 120, 164, 165, 342, 346.
- Crystal, simple — perforated, 83, 102, 108.
- Crystal, simple — with a central cavity, 342.
- Crystals, simple — with augmenting size towards the outer part of the bark, 103.
- Crystal skin. See cluster crystal and simple crystal.
- Cubebs, 225.
- Cusparia bark, 85.
- Cusparia febrifuga, 85.
- Cynips Gallae tinctoriae, 246.
- Cystolith hairs, 279, 285, 286, 287, 289.
- Cytology, 11.
- Daphne Mezereum, 102.
- Datura Stramonium, 190, 449.
- Digitalis Folia, 165.
- Digitalis purpurea, 165.
- Dilatation. See tangential expansion of secondary phloem.
- Dill fruit, 203.
- Drawings, 5, 24.
- East Indian Senna, 187.
- Elder fruit, 245.
- Elettaria Cardamom, 423.
- Empirical description, 10.
- Emulsine, 179, 421, 422.
- Emulsine cells, 179.
- Emulsine-sheath. See also Endodermis.
- Endodermis containing emulsine, 421.
- Endodermis containing schizogenous glands, 367, 368.
- Endodermis developed as a crystal-sheath, 71, 189, 288.
- Endodermis developed as a emulsine-sheath, 179.
- Endodermis developed as a protective-sheath 200, 304, 353, 372 (root), 381, 388 (root), 392 (root), 395 (rhizome), 403 (monocotyledonous rhizome), 416 (monocotyledonous rhizome).
- Endodermis developed as a sclerenchyma-sheath 372 (root), 386 (root).
- Endodermis developed as a starch-sheath, 75, 79, 84, 95, 100, 103, 114, 118, 122, 149, 157, 160, 168, 172, 179, 183, 185, 193, 196, 199, 251, 255, 260, 265, 267, 268, 270, 274, 281, 288, 293, 297, 305, 313, 318, 320, 324, 327, 332.
- Endodermis with passage cells, 392, 395.
- Endostomium, 419.
- Epidermal cells containing crystals, 222, 226, 258, 298, 302.
- Epidermal cells containing starch grains, 78.
- Epidermal cells, mother cells of —, 205, 208, 210, 219, 223, 226, 231, 234.
- Epidermal cells of branches living during a long time, 78.

- Epidermal cells, outer walls of — showing cuticular layers containing crystal sand, 177, 178.  
 Epidermal cells showing pitted outer walls, 160, 219.  
 Epidermal cells showing thickened lignified walls, 208.  
 Epidermal cells with an outer wall consisting for a large part of mucilage, 433, 444, 447, 451.  
 Epidermal cells, walls of — showing crystals, 325, 331.  
 Epidermis consisting of fibres 143, 162, 246, 251, 256, 298, 299, 300, 310, 424.  
 Epidermis consisting of thick-walled cells, 298, 302, 308, 331.  
 Epithema, 317.  
 Ergot, 417.  
 Ergota, 417.  
 Ergot of Rye, 417.  
 Erythraea Centaurium, 303.  
 Erythroxyllum Coca, 161.  
 Eucalyptus Globulus, 169.  
 Eucalyptus leaves, 169.  
 Eugenia caryophyllata, 69.  
 Eugenia Jambolana, 121.  
 Exodermis, 371.  
 Exostomium, 419.
- Farina**, 64.  
**Fennel**, 230.  
**Fennel fruit**, 230.  
**Fibrous layer**, e.g. 277.  
**Fibrovascular bundle** 294 (stem *Carbenia benedicta*), 313 (stem *Conium maculatum*), 399 (*Rhizoma Arnicae*), 402 (*Rhizoma Calami*), 413 (*Rhizoma Hydrastis*), 416 (*Rhizoma Zingiberis*).  
**File hairs**. See hairs.  
**Filix Mas**, 404.  
**Flaxseed**, 432.  
**Flores Anthemidis**, 135.  
**Flores Arnicae**, 129.  
**Flores Chamomillae romanae**, 135.  
**Flores Chamomillae vulgaris**, 137.  
**Flores Cinae**, 142.  
**Flores Tiliae**, 146.  
**Foeniculum vulgare**, 230.  
**Folia Althaeae**, 153.  
**Folia Buchu**, 157.  
**Folia Coccae**, 161.  
**Folia Digitalis**, 165.  
**Folia Eucalypti**, 169.  
**Folia Jaborandi**, 173.  
**Folia Laurocerasi**, 177.  
**Folia Menthae piperitae**, 180.  
**Folia Menyanthidis**, 197.  
**Folia Salviae**, 183.  
**Folia Sennae**, 187.  
**Folia Stramonii**, 190.  
**Folia Taraxaci**, 194.  
**Folia Trifolii fibrini**, 197.  
**Folia Uvae Ursi**, 200.  
**Foxgloves leaves**, 165.  
**Fructus Anethi**, 203.  
**Fructus Anisi**, 206.  
**Fructus Carui**, 210.  
**Fructus Colocynthis**, 213.  
**Fructus Conii**, 216.  
**Fructus Coriandri**, 221.  
**Fructus Cubebae**, 225.  
**Fructus Foeniculi**, 230.  
**Fructus Pimentae**, 236.  
**Fructus Piperis nigri**, 241.  
**Fructus Sambuci recentes**, 245.
- Gallae**, 246.  
**Galls**, 246.  
**Gigartina mamillosa**, 67.  
**Ginger**, 414.  
**Glandulae Lupuli**, 248.  
**Glandular hairs in intercellular spaces**, 406.  
**Glandular hairs, unicellular** —, 406.  
**Glycyrrhiza glabra**, 376.  
**Glycyrrhiza glabra var. glandulifera**, 376.  
**Glands, oblito-schizogenous** —, 70, 72, 73, 170, 171, 237, 240.  
**Glands, oblito-schizogenous** — with disk, 170, 171.  
**Glands, schizogenous** — 138, 139, 141, 148, 212, 218, 251, 253, 255, 256, 258, 293, 297, 299, 302, 313, 314, 318, 332, 367, 368, 369, 381, 383, 384, 397.  
**Glands, schizogenous** — with septa, 212.  
**Glands, schizo-lysigenous** — 91, 159, 174.  
**Glands with a perforated cover**, 236, 240.  
**Glossary**, 21.  
**Guaiacum officinale**, 335.  
**Guaiacum sanctum**, 335.  
**Guaiacum wood**, 335.  
**Guiding schemes**, 9, 12, 21, 25.  
**Gum pockets in medullary rays**, 108.
- Habit**, 3.  
**Hairs, file** —, 173, 175, 184, 187, 188, 191, 201, 202, 207.  
**Hairs not separated by a transverse partition wall from the cells bearing them**, 207.  
**Hairs, sclerotic** —, 207.  
**Hairs, stellate** —, 98, 151, 153, 154, 155.  
**Hairs, twin** —, 132.  
**Hamamelis Bark**, 97.  
**Hamamelis virginiana**, 97.  
**Henbane**, 319.  
**Herba Absinthii**, 249.  
**Herba Aconiti recens**, 258.  
**Herba Belladonnae recens**, 269.  
**Herba Cannabis indicae**, 278.  
**Herba Cardui Benedicti**, 291.  
**Herba Centaurii**, 303.  
**Herba Centaurii minoris**, 303.  
**Herba Conii**, 311.  
**Herba Hyoscyami**, 319.  
**Herba Sabiniae**, 328.  
**Histology**, 11.  
**Horseradish root**, 360.  
**Hour-glass-shaped cells**, 439.  
**Humulus Lupulus**, 248.  
**Hydrastis canadensis**, 408.  
**Hyoscyamus niger**, 319.

- Iceland Moss, 333.  
 Idioblasts forming longitudinal rows, 85, 271, 272, 274, 281, 283, 284, 287, 289, 340, 345, 358, 364, 371, 373, 386, 412.  
 Indian Hemp, 278.  
 Intercellular spaces, walls adjoining — showing in iodine and sulphuric acid 66 per cent. a thin outermost layer remaining colourless, 199.  
 Intercellular spaces, walls surrounding the — covered with a very thin lamella becoming brown in iodine and sulphuric acid 66 per cent. or covered with a cuticle, 397, 407 (cuticle).  
 Inula Helenium, 365.  
 Inulin 383, 384, 389, 390, 398, 399, 400.  
 Ipomoea Purga, 370.  
 Irish Moss, 67.  
  
 Jaborandi Leaves, 173.  
 Jalapa, 370.  
 Juniperus Sabina, 328.  
  
 Latex tubes, unbranched —, 283, 288.  
 Latex vessels, 195, 389.  
 Laticiferous tubes, 75, 76.  
 Laticiferous tubes, non-articulated — 83, 84.  
 Leaf traces, 329 (Juniperus Sabina).  
 Leaf-sheath with secondary cork tissue, 394.  
 Lenticels with closing bands, 83, 94, 99, 113, 116, 126, 377.  
 Lenticels, Collenchyma wanting under —, 83, 99, 117.  
 Libriform fibres, partitioned —, 271.  
 Lichen islandicus, 333.  
 „Lichtlinie“, 439.  
 Lignum Guaiaci, 335.  
 Lignum Quassiae, 338.  
 Lignum Santalinum, 342.  
 Lime-tree flowers, 146.  
 Linden flowers, 146.  
 Linnean method, 6.  
 Linnean method applied to Micrography 9.  
 Linseed, 432.  
 Linum usitatissimum, 432.  
 Liquorice root, 376.  
 Literature, 19, 48.  
 Lupulinum, 248.  
 Lycopodium, 346.  
 Lycopodium clavatum, 346.  
  
 Macroscopic characters, Description of —, 18.  
 Maize Starch, 62.  
 Male Fern, 404.  
 Manihot utilisissima, 61.  
 Maranta Arrowroot, 62.  
 Maranta arundinacea, 62.  
 Maranta starch, 62.  
 Margin of the blade of the leaf, 148, 159, 171, 188, 199, 201, 254, 256, 263, 317.  
 Marsdenia, 81.  
 Marshmallow leaves, 153.  
 Material, 19.  
 Matricaria Chamomilla, 137.  
 Medulla containing phloem bundles, 272, 275, 322, 325.  
 Medullary rays, compound — in secondary phloem, 124, 128.  
 Medullary rays, incipient — in wood, 341.  
 Medullary rays in secondary phloem in two kinds, 128.  
 Medullary rays in secondary phloem with clusters of sclereids, 85, 111, 129.  
 Medullary rays in secondary xylem broken off in a radial direction by a group of vessels, 339.  
 Mentha piperita, 180.  
 Menyanthes trifoliata, 197.  
 Metacribral bands of bast parenchyma, 77, 101, 104, 114, 119.  
 Metatracheal wood parenchyma, 340, 344.  
 Method of pen-portraits, 2.  
 Mezereum, 102.  
 Mezereon bark, 102.  
 Micrography, 6, 20.  
 Micrography of powder, 17, 73, 80, 91, 97, 120, 169, 190, 194, 230, 232, 336, 342, 346, 356, 359, 369, 376, 380, 396, 404, 427, 449, 455.  
 Microscopical Anatomy, 11.  
 Mucilage cells, 79, 80, 87, 88, 113, 117, 118, 149, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 162, 358, 359, 433, 444, 447, 451.  
 Mucilage cells larger than the surrounding cells, 79, 89.  
 Myristica fragrans, 434.  
 Myrosine cells, 361, 448.  
 Myrosine fibres, 362.  
  
 Nutgall, 246.  
 Nutmeg, 434.  
 Nux Vomica, 452.  
  
 Oblito-schizogenous glands. See glands.  
 Oil cells, 79, 80, 226, 227, 228, 230, 242, 244, 402, 415, 416, 425, 426, 428, 436.  
 Oil cells in groups, 242.  
 Oil cells having the same size and shape as the surrounding parenchyma cells, 79.  
 Operculum of seed, 426.  
 Oryza sativa, 63.  
 Ovule, anatropous —, 133, 141, 153, 419.  
 Ovule, campylotropous — 438, 444, 446, 450.  
 Ovule inserted in the top of the ovary, 419.  
 Ovule, semi-anatropous —, 290, 419, 428.  
  
 Palisade chlorenchyma, cells of — half as high as the others and with the other dimensions often somewhat larger, 159, 174.  
 Palisade chlorenchyma continued in the midrib of a leaf, 189, 318.  
 Palisade chlorenchyma present in 2 or 3 layers of cells 177, 185, 195, 198, 201, 306, 331.  
 Palisade chlorenchyma present on either side of the blade 171, 188, 254, 257, 296.  
 Palisade sclerenchyma 439, 440.  
 Panama bark, 108.  
 Papaver Rhoeas, 347.  
 Paratracheal wood parenchyma, 340, 344.  
 Partitioned sclerenchyma fibres of pericycle 271.  
 Passage cells in endodermis, 392, 395.

- Pen-portraits, 2, 13.  
 Pen-portraying method, 2.  
 Pepper, 241.  
 Peppermint, 180.  
 Pericycle, bundles of sclerenchyma fibres in — corresponding in place to a vascular bundle, 260.  
 Pericycle consisting of sclerenchyma fibres, sclereids and common parenchyma, 100.  
 Pericycle, on transverse section especially the cells of the outer layer of — in radial rows with the endodermal cells, 407.  
 Pericycle showing a tangential reticulum of isolated sclerenchyma fibres and bundles of these, 84.  
 Pericycle showing more sclerenchyma fibres in elder branches, 329.  
 Pericycle with partitioned sclerenchyma fibres, 271.  
 Pericyclic fibres, partitioned —, 271.  
 Pericyclic fibres with walls showing cellulose reaction 76, 114, 118, 186, 271, 282, 288, 324.  
 Periderm cells lying in transverse (horizontal) rows, 357.  
 Periderm cells often pretty regularly arranged in tangential rows, 367.  
 Periderm cells with lignified walls, 99, 103, 357.  
 Periderm cells with thickened walls, 93, 98.  
 Perisperm, 244, 426, 435.  
 Perisperm, secondary —, 436.  
 Perspective description, 10.  
 Petala Rhoeados, 347.  
 Petala Rosae gallicae, 348.  
 Petiole, 319.  
 Peucedanum graveolens, 203.  
 Phellem of secondary cork tissue in 2 kinds, 86.  
 Phellem with sclereids, 383.  
 Phelloderm 79, 83, 99, 106, 361, 363, 377, 389, 395, 409.  
 Phelloderm cells with tangentially directed raphides or simple crystals, 87.  
 Phelloderm with sclereids, 87, 361.  
 Phloem bundles in medulla, 272, 275, 322, 325.  
 Phloem forming a continuous ring in a stem, 114, 271.  
 Phloem, quaternary —, 375.  
 Phloem, tertiary —, 374.  
 Pimpinella Anisum, 206.  
 Picraena excelsa, 338.  
 Pictures, 5, 24.  
 Pilocarpus Jaborandi, 173.  
 Physostigma venenosum, 438.  
 Phytography, 6.  
 Pimenta officinalis, 236.  
 Pimento, 236.  
 Piper, 241.  
 Piper Cubeba, 225.  
 Piper nigrum, 241.  
 Podophyllum peltatum, 411.  
 Poelasari, 74.  
 Poelasari dedès, 74.  
 Pollen Lycopodii, 346.  
 Pome granate bark, 92.  
 Portraying descriptions, 13.  
 Potato flour, 64.  
 Potato starch, 74.  
 Primary cork, 93, 391.  
 Primary phloem in a stem in the shape of a continuous layer, 114.  
 Protective-sheath. See endodermis.  
 Protoxylem, 284.  
 Prunus Amygdalus var. amara, 422.  
 Prunus Amygdalus var. dulcis, 418.  
 Prunus Laurocerasus, 177.  
 Prunus, serotina, 105.  
 Pterocarpus santalinus, 342.  
 Pulpa colocynthidis, 213.  
 Pulvis Lycopodii, 346.  
 Punctum Vegetationis, periodical development of —, 261, 313.  
 Punica Granatum, 92.  
 Pyrethrum, 380.  
 Quassia wood, 338.  
 Quaternary phloem, 375.  
 Quaternary xylem, 375.  
 Quercus infectoria, 246.  
 Quercus lusitania, 246.  
 Quillaia Saponaria, 108.  
 Quillaia bark, 108.  
 Radix Aconiti, 350.  
 Radix Althaeae, 356.  
 Radix Armoraciae, 360.  
 Radix Belladonnae, 363.  
 Radix Helenii, 365.  
 Radix Jalapae, 370.  
 Radix Liquiritiae, 376.  
 Radix Pyrethri, 380.  
 Radix Sarsaparillae, 385.  
 Radix Taraxaci, 387.  
 Radix Valerianae, 390.  
 Ramenta, 405.  
 Raphide cells, 66, 87, 89, 386, 387.  
 Raphide cells somewhat larger than the surrounding common bast parenchyma cells, 89.  
 Raphide cells with radially directed raphides in medullary rays, 89.  
 Rasura Ligni Santalini rubri, 342.  
 Reagents, 19, 46.  
 Red-Poppy petals, 347.  
 Red-rose, 348.  
 Red-rose petals, 348.  
 Red Sandal-wood, 342.  
 Red Sanders wood, 342.  
 Resin canals, schizogenous —, 381.  
 Resin cells, 373.  
 Reticulate cells, 205, 218, 233.  
 Rhamnus Frangula, 111.  
 Rhamnus Purshiana, 115.  
 Rhizoma Arnicae, 396.  
 Rhizoma Aspidii, 404.  
 Rhizoma Calami, 400.  
 Rhizoma Filicis, 404.  
 Rhizoma Filicis maris, 404.  
 Rhizoma Hydrastis, 408.  
 Rhizoma Podophylli, 411.  
 Rhizoma Zingiberis, 414.

- Rhytidoma, 109.  
 Rhytidoma, annular —, 86, 122.  
 Rhytidoma, scaly —, 87, 93, 106, 122, 377, 381.  
 Rice starch, 63.  
 Roots of *Valeriana officinalis*, vertical —, 392, 393.  
 Roots of *Valeriana officinalis*, horizontal —, 392, 393.  
 Root with radially arranged inner cortical cell layers, 93, 367, 371, 381, 448.  
*Rosa gallica*, 348.
- Sabina**, 328.  
 Sacred Bark, 115.  
 Saffron, 456.  
 Sage leaves, 183.  
*Salvia* leaves, 183.  
*Salvia officinalis*, 183.  
*Sambucus nigra*, 245.  
*Santonica*, 142.  
*Sarsae radix*, 385.  
*Sarsaparilla*, 385.  
*Savin*, 328.  
 Schizogenous duct in medullary rays of secondary phloem, 129.  
 Schizogenous glands. See glands.  
 Schizo-lysigenous glands. See glands.  
 Sclereids, clusters of — in secondary phloem showing a fibrous shape, 107, 128.  
 Sclereids in phellem 283.  
 Sclereids in phelloderm, 87, 361.  
 Sclereids with thin outer or with thin outer and inner walls, 79, 87, 122, 356.  
 Sclerenchyma fibres, partitioned — of pericycle, 271.  
 Sclerenchyma fibres with branched ends, 246.  
 Sclerotic hairs. See hairs.  
*Sclerotium Clavicipitis purpureae*, 417.  
*Secale cornutum*, 417.  
 Secondary cork tissue appearing in epidermis, 75, 81, 82, 86, 103, 125.  
 Secondary cork tissue appearing in inner layers of primary cortex, 367 (root).  
 Secondary cork tissue appearing in outmost or in second or in third layer of primary cortex, 98, 105, 106, 109, 113, 116, 121, 371 (root), 388 (root), 394 (rhizome), 397 (rhizome), 401 (rhizome), 409 (rhizome), 412 (rhizome).  
 Secondary cork tissue appearing in pericycle 93 (root), 95 (stem), 357 (root), 360 (root), 363 (root), 377 (root), 388 (root).  
 Secondary cork tissue appearing in trichomes of epidermis, 103  
 Secondary cork tissue in leaf-scales of rhizomes, 394, 409.  
 Secondary cork tissue in scales, 78.  
 Secondary cork tissue, monopleuric —, 412.  
 Secondary cork with cells in tangential layers, 93.  
 Secondary perisperm, 436.  
 Secondary phloem consisting for the largest part of bast fibres, 110.  
 Secondary phloem consisting for the greater part of tissues belonging to the parenchymatic system, e.g. 107, 123.  
 Secondary phloem consisting of elements of the 3 systems e.g. 376, 378.  
 Secondary phloem only consisting of the cribral and the parenchymatic system 84, 94, 354, 357, 361, 364, 368, 373, 383, 389.  
 Secondary phloem showing tangential layers 88, 101, 109, 114, 118, 119, 120, 123.  
 Secondary phloem with clefts on both sides of medullary rays 107.  
 Secondary phloem with storied arrangement, 361.  
 Secondary xylem only consisting of elements of the tracheal and the parenchymatic system 355, 362, 373, 390.  
 Secondary xylem with storied arrangement, 336, 339, 343, 362.  
 Seed containing perisperm, 225, 229.  
 Seed with 2 integuments, 424, 432, 444, 446.  
 Semen *Amygdalae amarae*, 422.  
 Semen *Amygdalae dulcis*, 418.  
 Semen *Cardamomi*, 423.  
 Semen *Colchici*, 427.  
 Semen *Lini*, 432.  
 Semen *Lycopodii*, 346.  
 Semen *Myristicae*, 434.  
 Semen *Physostigmatis*, 438.  
 Semen *Sinapis*, 445.  
 Semen *Sinapis albae*, 443.  
 Semen *Sinapis nigrae*, 445.  
 Semen *Stramonii*, 449.  
 Semen *Strychni*, 452.  
 Semi-anatropous ovule, 290, 419, 428.  
 Septate bast fibres, 378.  
 Sieve-plates on lateral walls of sieve-tubes, 110, 114, 119, 123, 407.  
 Sieve-tubes united in anastomosing bundles, 361.  
 Simple crystal. See crystal.  
*Smilax*, 385.  
 Soap bark, 108.  
*Solanum tuberosum*, 64.  
 Spiral cells, 309, 403.  
 Spiral fibres, 428.  
 Spongy chlorenchyma in the midrib of a leaf, 318.  
 Sporae *Lycopodii*, 346.  
 Squill, 65.  
 Starch-sheath. See endodermis.  
 Starch-sheath in stem, cell walls of — lignified, 251, 260.  
 Stellate hairs. See hairs.  
 Statistical data, 22; 62, 64, 67, 80 (anatomical characters); 129, 137, 147, 423 (macroscopic characters).  
 Stellate hairs. See hairs.  
 Stigmata *Croci*, 456.  
 Stomata, compressed —, 263.  
 Stomata, cryptoporous —, 215, 331.  
 Stomata missing one of the guard-cells, 191.  
 Storied arrangement in secondary phloem, 361.  
 Storied arrangement in secondary xylem, 336, 339, 343, 362.  
 Storied cork, 109, 394 (rhizome), 397 (rhizome), 401 (rhizome), 415 (rhizome).  
 Striae, cell walls showing 2 intercrossing sets of —, 390, 409.

- Striation of cell wall, 358.  
 Stramonium leaves, 190.  
 Stramonium seeds, 449.  
 Strychnos Nux-vomica, 452.  
 Stylopode, 213, 220.  
 Suberized cell walls with a thin inner cellulose layer, e.g., 415.  
 Substitute fibres in secondary phloem, 354, 362 (forming the bulk of the bast parenchyma), 383, 389.  
 Substitute fibres in secondary xylem, 337, 344, 362 (forming the bulk of the wood parenchyma), 384 and 390 (forming the bulk of the wood parenchyma and of medullary rays).  
 Summitates Absinthii, 249.  
 Syzygium bark, 121.  
 Sweet flag root, 400.  
 Sweet almond, 418.  
 Tangential expansion of secondary phloem (dilatation), 87, 88, 94, 96, 106, 109, 123, 127, 351, 378, 383.  
 Tannin cells, 402, 437.  
 Tapioca starch, 61.  
 Taraxacum leaves, 194.  
 Taraxacum officinale, 194, 387.  
 Tertiary layer, 271, 455.  
 Tertiary phloem, 374.  
 Tertiary xylem, 374.  
 Thecae, line of dehiscence of —, 309, 327.  
 Thecae, septa of — 277.  
 Thistle, 291.  
 Thyloses, 359.  
 Tilia cordata, 146.  
 Tilia platyphyllos, 146.  
 Tinnivelly Senna, 187.  
 Topography, 12.  
 Tracheids in seed-coat, 440.  
 Trichomes, walls of — showing crystals, 325.  
 Triticum vulgare, 64.  
 Twin crystals, 193, 323, 327.  
 Twin hairs, 132.  
 Urginia Scilla, 65.  
 Vacuoles of a cell not all filled with the same sap, 130, 131, 132, 134, 139, 141, 149, 152, 166, 167, 191, 192, 198, 199, 253, 255, 257, 266, 293, 294, 295, 296, 348, 405, 406, 407, 430.  
 Valeriana officinalis, 390.  
 Valerian rhizome, 390.  
 Valerian root, 390.  
 Vascular bundle, amphivasal —, 403.  
 Vessels of xylem with walls showing no reaction with phloroglucin and hydrochloric acid, neither with anilin sulphate, 416.  
 Vessels united in anastomosing bundles, 362.  
 Vessel tracheids, 408.  
 Viburnum prunifolium, 125.  
 Virginian Prune bark, 105.  
 Vittae, anastomosing —, 207.  
 Vittae, bifurcating —, 207.  
 Vittae without transverse septa, 224.  
 Water-pores, 177, 262, 315, 322.  
 Wheat starch, 64.  
 White Mustard seed, 443.  
 Wild Cherry, 105.  
 Witch Hazel bark, 97.  
 Wood parenchyma, conjugated —, 337.  
 Wood parenchyma forming the main part of the secondary xylem, 359, 362, 365.  
 Wood parenchyma, metatracheal —, 340, 344.  
 Wood parenchyma, paratracheal —, 340, 344.  
 Wood parenchyma scattered among the libriform fibres, 337.  
 Xylem forming a continuous ring in a stem, 271.  
 Xylem, quaternary, 375.  
 Xylem, tertiary, 374.  
 Zea Mays, 62.  
 Zingiber officinale, 414.  
 Zingiberis, 414.