

Index

A

- ABTS decolorization assay, 201–202, 243
- ACE. *See* Angiotensin I-converting enzyme (ACE)
- Agua de chia*, 131
- Água-mel, 349
- Almond, 295, 297–298
- Amaranth, 192, 195
 - characteristics, 190–191
 - cultivars, 190
 - health benefits, 189
 - low commercial value, 189
 - nutraceutical compounds
 - fatty acids, 192
 - protein content, 192, 195
 - squalene, 192, 195
 - origin, 190–191
 - production yield of, 190
 - species, 190
- Amaranthus hypochondriacus* (*see* *Amaranthus*)
- Añapa, 108
- Andean grain, 119, 124
- Angiotensin I-converting enzyme (ACE), 201
- Antioxidants, 242
 - bioactive compounds, 6
 - Capsicum chinense* Jacq. genotypes (*see* *Capsicum chinense* Jacq. genotypes)
 - carotenoids, 242
 - chain-breaking antioxidants, 4
 - CMG, 281
 - consumers awareness, 4
 - consumption, 3
 - definition, 3
 - functional food, 4
 - honey, 331–333, 340–341
 - lipid antioxidants, 3
 - lipid peroxidation inhibitors, 241
 - natural, 4–6
 - phenolics, 242
 - polyphenols, 5
 - potential, 5
 - preventive antioxidants, 4
 - Salvia hispanica*, 194, 204–205
 - sources, 5–6
 - synthetic, 4
 - vitamin C, 242
- Argan oil
 - argan forest, 310–313
 - argan tree, 310
 - beauty, 313
 - chemical composition, 316–317
 - cosmetic, 313
 - edible, 313
 - industrial extraction method, 316
 - Morocco, 309–310
 - pharmacological properties of, 317–318
 - physical properties of, 316–317
 - semi-mechanized extraction method, 314–316
 - traditional extraction method, 313–314
 - types, 313
- Argentinean Food Codex, 108
- Arrope, 108
- Available starch (AS), 149–151
- Ayran, 90–92
- 2,2'-Azino-bis(3-ethylbenzthiazoline-6-sulfonic acid) (ABTS) assay, 201

B

- Bacillus* spp., 43–44
B. subtilis, 57
 Bacteriocins, 172, 177–178
 food preservation and human therapy, 178
 genus *Lactobacillus*, 179
 production and modeling, 179–181
 Bakmaz (folk medicine), 216
 Berbers, 311
 Betalains, 258
 Bifidobacteria, 11
 Bioactive compounds, 124–125, 339
 Biochemical characterisation, 212–215
 Black mulberry (*Morus nigra* L.)
 acid strawberry/tree strawberry, 210
 biochemical characterisation, 212–215
 botanical characterisation, 210–211
 full maturity, 210
 phytotherapeutic characterisation, 215–216
 phytotherapeutic purposes, 210
 plant parts utilised economically, 211–212
 Bryndza, 75, 80
 for human health, 84–85
 history, 81–82
 production technology, 82–83
 Bulgur, 95–97

C

- Cactus products, 257–258
 anatomy and morphology, 251
 cladodes
 chemical composition of, 254
 mucilage, 254–257
 nopal flour, 257–258
 nopalitos, 254
 spines, 254
 flowers, 260
 fruits, 258–259
 grow up, 251
 medicinal properties, 260–264
 native, 251
 natural bioactive substances, 253
 parts, 253
 physiological traits and distinctive
 appearances, 251
 phytochemicals, 253
 pre-Columbian culture, 251
 seeds and seed oil, 259–260
 Caffeine, 359
Camellia sinensis (see Tea)
 Cañahua
 agronomic classifications, 119
 cereal-like seed, 120

- characteristics, 120
 chemical composition
 bioactive compounds, 124–125
 carbohydrates, 122
 dietary fiber, 123
 lipids, 123–124
 minerals, 124
 proteins, 121
 starch, 122–123
 erect types, 120
 high Andes area, 119
 in global food, potential uses, 125–128
 in local food, traditional uses, 125
 semierect types, 120
 Capsaicinoids, 290–291
Capsicum chinense
 antimicrobial properties, 291
 antioxidant capacity, 291
 applications, 289
 capsaicinoids, 290–291
 nutrient composition, 289–290
 pungency, 290–291
 ripe fruits, 289
 unripe fruits, 289
 Yucatan Peninsula, 289
Capsicum chinense Jacq. genotypes, 242–247
 extracts
 ABTS decolorization assay, 243, 246
 ascorbic acid content, 243, 245
 carotenoid content, 242–245
 β -carotene decolorization assay,
 243–244, 246–247
 statistical analysis, 244
 total carotenoids, 243
 total polyphenols, 244
 selected colors and materials, 242
 Carob, 296, 298–300
 Carob flour (CF)
 carob pod, 107, 109
 carob protein isolate, 110–111
 carob seed, 109
Ceratonia, 110
 composition, 107
 emulsions and gels, 111–112
 in light foods, 108
 polyphenols, 109
Prosopis species
 American carob, 108
 añapa, 108
 Argentinean Food Codex, 108
Ceratonia, 110
 dietary fiber, 110
 gums, 109
 mesocarp, 108

- Mesquite pods, fraction of, 109
 - NOA region, 108
 - patay, 109
 - PER, 110
 - refined flour, 110
 - sugar, 108
 - toasted, 107
 - unsaturated fatty acids, 109
 - Carob protein isolate (CPI), 111
 - β -Carotene decolorization assay, 243–244, 246–247
 - Carotenoid, 234
 - content, 244–245
 - description, 242
 - Ceratonia siliqua*, 107, 109, 110, 296
 - Chenopodium pallidicaule* (see Cañahua)
 - Chestnut, 295, 296, 300
 - Chia. *See Salvia hispanica*
 - Chicha, 191
 - Chios Mastic Gum (CMG)
 - annual production yield, 273
 - antibacterial properties, 280
 - antimicrobial properties, 281
 - antioxidant activity, 281
 - application, 273
 - chemistry and physicochemical properties
 - biological properties of, 274–275
 - discoloration, 277
 - essential oil fraction, 273
 - P. lentiscus* var. *chia*, acidic fraction of, 275
 - polymeric fraction, 274
 - resin collection technique, 277
 - resin composition, 277
 - resin, natural fraction of, 275–277
 - triterpenes/triterpenoids, 273
 - cultivation of, 271
 - flavor, 272
 - food additive, 281
 - Greek product, 271
 - mastic tree, 271
 - medicinal properties, 278
 - Pistacia lentiscus* L. var. *chia*, 271
 - polymeric material, 272
 - traditional food products
 - alcoholic and nonalcoholic beverages, 284
 - chewing gum, 282–283
 - confectionery, desserts, and bakery products, 283–284
 - treatment
 - against *H. pylori*, 279
 - against *Helicobacter pylori*, 278
 - anticancer agent, 280
 - antimicrobial properties, 280
 - cardiovascular protection, 281
 - Crohn's disease, 279–280
 - dyspepsia, 279
 - gastric and duodenal ulcers, 278
 - heart attacks, risk reduce, 278
 - hepatoprotection, 281
 - peptic ulcers, 278
 - zinc, natural supplement, 280
 - Cobalamin. *See* Vitamin B₁₂
 - Corn Tortilla
 - ancient staple, 139–140
 - characteristics of
 - chemical, 143–144
 - physical, 141–142
 - thermal properties, 144–145
 - consumption of, 152
 - nixtamalisation
 - masa, 140
 - novel processes, 140–141
 - traditional process, 140
 - nutraceutical aspects, 150–152
 - nutritional aspects
 - proteins, 146–148
 - vitamins and minerals, 145–146
 - starch digestibility, 148–150
 - Cornelian Cherry (*Cornus mas* L.)
 - biochemical characterisation, 218–221
 - botanical characterisation, 217
 - dried leaves and fruits, 216
 - drienka'*, 216
 - economically used plant parts, 217–218
 - phytotherapeutic characterisation, 221–222
 - phytotherapeutic effects, 216
 - ripe fruits, 216
 - Cornus mas* L. (see Cornelian Cherry)
- D**
- Date seeds
 - antimutagenic properties, 233
 - antioxidant properties, 233
 - bread enrichment, 236–237
 - coffee-substitute hot beverages, 237–238
 - composition of, 234–235
 - energy source, 233
 - health attributes of, 235–236
 - staple food, 233
 - world production, 233
 - Degree of hydrolysis (DH)
 - ABTS decolorization assay, 201–202
 - ACE, 201
 - statistical analysis, 202

Dried fig, 301–302
 Dried fruits, 301–302
 DAFNE, 294
 functional properties
 fig, 301–302
 pear, 301–302
 plum, 302
 INE data, 294
 nutritional properties, 300–301
 Portugal, 295
 production, 295
 Dried pear, 301–302
 Dried plum, 302

E
Escherichia coli, 11, 58, 60, 168, 222, 281, 334, 359

F
 Fat burner, 354
 Fermentation process, 56
 Fermented foods, 39–40
 Fiber-rich fraction (FRF), 133, 134
 Fig (*Ficus carica*), 301
 Flow behavior index, 255
 Folin–Ciocalteu method, 235, 238, 242, 332
 Food fermentation, 9, 18, 21–23, 34, 40, 44, 46, 157, 172
Foods for Specific Health Use (FOSHU), 293
 Forgotten and unutilised plant species, 210–229
 black mulberry
 acid strawberry/tree strawberry, 210
 biochemical characterisation, 212–215
 botanical characterisation, 210–211
 full maturity, 210
 phytotherapeutic characterisation, 215–216
 phytotherapeutic purposes, 210
 plant parts utilised economically, 211–212
 Cornelian Cherry
 biochemical characterisation, 218–221
 botanical characterisation, 217
 dried leaves and fruits, 216
 economically used plant parts, 217–218
 phytotherapeutic characterisation, 221–222
 phytotherapeutic effects, 216
 ripe fruits, 216
 ecological species, 209
 in Slovakia, 210

Strategic Action Plan, 209
 True Service Tree, 222
 biochemical characterisation, 225–227
 economically exploited plant parts, 227–228
 genus *Sorbus* botanic characterisation, 223–225
 in Central Europe, 223
 origin of, 223
 phytotherapeutic characterisation, 228–229

Functional food, 4, 75, 293
 cereals or cereal constituent applications, 175
 definition, 9
 dried fruits, 300–303
 microbial GABA, 59
 nuts, 295–300
 Slovak products (*see* Slovak products)
 tarhana, 95

G
 γ -Aminobutyric acid (GABA), 58–59
 Genetically modified organisms (GMOs), 180
 Grape (*Vitis vinifera* L.), 301

H
 Habanero pepper. *See Capsicum chinense*
 Hazelnuts, 296–298
 Honey
 annual production, 323
 antibacterial activity, 333–335
 antioxidant, 331–333, 340–341
 applications, 340
 bioactive compounds, 339
 by-product, 342
 components, 325
 composition of, 340
 acidity, 328
 carbohydrates, 327–328
 electrical conductivity, 327
 hydroxymethylfurfural, 329–330
 pH value, 328
 phytochemicals, 325
 proteins and amino acids, 329
 water content, 325
 definition, 323
 elements, 330–331
 honey bees, 340
 nutraceuticals and medical ingredients, 339
 physicochemical parameters, 325

- prebiotic properties, 335–336
 - quality of, 325
 - secondary metabolites, 339
 - sweetener, 323
 - traditional honey products
 - Abbamele/Águamel, 344–350
 - mead, 342–344
 - traditional medicine, 340
 - types of, 324–325
- Horchata de Chufa
 - cultural and economic values, 371–372
 - functional characteristics, 372–373
 - manufacture process, 373–374
 - nutritional characteristics, 372–373
- Hydrocolloids, 255

- I**
- Insoluble dietary fiber (IDF), 133, 195
- Institute of Food Research (IFR), 298
- International Food Information Council (IFIC)*, 293

- K**
- Kefir, 92–94
- Klebsiella pneumoniae*, 30, 32

- L**
- Lactic acid bacteria (LAB)
 - antimicrobial compounds, 158
 - biopreservatives, 175
 - carbon dioxide, 176
 - diacetyl, 177
 - hydrogen peroxide, 176
 - low molecular weight, 177
 - organic acid, acetaldehyde, and ethanol, 176
 - bacteriocins, 177–178
 - food preservation and human therapy, 178
 - genus *Lactobacillus*, 179
 - production and modeling, 179–181
 - beneficial properties of, 168–171
 - in cereal-based fermented products, 158–166
 - morphology, 158
 - starter cultures
 - backslipping, 172
 - bacteriocins, 172
 - cereal-based probiotic foods, 173–175
 - commercial, 172
 - definition, 158
 - disadvantages, 172
 - factors, 172
 - functional starters, 172
 - Lactobacillus* spp. and *Bifidobacterium* spp., 171
 - natural, 172
 - potential health benefits, 158
 - raw material, 171
 - recombinant DNA technology, 173
 - wild-type strains, 172
 - Thai-fermented foods and beverages, 44
- Lactobacilli, 11, 168, 171
- Lipid antioxidants, 3
- Liquid collection method, 277
- Locust bean gum (LBG), 109
- Lokum (Turkish Delight), 101–103
- Loog-pang, 45

- M**
- Mastiha. *See* Chios Mastic Gum (CMG)
- Mediterranean diet, 294, 296, 297, 303
- Morus nigra* L. (*see* black mulberry)

- N**
- Nixtamalised corn flour (NCF), 140–141
- Nopalitos, 254
- Nutraceutical compounds, 189
 - amaranth
 - fatty acids, 192
 - protein content, 192, 195
 - squalene, 192, 195
 - Salvia hispanica*
 - antioxidants, 194
 - dietary fiber, 194–195
 - lipid content, 194
 - proteins, 195
- Nuts
 - cardiovascular protective properties, 297
 - DAFNE, 294
 - functional properties
 - almond, 297–298
 - carob, 298–300
 - chestnut, 300
 - hazelnuts, 297–298
 - pine, 300
 - walnuts, 298–299
 - INE data, 294
 - nutritional properties, 295–296
 - Portugal, 295
 - production, 295

O

- Ochratoxin A (OTA), 57
- Olive trees, 312
- Opuntia ficus-indica* (see Cactus products)
- OTA. See Ochratoxin A (OTA)
- Oxygen radical scavenging capacity (ORAC), 235, 237, 238, 362

P

- Pistacia lentiscus*, 280
- Particles classification, 272
- Pastirma
 - consumption, 89
 - functional properties, 89
 - health effects, 89
 - identification, 88
 - production, 88
- Patay, 109
- Pekmez, 98–99
- Phenolic antioxidants, 234
- Phenolic compounds, 343
- Phenolics, 242
- Phoenix dactylifera* L. (see Date seeds)
- Pine, 296, 300
- Plasticizer, 272
- Polyphenols, 5, 6, 109
- Polyunsaturated fatty acid (PUFA), 136
- Prebiotics
 - criteria for, 12–13
 - definition, 11
 - food applications, 13–14
 - gut microflora, 11
 - honey, 335–336
 - human colon, 9
 - intestinal microflora, 11
 - lactobacilli and bifidobacteria, 11
 - prebiotic carbohydrates, definition, 11
 - safety, 14
 - Thai-fermented foods and beverages, 59–64
- Prediction of glycaemic index (pGI), 150
- Probiotics
 - benefits, for human organism, 11
 - in capsules, microorganism, 10
 - criteria for, 12–13
 - definition, 10, 168
 - in fermented dairy products, 10
 - food applications, 13–14
 - gut microflora, 11
 - human colon, 9
 - LAB, 168
 - lactobacilli and bifidobacteria, 11
 - malabsorption, 11
 - safety, 14

Thai-fermented foods and beverages, 59–64

- Protein efficiency ratio (PER), 110, 146
- Protein-rich fraction (PRF), 133, 134

Q

- Quality protein maize (QPM), 143, 146

R

- Rapidly digestible starch (RDS), 148
- Recombinant DNA technology, 173
- Resistant starch (RS), 148, 152
- Retrograded resistant starch (RRS), 150
- RS. See Resistant starch (RS)

S

- Salvia hispanica*
 - ACE inhibitory activity, 203–204
 - agua de chia*, 131
 - Alcalase® 2.4 L FG, 200
 - antioxidant activity, 204–205
 - antioxidant capacity, 117
 - in Aztec Empire, 199
 - carbohydrate, 116
 - chia flour, 132
 - chia oil characterization, 133, 136
 - chia seed, 132
 - chia seed oil, 131
 - composition, 131
- DH
 - ABTS decolorization assay, 201–202
 - ACE, 201
 - statistical analysis, 202
 - diabetes, dietary treatment, 116
 - essential amino acids, 115
 - fatty acid composition, 116–117
 - Flavourzyme® 500MG, 200
 - FRF, 132–134
 - health benefits, 189
 - history, 115
 - Labiatae family, 115, 199
 - low commercial value, 189
 - millennia, 131
 - nutraceutical compounds
 - antioxidants, 194
 - dietary fiber, 194–195
 - lipid content, 194
 - proteins, 195
 - potential health benefits, 199
 - PRF, 132–134
 - protein content, 115
 - protein-rich fraction, 200, 202–203

- proximate composition, 132–134
- statistical analysis, 133
- White seeds, 199
- Sattoo, 191
- Shalgam drink, 99–101
- Short-chain fatty acids (SCFAs), 148, 152
- Slovak products, 76–85
 - bryndza, 75, 80
 - for human health, 84–85
 - history, 81–82
 - production technology, 82–83
 - plum jam, 75
 - for human health, 80
 - plum land varieties fruit, characteristics of, 76–77
 - traditional production technology of, 77–79
- Slovak sheep cheese. *See* Bryndza
- Slovenian honey. *See* Honey
- Slowly digestible starch (SDS), 148
- Soluble dietary fiber (SDF), 195
- Sorbus domestica* L (*see* True Service Tree)
- Spontaneously hypertensive rats (SHRs), 204
- Staphylococcus* spp., 45
- Sterols, 234
- Sucuk
 - consumption, 90
 - functional properties and health effects, 90
 - identification, 89
 - production, 89–90
- T**
- Tarhana, 94–95
- Tea, 354–358, 362–363
 - antibacterial and antiviral agent, 359
 - antioxidant properties, 353
 - beneficial impact, 358
 - bioavailability and absorption of, 360–361
 - biological activity, 358
 - Biological Activity and Health Effects
 - ancient chewing gum, 358
 - caffeine, 359
 - catechins toxicity, 360
 - commercial products, 363
 - cultivation and manufacture
 - average daily consumption, 354
 - chemical composition, 355–357
 - Chinese tea shrub, 354
 - classifications, 354
 - fermentation, 354–355
 - green tea, 354
 - inactivate oxidative enzymes, 355
 - machines, 355
 - mountainous, 354
 - oolong tea, 354
 - picking, 354
 - polyphenol content, 357
 - rolling process, 354
 - var. *assamica*, 354
 - discovery, 353
 - health and prevention of many diseases, 353
 - negative effect of, 359
 - polyphenols, antioxidative activity, 361–362
 - flavonoids, 361, 362
 - properties, in food, 362–363
 - polyphenols, in body weight regulation, 360
 - therapeutic value, 358–359
- Thai-fermented foods and beverages
 - classification of, 40
 - cottage and small-scale levels, 46
 - fermented fish (pla-ra), 39
 - fermented minced fish cake (Som-fug), 46–49
 - fermented pork (nham), 39
 - fish sauce (nam-pla), 39
 - khanom-jeen/fermented rice noodles, 50–51
 - khao-mak/sweet alcoholic rice paste, 55–56
 - microorganisms, 46
 - Bacillus* spp., 43–44
 - Enterococcus thailandicus*, 46
 - LAB, 44
 - Lentibacillus halophilus*, 46
 - physio-biochemical modifications, 40
 - Piscibacillus salipiscarius*, 46
 - Staphylococcus* spp., 45
 - Thai sweet-low alcoholic snack (khao-mak), 46
 - yeast and mold, 44–46
 - nham/fermented pork sausage, 49–50
 - phak-gad-dong/fermented mustard leaf, 53–54
 - potential health benefits
 - antimicrobial substances, 58
 - antioxidative activity, 59
 - bioavailability, 64–65
 - food safety and food preservation, 56
 - functional active compounds, 56
 - GABA, 58–59
 - microbial enzymes, 56–57
 - micronutrient synthesis, 64–65
 - prebiotics, 59–64
 - probiotics, 59–64
 - qualitative attributes, 40
 - rice wines, 53–54
 - thua-nao/fermented soybean, 52–53

- Tiger nut. *See* Horchata de Chufa
- Traditional foods, 3, 9–14, 88–103
 and *Prosopis* species, 108–109
 antioxidants (*see* Antioxidants)
 definition, 87
 prebiotics
 bifidobacteria, 11
 criteria for, 12–13
 definition, 11
 gut microflora, 11
 human colon, 9
 intestinal microflora, 11
 lactobacilli, 11
 prebiotic carbohydrates, definition, 11
 safety, 14
 synbiotics, 12
 probiotics
 benefits, for human organism, 11
 criteria for, 12–13
 definition, 10
 fermented dairy products, 10
 food applications, 13–14
 gut microflora, 11
 human colon, 9
 in capsules, microorganism, 10
 lactobacilli and bifidobacteria, 11
 malabsorption, 11
 safety, 14
 Turkish (*see* Turkish traditional foods)
- Trolox equivalent antioxidant coefficient (TEAC), 202, 246
- True Service Tree (*Sorbus domestica* L), 222
 biochemical characterisation, 225–227
 economically exploited plant parts, 227–228
 genus *Sorbus* botanic characterisation, 223–225
 in Central Europe, 223
 origin of, 223
 phytotherapeutic characterisation, 228–229
- Turkish traditional foods, 88–92, 94–101
 dairy-based traditional products, 92–94
 ayran, 90–92
 sucuk
- kefir, 92–94
- legume-based traditional products
 bulgur, 95–97
 tarhana, 94–95
- lokum/Turkish delight), 101–103
- meat-based traditional products
 pastirma, 88–89
 sucuk, 89–90
- vegetable- and fruit-based traditional products
 pekmez, 98–99
 shalgam drink, 99–101
- V**
- Viscosity coefficient, 255
- Vitamin B₁₂
 anemias, fermented fish, 23–25
 bacterial strains, 25
 biosynthesis, 18
 deficiency, 17
 dietary sources of, 17
 food fermentation, 21–23
 in microorganisms, 18–19
Klebsiella spp., bioenrichment
 tempe, 29–30
 thua-nao, 30–34
 lime fruit waste products, 25
 metabolic process, 17
 non-vegetarian foods, 19–20
Propionibacterium spp., bioenrichment, 25
 kefir, 27–28
 kimchi, 28–29
 tempe, 26–27
 recommended dietary intake, 18
 requirement, 18
 soybean paste, 25
 vegetarian foods and deficiency, 20–21
- Vitamin C, 242
- W**
- Walnuts, 295, 296, 298–299