

# Index

## A

- ABA-responsive element (ABRE), 82
- Abiotic stresses, 9, 12, 17–19, 253
- Abscisic acid (ABA), 75, 181, 229, 253, 289
- Acaryochloris*, 43
- Acetolactate synthase gene (OsALS), 21
- Adenosine triphosphate (ATP), 178
- ADP glucose pyrophosphorylase (ADPGPPase), 11
- Advanced yield trial (AYT), 130
- Aeluropus littoralis*, 228
- Aerobic rice development, 154
- Agronomic nutrient management (ANM) system, 352
- Agronomic traits, 456
- Alcohol dehydrogenase (ADH), 181
- Alcoholic fermentation (AF) pathway, 178
- Aldehyde dehydrogenase (ALDH), 181
- Allium sativum*, 406
- Alternate wetting and drying (AWD), 153, 303, 461
- Amiprofos-methyl (APM), 431
- Anaerobic germination (AG)
  - breeding rice, 193–195
  - crop establishment, 178
  - physiology and molecular mechanisms
    - AF, 178
    - amylases, 180
    - anaerobic response index, 178
    - anoxia, 178–181
    - ATPs, 178
    - carbohydrate, 180
    - carbon, 181
    - cellulose microfibrils, 182
    - efficient water imbibition, 180
    - energy, 181
    - ethylene, 181
    - expression analysis, 181
    - genetic factors, 182, 183, 185
    - genotypes, 179, 180
    - hypoxia, 181
    - hypoxic conditions, 178
    - matrix polysaccharides, 182
    - mechanism of action, 180
    - nonenzymatic proteins, 182
    - O<sub>2</sub> concentration, 179
    - water stress, 179
  - rice genotypes, 178
  - seed longevity
    - crop yield, 186
    - genetic factors, 187, 188
    - life cycle, 186
    - nongenetic factors, 186
    - physiology and molecular mechanisms, 186, 187
    - rice varieties, 186
  - seedling vigor
    - crop establishment, 188
    - genetic factors, 190, 193
    - physiology and molecular mechanisms, 188, 190
    - soil/water, 188
    - weed competition, 188
  - seeds, 178
    - weed control mechanism, 178
- Androgenesis, 426–430, 436, 442
- Anther culture, 426, 427, 429–431, 434, 436, 437, 441, 442
- Antioxidant enzymes, 186
- Antioxidant polyphenols, 187
- Antioxidant production, 186
- Apparent amylose content (AAC), 133

- Arabidopsis phytochrome*, 8  
*Arabidopsis*, 37, 44, 47, 48, 187, 245, 473  
 Arsenic contamination, 287, 302, 303, 305  
 Arsenic-induced toxicity symptoms  
   germination, 289, 291  
   reproductive growth, 291, 292  
   vegetative growth, 291  
 Arsenic speciation, 288, 305  
 Arsenic stress responses  
   arsenate uptake, 297  
   arsenite uptake, 295, 296  
   biological function, 299  
   biological system, 284  
   biotic and abiotic stresses, 282  
   cereal crop, 282  
   climate change, 283  
   detoxification, 299  
   enzymatic reduction, 299  
   factors, 295  
   food demand, 283  
   global food security, 283  
   grain and straw, 283  
   groundwater, 283  
   growth stages, 290  
   health-related concerns, 283  
   heavy metal contamination, 283, 285  
   human agricultural advancement, 282  
   mitigation strategies, 303, 304  
   oxidative stress response, 300, 301  
   paddy soil  
     climate change, 286  
     fungicides, 286  
     groundwater, 286  
     pesticides, 286  
     plant nutrients, 288  
     rice ecosystem, 287, 288  
     subsurface mobilization, 286  
     surface water, 286  
 QTL, 292–294  
 rainfed ecosystem, 282  
 rice production and consumers, 282  
 rice-growing environments, 288  
 root epidermis and exodermis, 299  
 root plaque formation, 301, 302  
 state of knowledge gaps, 302, 303  
 translocation, 296–298  
 uptake and storage, 298  
 worldwide distribution, 287  
*Arthrobacter globiformis*, 18  
 Artificial genome doubling, 431  
 Artificial intelligence (AI), 362  
 Ascorbate peroxidase (APEX), 229  
 Asian rice gall midge (ARGM), 381  
 Atmospheric CO<sub>2</sub> concentrations, 35  
 Atmospheric methane, 173  
 ATP-binding cassette (ABC) transporter, 300
- B**  
*Bacillus thuringiensis*, 15, 16, 400  
 Backcross-selective introgression lines  
   (EB-SILs), 63, 81  
 Bacterial blight (BB), 321, 322  
 Bacterial leaf streak (BLS), 325, 350  
 Bangladesh Rice Research Institute  
   (BRRI), 302  
 Bayesian network analysis, 456  
 Biochemical factors, 187  
 Biochemical mechanisms, 473  
 Biofortification, 438, 439, 450  
   agronomic traits, 481  
   conventional plant breeding, 472, 473  
   endogenous genes, 481  
   Fe concentration, 472, 480, 481  
   genetic engineering, 475, 480  
   genetically modified (GM) crops, 480  
   genome editing, 481  
   genome editing technology, 481  
   genome-edited crops, 481  
   homozygous condition, 481  
   iron deficiency, 472  
   iron uptake, 473, 475  
   micronutrient deficiency, 472  
   molecular characterization, 481  
   multi-location and multi-season field  
     trials, 481  
   nutritional deficiency, 472  
   nutritional traits, 481  
   transgenic plants, 481  
   translocation, 473, 475  
 Biofortification Priority Index, 451  
 Bioinformatics  
   AI, 362  
   datasets, 360  
   genome databases, 360  
   GWAS, 361  
   mapping QTLs and genes, 361  
   ML, 362  
   scientific community, 360  
   speeding up breeding, 362  
   whole-genome sequence, 360  
 Biological system, 284  
 Biophysical stresses, 12  
 Biosynthesis, 41  
 Biotic processes, 204  
 Biotic stress, 12, 13, 78, 79

- Bipolaris oryzae*, 353  
 Blast disease, 322  
 Breeding approaches  
   genetic transformation, 264, 265  
   genome editing, 265, 266  
   LTS tolerance, 261–263  
   plant breeders, 261  
   SIB, 263, 264  
   strategies, 261  
 Breeding rice, 193–195  
 Breeding strategies, 346, 347  
 Broad-spectrum resistance genes, 351, 352  
 Brown planthopper (BPH), 16, 436
- C**  
 Calcineurin B-like protein kinase  
   (CIPK15), 181  
 Callus, 427–431  
 Calmodulin-binding transcription activators  
   (CAMTAs), 253  
 Calvin–Benson–Cycle enzymes, 40, 41  
 Candidate gene cloning, 408  
 Carbohydrates, 8  
 Carbon concentrating mechanism (CCM), 39  
 Carboxysome, 40  
 Catalase (CAT), 229  
 Cellular metabolic activities, 186  
 Cell-wall components, 353  
 CFTP, *see* Critical sterility temperature  
   point (CSTP)  
 Characterization-based approach, 408  
 Chemosensory protein (CSP) genes, 398  
 Chloroplast, 49, 50  
 Chloroplast electron transport capacity, 42  
 Chromosome, 253  
 Chromosome segment substitution lines  
   (CSSLs), 395  
 Climate change, 173–176, 204, 205  
 Climatic environments, 205  
 Clustered regularly interspaced short  
   palindromic repeats (CRISPR), 21,  
   356, 357  
 Clustered regularly interspaced short  
   palindromic repeats/CRISPR-  
   associated protein (CRISPR/  
   Cas), 265  
 CO<sub>2</sub> concentration, 32  
 CO<sub>2</sub> diffusion, 49, 50  
 CO<sub>2</sub> transport, 32, 38  
 Cold stress (CS), 75  
*COLD1* gene, 254  
 Cold-responsive (COR) genes, 230, 245  
 Cold-stress response, 257  
 Cold-tolerant genotypes, 262  
 Comparative transcriptome analysis, 256  
 Conventional breeding-selection  
   approaches, 15  
 Conventional plant breeding, 472, 473  
 Critical sterility temperature point  
   (CSTP), 111  
   controlled-temperature screening  
     conditions, 113  
   field screening, 113, 114  
   seed production, 113  
   sequential seeding, 113, 114  
   temperatures, 112  
   tracking technique, 112  
 Crop plants, 5  
 Crop protection, 396  
 Crop simulation models, 173–174  
 Crystalline, 400  
 Culture technology, 437  
 Cumulative drought stress response index  
   (CDSRI), 151  
 Cyanobacterium, 33  
 Cyclic electron transport, 42  
 Cytokinin biosynthesis, 257  
 Cytoplasmic male sterile (CMS), 3, 100, 440
- D**  
 Damage-associated molecular patterns  
   (DAMPs), 348  
*Datura innoxia*, 427  
 Defense-related genes (DR), 348  
 Deoxymugineic acid synthase (DMAS)  
   enzymes, 473  
 Designed QTL pyramiding (DQP), 65, 81, 85  
 Differentially accumulated proteins  
   (DAPs), 121  
 Differentially expressed genes (DEGs), 257  
 Dimethylarsinic acid (DMA), 288  
*Dioscorea batatas*, 406  
 Direct-seeded rice (DSR), 74, 303  
   climate change, 173–176  
   crop establishment, 173, 196  
   ecosystems, 172  
   environment, 173  
   rainfed areas, 172  
   rice production, 195  
   topographical location, 172  
   water and labor requirements, 195–196  
   water scarcity, 173–176  
   water supply, 172  
   worldwide rice supply, 171

## Disease resistance

## breeding

- bacterial blight, 337, 344
- blast resistance, 344
- broad-spectrum resistance, 337
- conventional breeding methods, 326
- disease-resistant cultivars, 326
- DNA sequencing, 344
- gene-linked molecular markers, 337
- MABB, 344
- marker-assisted selection, 337
- MAS, 344
- molecular markers, 337
- multiple disease, 346, 347
- pathotypes, 337
- pyramiding, 345, 346
- varieties, 346

genetics, 325, 326 (*see also* Molecular mechanisms)

## Doubled-haploid (DH) approach, 125, 454

advantages, 427

albinos, 430

androgenesis, 426, 427, 430

## application

- biofortification, 438, 439
- cytoplasmic and nuclear genomes, 440
- development, 437, 438
- food security, 434
- gene stacking, 440
- GS, 441
- hybrid rice, 434, 435
- MAS, 441
- photosensitive rice genotypes, 434
- QTLs, 435–437
- reverse breeding, 440
- transgenic research, 441

basal media, 429

biotechnological approaches, 427

breeding cycle, 426

carbohydrate source, 429

chemical and physical factors, 428

chromosome doubling, 427

conventional breeding, 427

crop improvement, 426, 427

culture efficiency, 430

factors, 427

genomic regions, 442

genomics, 426

gynogenesis, 426

high-yielding rice cultivars, 426

*in vitro* morphogenic responses, 430

*indica* rice, 426, 442

*japonica* cultivars, 426

microspore development, 427, 429

microspore stage, 427

microspores, 430, 442

nitrogen composition, 429

phenomics, 426

physiology, 427

plant growth regulators, 429

## ploidy analysis

chromosome, 432

cytological approach, 433

flow cytometry, 433

high-throughput cell analysis, 433

homozygous lines, 431

molecular markers, 434

morphological evaluation, 432

nuclear DNA content, 432

pollen fertility analysis, 433

technology, 426

Double-stranded RNAs (dsRNA), 396

Drought, 149–152, 156, 157

**E**

Early seed germination (ESG), 74

Early seedling vigor (ESV), 74

Eating and cooking quality (ECQ), 133

Economic water scarcity, 174

Effector-binding elements (EBEs), 358

Effector-triggered immunity (ETI), 348, 408

Efficient screening technologies, 352

Electron transport, 41–45, 47, 48

Endogenous switching regression (ESR)

model, 86

Endomitosis, 431

Environment-sensitive genic male sterility

(EGMS), 4

Enzymes, 451

*Erwinia uredovora*, 20

*Escherichia coli*, 161

Estimated Average Requirement (EAR), 472

Ethyl methane sulfonate (EMS), 320, 395

**F**

Ferric-chelate oxidase 2 (FRO<sub>2</sub>), 473

Fertility-sterility alteration, 115

Flow cytometry, 433

Food-based solutions, 452

Food security, 60

Free-air CO<sub>2</sub> enrichment (FACE)

conditions, 41

Functional genetic units (FGUs), 84

Fungicides, 286, 353

**G**

- Galanthus nivalis* agglutinin (GNA), 406
- Gall midge (GM), 16
- Gelatinization temperature (GT), 133
- Gel-based protein, 259
- Gene editing, 21
- Gene expression analysis, 264
- Gene transfer, 399, 400, 407
- General combining ability (GCA), 130
- Genes, 292, 293, 297, 300, 303, 304
- Genetic basis, 473
- Genetic breeding, 261
- Genetic engineering, 406, 475, 480
- Genetic modification (GM) technology, 320
- Genetic resources, 208, 209
- Genetic transformation, 264, 265
- Genetics, 325, 326
- Genome editing, 265, 266
- Genome editing approach, 398, 399
- Genome editing tools (GETs), 265
  - biotic stress factors, 358, 359
  - CRISPR, 356, 357
  - disease resistance, 354
  - genome engineering, 354
  - introgress disease resistance, 354
  - MNs, 355
  - nonbiotic stress-related phenotypes, 359
  - ZFNs, 355, 356
- Genome-wide association studies (GWAS), 183, 232, 255, 256, 361, 456
- Genomic estimated breeding value (GEBV), 159, 457
- Genomic resources, 256
- Genomic selection (GS), 160, 441, 457
- Genomic tools, 17
- Genomics
  - genome editing approach, 398, 399
  - RNAi approach, 396, 398
- Genotyping by sequencing (GBS), 79
- Germplasm enhancement, 459, 460
- Gibberellic acid (GA), 181, 257, 289
- Global climate variability (GCV), 60, 222
- Global warming, 204, 205
- Glutathione, 299
- Glutathione S-transferase (GST) genes, 397
- Glycine decarboxylase (GDC), 37
- Glyoxalase, 36
- Grain, 472, 473, 475, 480, 481
- Grain quality, 76–78, 437
- Green Revolution (GR), 2, 3, 73, 173
- Green super rice (GSR), 9, 10, 264, 345
  - ABA signaling pathways, 82
  - abiotic stresses, 80
  - achievements, 86
  - advanced genotyping technology, 80
  - alkalinity, 61
  - biotic and abiotic stress, 62
  - breeding and population
    - development, 63, 65
  - breeding strategy, 80
  - candidate genes, 68
  - complex abiotic stress tolerances, 62
  - conventional and marker-assisted breeding
    - approaches, 62
  - conventional breeding approaches, 62
  - crop growth stages, 61
  - crop yield, 61
  - development, 62
  - distribution, 81
  - drought, 61
  - ecosystems, 62
  - food demand, 60
  - food security, 60
  - genetics
    - biotic and abiotic stresses, 67
    - biotic stress, 78, 79
    - drought tolerance, 67, 69, 70
    - grain quality, 76–78
    - LTS, 75, 76
    - NUE, 73
    - QTLs and breeding materials, 67
    - salinity tolerance, 70, 71
    - submergence tolerance, 71, 72
    - weed-competitive (WC) ability
      - traits, 74, 75
  - genotyping technologies, 79
  - grain quality, 62, 63
  - green traits, 62, 63
  - growth of population, 60
  - list of donors, 66–67
  - marker-assisted selection, 80
  - marker-trait association studies, 81
  - molecular genetics, 63
  - multiple stress tolerance, 79, 80, 82
  - multi-stress-tolerant varieties, 62
  - nutritional and health benefits, 61
  - physiological mechanisms, 63
  - physiological processes, 61
  - pyramiding QTLs, 80
  - QTLs, 68
  - rained environments, 80
  - rice cultivars, 61
  - rice varieties, 64
  - soil salinity, 61
  - stresses, 61
  - stress-tolerance genes, 82

Green super rice (GSR) (*cont.*)  
 stress-tolerant rice varieties, 61  
 sustainable development programs, 60  
 transcription factors, 82  
 Green traits, 62  
*See also* Green super rice (GSR)  
 Greenhouse gases (GHG), 173, 204

## H

Haploidy diploidization  
 artificial genome doubling, 431  
 spontaneous genome doubling, 431  
 HC-ATPase (HA) genes, 473  
 Heat shock proteins (HSPs), 121  
 Heat tolerance  
 agricultural production, 204  
 breeding, 213, 214  
 climate change, 204, 205  
 cultivars, 215  
 cultural practices, 214  
 damage, 205, 206  
 evaluation, 204  
 genetic efforts, 215  
 genetic resources, 208, 209  
 genetics, 211, 212  
 genotypic and morphological  
 characters, 215  
 global warming, 204, 205  
 marker-assisted gene, 215  
 marker-assisted recurrent selection, 215  
 physiology, 209–211  
 plant hormones, 214  
 reproductive stages, 204  
 rice, 206–208  
 rice germplasm, 214  
 rice production, 205, 206  
*Helicoverpa armigera*, 398  
 Herbicides, 286  
 High temperature, 204–210, 212, 214  
 High-throughput cell analysis, 433  
 Host-plant resistance, 15  
 Host-plant resistance gene deployment  
 classical breeding methods, 381  
 crambidae, 395  
 diopsidae, 395  
 gall midge, 381, 382, 394  
 leafhoppers, 394, 395  
 noctuidae, 395  
 phenotypic selections, 381  
 planthoppers, 394, 395  
 resistance genes, 395  
 Host-plant resistance mechanism, 78  
 Human health, 285

Hybrid Rice Development Consortium  
 (HRDC), 100, 133  
 Hybrid rice technology, 3, 4, 100  
 Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), 300  
 Hyperplasia, 381  
 Hypersensitive reaction (HR), 348, 382, 408  
 Hypertrophy, 381

## I

Indian Council of Agricultural Research-  
 National Rice Research Institute  
 (ICAR-NRRI), 149  
 Indole-2-acetic acid (IAA) homeostasis,  
 253, 257  
 Inducer of CBF expression (ICE), 230  
 Inductively coupled plasma optical emission  
 spectrometry (ICP-OES), 458  
 Industrial and urban sectors, 174  
 Infectious diseases, 353  
 Infrastructure, 174, 175  
 Insect pest management  
 biotechnology, 381  
 biotic stress, 380  
 chronic injuries, 380  
 dsRNA resistance, 410  
 feeding guilds, 380  
 genomic level  
 planthopper genomes, 407  
 rice-gall midge interactions,  
 408, 409  
 rice-planthopper interactions, 408  
 global warming, 380  
 host-plant resistance, 380  
 immune-level R-genes, 410  
 molecular approaches, 410  
 molecular biology, 381  
 molecular marker-based approaches, 381  
 pesticides, 410  
 production environments, 380  
 resistance genes, 385–388, 390, 392  
 rice crop, 410  
 rice ecosystem, 380  
 siRNA approach, 397  
 stem borers, 380  
 transgenes, 401–406  
 transgenic approach, 399, 400, 407  
 Insect pests, 15, 16  
 Insect-plant interaction, 407–409  
 Inter-connected breeding (IB) population, 76  
 Intergovernmental Panel on Climate Change  
 (IPCC), 60  
 International Center for Tropical Agriculture  
 (CIAT), 473

International Rice Research Institute (IRRI), 2,  
100, 178, 206, 394, 451, 472  
Interval mapping (IM), 185  
Introgression lines (ILs), 71, 264  
Iron deficiency, 472  
IRRI Genetic Resources Center, 182  
Isopentyltransferase gene (IPT), 161

## L

Leafhoppers, 394, 395  
Lesion length (LL), 322  
Linear electron transport, 42  
Lipid peroxidation, 300  
Liquid chromatography coupled with tandem  
mass spectrometry (LC-MS/  
MS), 258  
Low-temperature stress (LTS), 75  
  abiotic stresses, 222  
  air temperature, 224  
  antioxidative enzymes, 229  
  aquaporins, 229  
  biochemical parameters, 229  
  bioinformatics, 267  
  booting/flowering stage, 243, 244  
  breeding varieties, 229  
  calcium signals, 230  
  cellular metabolism, 230  
  chromosome, 236  
  cold stress, 229  
  cold-water screening, 224  
  colocalization, 238–241  
  crop growth stages, 223  
  diverse ecosystems, 222  
  food security, 222  
  genetic and genomic resources, 223  
  genomics, 266  
  germination, 223, 224, 232, 235, 236  
  growth stages, 224, 226–227,  
    233–235, 266  
  GS-related traits, 232  
  high-elevation areas, 224  
  high-throughput molecular  
    approaches, 225  
  high-throughput screening techniques,  
    225, 227  
  intracellular levels, 230  
  linolenic acid, 229  
  low-temperature stress (LTS)  
    systems biology, 267  
  mechanisms, 230  
  metabolome-wide association studies, 266  
  metabolomics, 266  
  metabolomics quantitative trait locus, 266

  molecular and physiological levels, 230  
  molecular marker technology, 227  
  molecular mechanisms, 266  
  natural cold-screening hotspots, 224  
  next-generation sequencing technology  
    tools, 223  
  omics-related technologies, 266  
  *OsNAC5* gene, 229  
  phenotypic evaluation, 225  
  phenotypic techniques, 225  
  photosynthetic activity, 227  
  photosynthetic process, 228  
  physiological and biochemical  
    methods, 229  
  physiological and metabolic activities, 230  
  plant growth period, 225  
  plant growth stages, 223  
  polymorphic SNPs, 267  
  protocols, 227  
  PSII, 228  
  QTLs, 237  
  reproductive stage, 224  
  rice production, 222  
  rice varieties, 228  
  screening methods, 224  
  seedling stage, 237, 241, 242  
  tolerance genes and stages, 246–252  
  water control, 223  
*LOX3* gene, 21  
Lysine-rich protein (LRP), 15

## M

Machine learning (ML), 362  
*Magnaporthe oryzae*, 13, 322  
Major resistance (MR) genes, 348  
Malnutrition, 450, 463  
MAMP-triggered immunity (MTI), 348  
Marker-assisted backcross (MABC), 120, 126  
Marker-assisted backcross breeding  
  (MABB), 344  
Marker-assisted backcrossing (MAB),  
  158, 254  
Marker-assisted breeding (MAB), 18, 156, 394  
Marker-assisted pyramiding, 158, 159  
Marker-assisted recurrent selection  
  (MARS), 159  
Marker-assisted selection (MAS), 125, 158,  
  178, 320, 441  
Matrix-assisted laser desorption/ionization  
  (MALDI-TOF MS), 258  
Meganucleases (MNs), 265, 355  
Metabolite quantitative trait locus  
  (mQTL), 261

- Metabolome-wide association studies (MWAS), 261
- Metabolomics, 260, 261
- Metal chelator, 475–480
- Microarray studies, 408
- Microbe-/pathogen-associated molecular patterns (MAMPs/PAMPs), 348
- Micronutrient, 452, 472, 473
- Mitogen-activated protein (MAP) kinase activity, 348
- Molecular approaches
  - agricultural crop plants, 321
  - bacterial blight, 316, 327–331
  - BB, 321, 322
  - biotic disease resistance, 326, 352–354
  - biotic stresses, 316
  - blast disease, 322
  - blast resistance genes, 332–337
  - BLS, 325
  - climatic conditions, 316
  - conventional breeding methods, 316
  - crop protection, 316
  - diseases, 316–319
  - DNA markers, 320
  - false smut, 324
  - food demand, 363
  - food security crop, 316
  - frequency of variations, 321
  - gamma-rays ( $\gamma$ -rays), 320
  - genetic improvements, 363
  - genome engineering technologies, 321
  - genome-editing technologies, 321
  - genome-editing tools, 321
  - genomic-assisted breeding tools, 363
  - genomics, 321
  - GM technology, 320
  - management practices, 316
  - methods, 316
  - mutation breeding, 320
  - nutrient fertilizers, 352–354
  - plant pathogens, 316
  - QTL, 320, 332–337
  - resistance, 320
  - R*-genes, 320
  - rice cultivation, 316
  - rice grain yield, 316
  - sheath blight (ShB) disease, 316, 322, 323
  - sheath blight-resistance genes, 338–343
  - ShR, 323, 324
  - techniques, 359, 360
  - tungro disease, 324, 325
- Molecular breeding
  - antioxidants, 156
  - GS, 160
  - MABC, 158
  - marker-assisted pyramiding, 158, 159
  - MARS, 159
  - MAS, 158
  - proteins, 156
  - QTL mapping, 156, 157
- Molecular genetic mechanisms, 302
- Molecular level, 17
- Molecular marker studies, 117
- Molecular markers, 337, 381, 394, 434
- Molecular mechanisms, 120, 121
  - bacterial blight, 349, 350
  - BLS, 350
  - broad-spectrum resistance genes, 351, 352
  - calcium-binding proteins, 245
  - cold-sensing calcium channel, 245
  - cold-stress conditions, 245
  - defense mechanisms, 347
  - elicitors, 349
  - gene expression patterns, 245
  - genes and transcription factors, 245
  - GWAS, 255, 256
  - innate immune system, 348
  - metabolomics, 260, 261
  - molecular patterns, 348
  - molecules, 348
  - pathogens, 347
  - pathogen-specific factor, 348
  - phosphorylation, 245
  - physiological and metabolic compounds, 245
  - plant resistance, 349
  - proteomics, 258, 259
  - rice blast, 350
  - sheath blight, 351
  - signaling pathways, 253–255
  - transcription factors, 245
  - transcriptomics, 256–258
- Monomethylarsonic acid (MMA), 288
- Mugineic acid, 473
- Multi-environment trials (METs), 130
- Multi-parent Advanced Generation Inter-Cross (MAGIC)-derived lines, 452
- Multiple stress tolerance
  - breeding products, 84–86
  - rice hybrids, 87
  - stress-regulated mechanisms, 82–84
- Multi-pronged approaches, 5
- Multi-trait and multi-environment (MTME) models, 457–458
- Multi-trait genomic selection, 457, 458
- Myeloblastosis (MYB), 257



**N**

NADH dehydrogenase-like (NDH) complex, 47  
 NADH quinone oxidoreductase (NQO), 406  
 Natural resistance-associated macrophage protein (NRAMP), 475  
 Near-isogenic line (NIL), 8, 212, 435  
 New breeding techniques (NBT), 481  
 New plant type (NPT), 7, 8  
*Nicotiana tabacum*, 37  
 Nicotianamine aminotransferase (NAAT), 473  
 Nicotianamine synthase (NAS), 473  
 Nicotinamide adenine dinucleotide (NAD), 181  
 Nitrogen, 352  
 Nitrous oxide, 173  
 Nucleotide-binding-leucine-rich repeat (NB-LRR), 348  
 Nucleotides, 398  
 Nutrient deficiency, 19  
 Nutrient fertilizers, 352–354  
 Nutrient-use efficiency (NUE), 73  
 Nutritional deficiency, 472  
 Nutritive quality, 19, 20

**O**

Observation yield trial (OYT), 130  
 Oligosaccharyl transferase (*OoOST*), 409  
 Optimum temperature, 205  
*Oryza australiensis*, 38  
*Oryza meridionalis*, 208  
*Oryza rufipogon*, 156  
 Oryzalin, 431  
 Oxidized macromolecules, 186

**P**

Pattern recognition receptors (PRRs), 348  
 Pattern-triggered immunity (PTI), 348, 408  
 Peroxidases, 181  
 Pesticides, 286  
 Phenotypic screening, 326, 346  
 Phenotypic traits, 183  
 Phenotypic variation (PV), 156  
 Phloem protein2 promoter (pPP2), 9  
 Phosphoenolpyruvate carboxylase (PEPC), 40  
 Phosphoglycerate (PGA), 33  
 Phosphoglycolate (PGO), 33  
 Photoperiod-sensitive genic male sterile (PGMS), 101  
 Photoperiod-sensitive male sterility (PGMS), 4

Photorespiration, 11, 35, 37  
 Photosynthesis, 11, 12, 473  
   air temperature, 31  
   Calvin–Benson–Cycle enzymes, 40, 41  
   CO<sub>2</sub> concentrations, 38–40  
   crop productivity, 31  
   crop yield, 32  
   electron transport, 41–44  
   environmental conditions, 32  
   greenhouse gases, 32  
   natural environments  
     Calvin–Cycle enzymes, 48, 49  
     CO<sub>2</sub> diffusion, 49, 50  
     electron transport, 45, 47, 48  
     genetic engineering approaches, 45  
     light intensity, 44  
     non-steady-state, 44  
     photosynthetic induction, 44  
     photorespiration, 35–37  
     Rubisco kinetics, 33, 35  
     steady-state and non-steady-state, 46  
     thermotolerance, 37, 38  
 Photosynthetic carbon metabolism (PCM), 12  
 Photosynthetic efficiency, 33  
 Photosystem-II (PSII), 228  
 Physiological and metabolic pathways, 353  
 Physiological characterization, *see*  
   Temperature-sensitive genetic male sterile (TGMS)  
 Phytochelatins (PCs), 299  
 Phytotoxicity, 302  
 Plant growth facility (PGF), 125  
 Plant growth regulators (PGRs), 428  
 Plant resistance, 349  
 Planthopper genomes, 407  
 Planthoppers, 394, 395  
 Pollen grains, 210  
 Polycistronic tRNA-gRNA (PTG), 356  
 Posttranslational changes, 259  
 Prime editing guide RNA (pegRNA), 357  
 Pronamide, 431  
 Prorich protein (PRP), 253  
 Protein content (PC), 19  
 Proteins, 258  
 Proteomic analysis, 212  
 Proteomics, 258, 259  
 Protospacer adjacent motif (PAM), 356  
*Pseudomonas*, 409  
*Pseudomonas fuscovaginae*, 324  
 Pyramiding lines (PDLs), 65  
 Pyrosequencing approach, 408  
 Pyrosequencing-based transcriptome analysis, 409  
 Pyruvate decarboxylase (PDC), 181

**Q**

Quantitative trait loci (QTL), 156, 157, 183, 211, 292–294, 320, 435–437, 453–455

**R**

Rapid generation advancement (RGA), 125  
 Rapid viscosity analysis (RVA), 133–134  
 Reactive oxygen species (ROS), 17, 70, 186, 300, 348  
 Receptor-like kinase (RLK), 160  
 Recombinant inbred lines (RILs), 183, 435  
 Regeneration, 426–431, 434, 441  
 Regional high-temperature damage, 205  
 Relative humidity (RH), 207  
 Repeat variable di-residues (RVDs), 356  
 Resistance mechanism, 382  
 Respiration, 473  
 Restriction fragment length polymorphism (RFLP) markers, 183  
 Reverse breeding, 440  
*Rhizoctonia solani*, 15, 322  
 Ribonucleoproteins (RNPs), 359  
 Rice (*Oryza sativa* L.)  
   food-deficit Asia, 1  
   gene editing, 21  
   gene pool, 5  
   GSR, 9, 10  
   huge demand projection, 1  
   hybrid rice technology, 3, 4  
   ideotype breeding, 7, 8  
   molecular and physiological breeding strategies, 2  
   physiological breeding approaches  
   abiotic stresses, 17–19  
   biophysical stresses, 12  
   biotic stresses, 13  
   diseases, 13–15  
   dwarfing genes, 10  
   insect pests, 15, 16  
   photosynthesis, 11, 12  
   starch biosynthesis, 11  
   plant architecture, 7, 8  
   root architecture, 9  
   self-sufficiency, 1  
   shoot and panicle architecture, 8, 9  
   strategies, 5  
   systematic breeding, 2  
   wild-weedy gene pools, 5, 7  
 Rice breeding programs, 452  
 Rice Diversity Panel 1 (RDPI), 255

Rice establishment methods

  direct-seeded rice (DSR), 177  
   puddled transplanted rice, 176

Rice leafroller (RLF), 16

Rice tungro bacilliform virus (RTBV), 14, 324

Rice tungro disease (RTD), 359

Rice tungro spherical virus (RTSV), 14, 324

Rice tungro virus (RTV), 13, 14

RNA interference (RNAi), 14, 396, 398

RNA-induced silencing complex (RISC), 396

Root architecture, 9

Root-related traits, 9

**S**

Salt overly sensitive (SOS) pathway, 18

Sarocladium oryzae, 323

Saturated soil culture (SSC), 149, 154

Secreted salivary gland proteins (SSGPs), 408

Seed production

  self-seed multiplication, 134, 135  
   two-line hybrid rice technology, 134

Seedling survival percentage (SSP), 227

Selective introgression breeding (SIB), 263, 264

Selective introgression lines (SILs), 84–85

Self-fertility, 135

Serial analysis of gene expression (SAGE), 360

Sexual barriers, 5

Sheath blight (ShB), 14, 322, 323

Sheath blight resistance (ShBR), 436

Sheath rot (ShR), 323, 324

*Silene vulgaris*, 187

Simple sequence repeat (SSR), 361, 434

Single marker analysis (SMA), 185

Single nucleotide polymorphisms (SNPs), 183

Single-guided RNA (sgRNA), 265

siRNA approach, 397

Site-specific nucleases (SSNs), 321

Small heat shock proteins (sHSP), 212

Small interfering RNAs (siRNAs), 396

Soil-and irrigation-based strategies

  aerobic rice development, 154  
   AWD, 153  
   depletions, 154, 155  
   environmental changes, 153  
   intensification, 155  
   outflows, 154, 155  
   saturated soil culture, 154  
   sprinkler irrigation, 155

Soil mulching, 155

Soil plant analysis development (SPAD), 227

Solar radiation, 204  
 Specific locus amplified fragment sequencing (SLAF-seq) technology, 255  
*Spodoptera exigua*, 398  
*Spodoptera litura*, 399  
 Spontaneous genome doubling, 431  
 Standard evaluation system (SES) scale, 322  
*Streptomyces pyogenes*, 356  
 Stress-regulated mechanisms, 82–84  
 Stress-responsive genes, 161  
 Superoxide dismutase (SOD), 229, 300  
 Susceptible host (SH), 408  
 Sustainable development goal, 162, 450  
 Systemic acquired resistance (SAR), 408

## T

Temperature stress, 205  
 Temperature-sensitive genetic male sterile (TGMS)  
   advantages, 111, 112  
   breeding pollen parents, 126  
   breeding trials, 130  
   China, 100  
   combining ability nurseries, 130  
   disadvantages, 111, 112  
   disease resistance, 133  
   food demand, 100  
   genetically modified (GM) domain, 137  
   genetics  
     identification, 117, 120  
     molecular mechanisms, 120, 121  
   genome editing tools, 137  
   grain quality, 133, 134  
   hybrid rice technology, 100  
   *indica/japonica* hybrids, 126  
   insect pest, 133  
   international scientific communities, 100  
   male-sterile lines, 138  
   multiplex gene editing, 137  
   mutation breeding, 122  
   pedigree breeding, 122  
   physiological characterization  
     fertility-sterility alteration, 115  
     pollen parental lines, 116, 117  
     sterility-fertility alteration, 115  
   public and private sectors, 101  
   pyramiding, 123  
   rapid fixation, 125, 126  
   rice production, 100  
   transfer, 123  
   two-line hybrid rice technology, 101, 111, 136, 137  
   two-line rice hybrids, 127

Thermotolerance gene (*TT1*), 212  
 Thioredoxin-mediated redox regulation, 49  
 Tiller angle (TA), 14  
 Tissue culture and genetic transformation protocols, 399  
 Total drought response index (TDRI), 151  
 Transcription activation-like effectors (TALEs), 349  
 Transcription activator-like effector nucleases (TALENs), 21, 265, 321, 455  
 Transcription factors (TFs), 82  
 Transcriptional/hormonal regulation, 9  
 Transcriptional regulation pathways, 253  
 Transcriptomics, 120, 256–258  
 Transformation-regeneration protocols, 18  
 Transgenic approach, 21, 399, 400, 407  
 Transgenic maize plants, 396  
 Transgenic plants, 32, 36  
 Transgenic strategies, 160, 161  
 Transgenics, 18  
 Trifluralin, 431  
 Tunable genotyping-by-sequencing (tGBS), 65  
 Two-dimensional gel electrophoresis (2-DE), 258  
 Two-line hybrid rice technology, 101, 111

## U

Uniform Blast Nursery (UBN), 322  
*Ustilaginoidea virens*, 324

## V

*Vigna aconitifolia*, 18  
 Vitamin deficiency–related health disorders, 20

## W

Water demand  
   abiotic stresses, 161  
   aerobic rice varieties, 148  
   climate changes, 161  
   environmental conditions, 148  
   food security, 148  
   human population, 148  
   plant-based strategies  
     cultivars/varieties, 150, 151  
     decreased stand density, 152  
     planting date, 152  
   plants, 149  
   rainfed ecosystem, 148  
   rice cultivation, 148  
   rice varieties, 161

Water demand (*cont.*)

- stress-tolerant cultivars, 148
- transgenic strategies, 160, 161
- water availability, 148
- water-deficient conditions, 148, 150
- water-limiting conditions, 149, 150
- water management, 148

## Water scarcity, 173–176

## Water-use efficiency, 40

## Weed Tolerant Rice-1 (WTR-1), 65

White-backed planthopper (WBPH)  
resistance, 436

## Whole-genome sequencing (WGS), 80

## Whole-genome sequence, 255

## Wide compatibility gene (WCG), 4

## Wide genetic variation, 207

## Wide-compatibility (WC) gene, 126

## Wild abortive (WA), 3, 112

*Wolbachia*, 407, 409**X***Xanthomonas*, 355*Xanthomonas oryzae*, 353*Xanthomonas oryzae* *pv.* *oryzae* (*Xoo*), 13

## X-ray fluorescence (XRF), 458

**Y**

## Yeast-like symbiont (YLS), 407

Yellow stem borer (*Scirpophaga incertulas*), 15**Z**

## Zinc-biofortified rice

## Bayesian network analysis, 456

## biofortification, 450

## breeding techniques, 451

## deficiencies, 450

## deficiency-related health

## consequences, 450

## donor identification, 452, 453

## food-based approach, 450

## food habits, 459

## genomic tools, 451

## genotype × environment effects, 461

## germplasm enhancement, 459, 460

## high grain Zn, 451, 452

## human health, 451

## mainstreaming, 460, 461

## micronutrients, 450

## multiple interventions, 450

## multi-trait genomic selection, 457, 458

## network of metal homeostasis genes, 455

## next-generation multi-nutrient, 462

## phenotyping, 458

## pre-breeding, 459, 460

## product control, 462

## QTLs, 453–455

## rice traceability, 462

## varieties, 462

## yield, 453

Zinc-finger nucleases (ZFNs), 265, 321, 355,  
356, 455