Some Other ELBS Low-Priced Editions

Abercrombie, M., Hickman, C. J. and Johnson, M. L. Andrewartha, H. G.	A DICTIONARY OF BIOLOGY	<i>Penguin</i>
Bailey, N. T. J.	INTRODUCTION TO THE STUDY OF ANIMAL POPULATIONS STATISTICAL METHODS IN BIOLOGY	<i>Chapman & Hall</i> <i>English Universities</i> <i>Press</i>
Barrington, E. J. W. Chapman, R. F.	INVERTEBRATE STRUCTURE THE INSECTS: STRUCTURE AND FUNCTION THE ECOLOGY OF INSECT POPULA- TIONS IN THEORY AND PRACTICE	<i>Nelson</i> <i>English Universities</i> <i>Press</i> <i>Chapman & Hall</i>
Clark, L. R., Geier, P. W., Hughes, R. D. and Morris, R. F. Hayes, W.	THE GENETICS OF BACTERIA AND THEIR VIRUSES	<i>Blackwell Scientific</i>
Elton, C. S.	THE ECOLOGY OF INVASIONS BY ANIMALS AND PLANTS	<i>Methuen</i>
Hughes, G. M.	COMPARATIVE PHYSIOLOGY OF VERTEBRATE RESPIRATION	<i>Heinemann Educational</i>
Imms, A. D.	A GENERAL TEXTBOOK OF ENTO- MOLOGY	<i>Methuen</i>
Lockwood, A. P. M.	ANIMAL BODY FLUIDS AND THEIR REGULATION	<i>Heinemann Educational</i>
Manning, A.	AN INTRODUCTION TO ANIMAL BEHAVIOUR	<i>Arnold</i>
Marshall, A. J. and Williams, W. D. Paul, J. Soulsby, E. J. C.	A TEXTBOOK OF INVERTEBRATE ZOOLOGY CELL BIOLOGY MÖNNIG'S HELMINTHS, ARTHRO- PODS AND PROTOZOA OF DOMESTI- CATED ANIMALS	<i>Macmillan</i> <i>Heinemann Educational</i> <i>Baillière Tindall</i>
Stephenson, G.	MATHEMATICAL METHODS FOR SCIENCE STUDENTS	<i>Longman</i>
Weatherburn, C. E.	A FIRST COURSE IN MATHE- MATICAL STATISTICS	<i>Cambridge University</i> <i>Press</i>
Wigglesworth, V. B.	THE PRINCIPLES OF INSECT PHYSIOLOGY	<i>Methuen</i>
Young, J. Z.	THE LIFE OF VERTEBRATES	<i>Oxford University Press</i>

Textbook of Zoology Vertebrates

Revised and largely rewritten by the late A. J. MARSHALL, D.Phil., D.Sc.
sometime Professor of Zoology and Comparative Physiology in Monash University

This is the Seventh Edition of *A Textbook of Zoology*, Vol. II
by the late T. JEFFERY PARKER, D.Sc., F.R.S.
sometime Professor of Biology in the University of Otago, Dunedin

and the late WILLIAM A. HASWELL, M.A., D.Sc., F.R.S.
sometime Professor of Biology in the University of Sydney

English Language Book Society and Macmillan Education

© Macmillan & Co Ltd 1962

All rights reserved. No part of this publication may be reproduced or transmitted, in any form or by any means, without permission.

First Edition	1898
Second Edition	1910
Third Edition	1921
Fourth Edition	1928
Fifth Edition	1930
Sixth Edition	1940
Reprinted	1943, 1947, 1949, 1951
Seventh Edition	1962
Reprinted	1963
Reprinted with corrections	1964
Reprinted	1966, 1967, 1972

ELBS edition first published 1972

Published by
THE MACMILLAN PRESS LTD
London and Basingstoke
Associated companies in New York
Toronto Dublin Melbourne
Johannesburg and Madras

SBN 333 05391 5 (hard cover)
333 02829 5 (paper cover)

ELBS edition: SBN 333 14085 0 (paper cover)

ISBN 978-1-349-00200-9 ISBN 978-1-349-00198-9 (eBook)
DOI 10.1007/978-1-349-00198-9

The paperback edition of this book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent, resold, hired out, or otherwise circulated without the publisher's prior consent, in any form of binding or cover other than that in which it is published and without a similar condition including this condition being imposed on the subsequent purchaser.

* * *

In conclusion, it must be pointed out that in order adequately to solve the problems of Zoology they must be approached from all sides. From the time of Cuvier to that of Owen comparative anatomy was the dominant branch of the science, and there was a tendency to depreciate the work of the 'mere' systematist and outdoor naturalist. For the last five and twenty years embryology has been in the ascendant, and the 'mere' anatomist has been somewhat overshadowed. To-day, hopeful signs of a renewed interest in ethology—the study of living animals under natural conditions—are accompanied by a tendency to look upon all laboratory work as necrology rather than biology—the study of corpses rather than of living things. But nothing is more certain than that if the new 'natural history' is to be superior to the old—more scientific, more concerned with the solution of general problems—it can only be by utilising to the full all that has been learnt in the laboratory in the departments of anatomy, physiology and embryology. (*Concluding paragraph, Parker and Haswell, 1st ed., 1898.*)

PREFACE TO THE SEVENTH EDITION

IT is now more than sixty years since the initial publication of 'Parker and Haswell'. Many changes were made in Volume II by Forster-Cooper in his Sixth Edition of 1940, but much of the original nineteenth-century text has nevertheless remained relatively unaltered. Consequently no apology is offered for the almost complete revision that follows.

When the publishers invited me to undertake the Seventh Edition they were kind enough to give me a free hand. For a while I thought of altering the whole scheme of the volume to one of my own devising. However, I decided (like my immediate predecessor) against the disruption of the 'general plan of a text-book that has served well so many generations of students'.

In general I have attempted, within the limits of available space, to retain basic morphology and yet at the same time emphasise functional aspects and, where possible, present animals as living creatures rather than as laboratory specimens. In this I have been helped by my wife, Jane Marshall, whose otherwise unacknowledged pen-drawings are scattered through the text: a collaboration, incidentally, that was responsible once or twice for minor domestic dissension owing to the conflicting claims of art and accuracy.

An innovation in the present volume is the inclusion of the elements of a classification under each illustration of a whole, or almost whole, animal. This may help the student more or less painlessly to familiarise himself with the animal's systematic position. Some trouble too, has been taken to provide a lavish cross-reference system so that comparative information can be found quickly and without tediously repetitive consultation of the Index or Contents.

It was at first hoped to include an appendix on the principles (and pitfalls) of systematics but there was no room for it. In any case, students can nowadays easily consult Mayr, Linsley and Usinger's *Methods and Principles of Systematic Zoology* (1953) or Simpson's *The Principles of Classification and a Classification of Mammals* (1945). (The last-named is additionally noteworthy in its lack of the solemnity that so many scientists believe to be indispensable to good scholarship.) Today, with such books readily available, only egotism or perversity, rather than ignorance, can be responsible for an extension of the taxonomic morass into which so many groups have been thrust: an unhealthy swamp that will need the labour of a century to remove.

The question of references has been one of some difficulty. Clearly it is

impossible to cite every source of information, most of which has been taken from original papers. A compromise has been made. In the first place, key references, leading to many others, have been included in the bibliography. Where an authority has been cited personally in the text, the publication date has been deliberately omitted. It is believed that it will provide a mild exercise in ingenuity and bibliographic skill for the honours man to discover the source from which he requires further information.

Another debatable point has been the appropriate amount of palæontology that should be given in a text-book of zoology. Some of my colleagues are of the opinion that there was already included rather too much information about fossil forms. I believe, however, that it is impossible to gain a good general understanding of living animals—either in form or function—without an adequate appreciation of what is known of their ancestry. Therefore the fossil sections have been expanded.

The elimination of *Neoceratodus* as the lung-fish 'type', and the substitution of *Protopterus*, needs brief explanation. The former is so rare, and rigidly protected, that not even an Australian student can confidently hope ever to dissect one. Living *Protopterus*, on the other hand, can be bought in unlimited numbers in fish-markets over wide areas of Africa. Thus they are available (for demonstration purposes at least) to university departments anywhere. It may seem surprising, in this view, that *Salmo* is retained as the teleost 'type'. However, there is employed for the purpose today (in the many countries where this book is used) such a wide diversity of bony fishes that any new choice would have only limited and local value.

Had I not had my first systematic instruction in zoology and physiology from students of Haswell, and made my first supervised dissections in the school that he founded, I would never have essayed the present revision. Yet, once begun, the task became surprisingly agreeable. It has been lightened by the rigorous but kindly criticism of friends who read sections of the manuscript dealing with subjects of their special interests. These are as follows:

E. C. Amoroso	A. J. E. Cave	E. W. Knight-Jones	D. L. Serventy
F. J. Aumonier	G. E. H. Foxon	K. A. Kermack	V. B. Sheffer
E. W. Baxter	F. C. Fraser	Dennis Lacy	J. R. Simons
A. d'A. Bellairs	P. H. Greenwood	D. A. McDonald	R. Strahan
Ruth Bellairs	Frank Goldby	J. D. Macdonald	E. I. White
R. J. H. Brown	I. Griffiths	N. B. Marshall	H. P. Whiting
G. H. O. Burgess	R. W. Hayman	J. A. C. Nichol	
P. M. Butler	William Holmes	H. K. Pusey	
D. B. Carlisle	C. Burdon-Jones	W. D. L. Ride	

If this edition is found to be deserving of any commendation, much of it is properly due to them. Some rewrote special pieces, and a few, in addition,

provided illustrations (below which acknowledgement is made). Almost all drew my attention to errors or half-truths, including, I have to confess, a few that I succeeded in introducing myself. On occasions, my benefactors disagreed in a most stimulating way among themselves—principally on minor matters of taxonomy and terminology—and so it is more than ever necessary to add on their behalf the customary disclaimer concerning responsibility. Nobody other than myself has seen more than a comparatively small part of the manuscript.

It is a final pleasure to acknowledge the loan by the British Museum (Natural History) of specimens for illustration, and the constant assistance given me by the Library staffs of University College and, particularly, the Zoological Society of London.

A. J. M.

Victoria

Australia 1960

NOTE ON THE 1964 REPRINT

The 1964 reprint with corrections has been improved by the further criticism of some of the above-named, and additionally by valuable comments from Drs. D. F. Dorward, A. K. Lee, and J. W. Warren.

A. J. M.

Victoria

Australia 1964

CONTENTS

PHYLUM CHORDATA	1
SUB-PHYLUM ADELOCHORDATA (HEMICHORDATA)	5
CLASS ENTEROPNEUSTA	5
External characters and coelom (p. 5); alimentary canal and associated structures (p. 8); stomochord (proboscis or buccal diverticulum) (p. 10); blood-vascular system (p. 10); nervous system (p. 11); locomotion (p. 11); reproductive organs (p. 11); development (p. 11); tornaria (p. 13).	
CLASS PTEROBRANCHIA	14
Blood-vascular system (p. 17); reproduction (p. 17); affinities (p. 20).	
SUB-PHYLUM TUNICATA (UROCHORDATA)	20
CLASS ASCIDIACEA	21
Example of the Class. (The Ascidian or Sea-squirt (<i>Ascidia</i>) (p. 21); body-wall and atrial cavity (p. 22); pharynx (p. 23); alimentary canal (p. 25); blood-vascular system (p. 25); excretion (p. 27); nervous system (p. 27); neural gland (p. 27); reproductive system (p. 27); development and metamorphosis (p. 28).	
General Organisation of Ascidians: General features (p. 34); composite ascidians (p. 36).	
CLASS THALIACEA	36
Orders Doliolida (<i>Cyclomyaria</i>) (p. 36); Pyrosomida (p. 39); Salpida (<i>Desmomyaria</i>) (p. 41).	
CLASS LARVACEA (APPENDICULARIA)	44
General features, affinities (p. 46).	
SUB-PHYLUM ACRANIA (CEPHALOCHORDATA)	47
Example of the Sub-phylum. —The Lancelet (<i>Amphioxus</i>) (<i>Branchiostoma</i>) (p. 47); body-wall (p. 49); skeleton (p. 50); alimentary canal and associated structures (p. 51); atrium (p. 53); ingestion, digestion, and absorption (p. 53); coelom (p. 54); blood-vascular system (p. 54); excretory organs (p. 56); nervous system (p. 57); organs of special sense (p. 59); reproductive system (p. 59); development (p. 60); affinities (p. 67).	
GENERAL INTRODUCTION TO CRANIATE (VERTEBRATE) FORMS	69
Craniate Tissues: epithelial tissue (p. 70); connective tissue (p. 70); adipose tissue (p. 70); blood (p. 70); lymph (p. 71); cartilage (p. 71); bone (p. 71); muscle (p. 71); neural tissue (p. 71); ectodermal derivatives (p. 71); mesodermal derivatives (p. 71); endodermal derivatives (p. 71).	
The Craniate Body: External characters (p. 78); skeleton (p. 88); endoskeleton (p. 89); alimentary canal and associated structures (p. 102); organs of respiration (p. 105); blood-vascular system (p. 108); lymphatics (p. 115); nervous system (p. 116); somatic motor nerves (p. 130); branchial arch nerves (p. 130); sensory organs (p. 135); olfactory organ (p. 137); eye (p. 139); development of the eye (p. 142); apparatus of audition and equilibration (p. 143); endocrine organs (p. 147); urinogenital organs (p. 154); development (p. 160); metamerism (p. 161).	
SUB-PHYLUM CRANIATA (VERTEBRATA)	163

SUPER-CLASS AGNATHA	164
CLASS EUPHANERIDA	167
CLASS HETEROSTRACI (PTERASPIDA)	168
CLASS ANASPIDA	170
CLASS OSTEOSTRACI (CEPHALASPIDA)	171
CCELOLEPIDA	174
CLASS PETROMYZONTIA	175
Example of the Class. —The Lamprey (<i>Petromyzon</i>)	176
External Characters (p. 177); endoskeleton (p. 180); muscles (p. 184); alimentary canal and associated structures (p. 184); respiratory organs (p. 185); circulatory system (p. 186); nervous system (p. 187); organs of special sense (p. 190); urinogenital system (p. 193); development (p. 194).	
CLASS MYXINOIDEA	197
Comparison of the Myxinoidea with the Lamprey	198
SUPER-CLASS GNATHOSTOMATA	203
CLASS PLACODERMI (APHETOHYOIDEA)	205
Sub-class Acanthodii	207
Sub-class Arthrodira	210
Sub-class Petalichthyida	212
Sub-class Antiarchi (Pterichthyomorphi)	214
Sub-class Rhenanida	216
Sub-class Palæospondylia	217
CLASS ELASMOBRANCHII (CHONDRICHTHYES) (SELACHII)	219
Orders Cladotelachii (Pleuropterygii) (p. 222); Pleuracanthodii (Ichthyotomi) (p. 224); Protoselachii (p. 224); Euselachii (p. 226).	
Example of the Class. —A Dogfish (<i>Scyliorhinus</i> (= <i>Scyllium</i>), <i>Squalus</i> , <i>Brachælorus</i> , etc.) (p. 229); general external features (p. 229); skeleton (p. 231); alimentary canal and associated structures (p. 237); respiration (p. 239); blood-vascular system (p. 240); nervous system (p. 245); organs of special sense (p. 249); endocrine organs (p. 251); excretion and osmoregulation (p. 252); reproduction (p. 254).	
Sub-class Bradyodonti	256
Order Holocephali	256
External characters (p. 258); endoskeleton (p. 259); alimentary canal (p. 261); respiratory organs, heart, brain (p. 262); urinogenital organs and reproduction (p. 263).	
General Organisation of Elasmobranchs: External characters (p. 265); integument (p. 267); endoskeleton (p. 267); skull (p. 268); electric organs (p. 271); alimentary canal and associated structures (p. 272); respiratory organs (p. 273); blood-vascular system (p. 273); brain (p. 274); organs of special sense (p. 274); audio-equilibration (p. 275); endocrine organs (p. 276); urinogenital organs and reproduction (p. 276); placentation (p. 278); development (p. 279).	
CLASS OSTEICHTHYES	282
Sub-class Actinopterygii	282
Super-order Chondrostei	285
Orders Palæoniscoidea (p. 285); Acipenseroides (p. 288); Polypterini (Cladistia) (p. 290).	
Super-order Holostei	291
Orders Semionotoidea (p. 292); Amioidea (p. 292).	
Super-order Teleostei	293
Soft-rayed orders (p. 296); Intermediate orders (p. 296); Spiny-rayed and allied orders (p. 297).	
Example of the Sub-class. —The Brook or Brown Trout (<i>Salmo fario</i>)	298
External characters (p. 298); skin and exoskeleton (p. 300); endoskeleton (p. 300); muscles (p. 309); coelom (p. 309); alimentary canal and associated structures	

CONTENTS

xiii

(p. 309); nervous system (p. 312); autonomic nervous system (p. 314); organs of special sense (p. 315); osmoregulation and excretion (p. 317); endocrine organs (see p. 346); reproduction (p. 318); development (p. 319).	
General Organisation of Actinopterygians: External form (p. 320); exoskeleton (p. 329); endoskeleton (p. 330); electric organs (p. 334); alimentary canal and associated structures (p. 335); respiratory organs (p. 337); swim-bladder (p. 339); blood-vascular system (p. 342); brain (p. 344); exteroception (p. 344); endocrine organs (p. 346); urinogenital organs (p. 347); reproduction (p. 349); development (p. 352).	
Sub-class Crossopterygii (Choanichthes)	353
Classification (p. 354).	
Order Rhipidistia	354
Sub-orders Osteolepidoti (p. 354); Coelacanthini (Actinistia) (p. 356).	
Order Dipnoi (Dipneusti, Dipneumones)	361
Sub-orders Monopneumona (p. 364); Dipneumona (p. 365).	
Example of the Sub-class. — <i>Protopterus aethiopicus</i>	365
Endoskeleton (p. 367); alimentary canal and associated structures (p. 369); respiratory system (p. 370); blood-vascular system (p. 370); nervous system (p. 374); general metabolism and excretion (p. 375); reproduction (p. 376); development (p. 377).	
General Organisation of Dipnoi	379
CLASS AMPHIBIA	381
Example of the Class. —A Frog (<i>Rana</i>)	389
External characters (p. 389); endoskeleton (p. 391); alimentary canal and associated structures (p. 400); respiratory system (p. 403); blood-vascular system (p. 404); lymphatic system (p. 410); nervous system (p. 410); organs of special sense (p. 412); endocrine glands (p. 415); urinogenital organs (p. 415); reproduction (p. 416); development (p. 418).	
Interrelationship of Amphibian Groups	421
Sub-class Apsidospondyli	421
Super-order Labyrinthodontia	422
Order Ichthyostegalia	423
Order Rhachitomi	424
Order Stereospondyli	425
Order Embolomeri	426
Order Seymouriamorpha	427
Super-order Salientia (Anura, Batrachia)	428
Sub-class Lepospondyli	430
Order Aistopoda	430
Order Nectridia	430
Order Microsauria (Adelospondyli)	431
Order Phyllospondyli	431
Order Urodela (Caudata)	432
Order Apoda (Gymnophiona, Caecilia)	434
General Organisation of Amphibians: External characters (p. 435); exoskeleton (p. 437); endoskeleton (p. 438); musculature (p. 442); alimentary canal and associated structures (p. 442); nervous system and organs of special sense (p. 447); endocrine system (p. 450); urinogenital system and osmoregulation (p. 450); reproduction and development (p. 452).	
CLASS REPTILIA	457
Example of the Class. —A Lizard (<i>Lacerta</i>)	462
Exoskeleton (p. 463); endoskeleton (p. 463); alimentary canal and associated structures (p. 471); blood-vascular system (p. 474); respiratory system (p. 477); brain (p. 478); endocrine glands (p. 481); organs of special sense (p. 481); urinogenital system (p. 482).	
Interrelationship of Reptilian Groups	484

Sub-class Anapsida	486
Order Cotylosauria	486
Sub-orders Captorhinomorpha (p. 486); Diadectomorpha (p. 486).	
Order Chelonia	486
Sub-class Ichthyopterygia	490
Order Mesosauria	490
Order Ichthyosauria	491
Sub-class Synaptosauria	492
Order Protorosauria	492
Order Sauropterygia	492
Sub-orders Nothosauria (p. 492); Placodontia (p. 493); Plesiosauria (p. 493).	
Sub-class Lepidosauria	495
Order Eosuchia	495
Order Rhynchocephalia	495
Squamata	496
Sub-orders Lacertilia (p. 496); Ophidia (Serpentes) (p. 497).	
Sub-class Archosauria	498
Order Thecodontia	499
Order Crocodilia (Loricata)	499
Order Pterosauria (Pterodactyla)	502
Dinosaurs	504
Order Saurischia	505
Sub-orders Theropoda (p. 505); Sauropoda (p. 507).	
Order Ornithischia	507
Sub-orders Ornithopoda (p. 508); Stegosauria (p. 510); Ankylosauria (p. 511); Ceratopsia (p. 511).	
Sub-class Synapsida	513
Order Pelycosauria	513
Order Therapsida	514
Order Ictidosauria	517
General Organisation of Recent Reptiles: External features (p. 518); integument and exoskeleton (p. 522); endoskeleton (p. 524); alimentary canal and associated structures (p. 533); respiratory system and voice (p. 540); blood-vascular system (p. 541); nervous system (p. 542); organs of special sense (p. 543); endocrine glands (p. 548); excretion (p. 548); reproduction (p. 549); development (p. 551).	
CLASS AVES	555
Example of the Class.—The Common Pigeon (<i>Columba livia</i> , var. <i>domestica</i>)	565
External characters (p. 566); exoskeleton (p. 567); endoskeleton (p. 571); muscular system (p. 580); locomotion (p. 582); alimentary canal and associated structures (p. 583); respiratory and vocal organs (p. 585); thyroid, parathyroid, and thymus glands (p. 589); blood-vascular system (p. 589); lymphatic system (p. 592); nervous system and hypophysis (p. 592); organs of special sense (p. 594); excretory and reproductive organs, and adrenal glands (p. 596).	
General Organisation of Birds	598
Sub-class Archæornithes	599
Order Archæopterygiformes	599
Sub-class Neornithes	602
Super-order Odontognathæ	603
Orders Hesperornithiformes (p. 604); Ichthyornithiformes (p. 605).	
Super-order Palæognathæ	606
Orders Struthioniformes (p. 607); Rheiformes (p. 607); Casuariiformes (p. 607); Apterygiformes (p. 608); Dinornithiformes (p. 608); Zepyornithiformes (p. 609); Tinamiformes (p. 609).	
Super-order Impennæ	610
Order Sphenisciformes	610
Super-order Neognathæ	612
Orders Gaviiformes (p. 613); Podicipitiformes (p. 613); Procellariiformes (p. 613); Pelicaniiformes (p. 614); Ciconiiformes (p. 614); Anseriformes (p. 614); Falconi-	

CONTENTS

xv

formes (p. 614); **Galliformes** (p. 615); **Gruiformes** (p. 615); **Diatrymiformes** (p. 616); **Charadriiformes** (p. 616); **Columbiformes** (p. 617); **Psittaciformes** (p. 617); **Cuculiformes** (p. 617); **Strigiformes** (p. 618); **Caprimulgiformes** (p. 618); **Apodiformes (Micropodiformes)** (p. 618); **Coliiformes** (p. 618); **Trogoniformes** (p. 619); **Coraciiformes** (p. 619); **Piciformes** (p. 619); **Passeriformes** (p. 620).
General Organisation of Neornithes: External characters (p. 620); pterylosis (p. 626); endoskeleton (p. 628); alimentary canal and associated structures (p. 636); respiratory and vocal organs (p. 638); circulatory organs (p. 639); nervous system and organs of special sense (p. 639); excretion (p. 643); reproduction (p. 644); development (p. 646).

CLASS MAMMALIA	652
Example of the Class. —The Rabbit (<i>Oryctolagus cuniculus</i>)	654
External characters (p. 654); skeleton (p. 655); cœlom (p. 666); alimentary canal and associated structures (p. 666); blood-vascular system (p. 670); lymphatic system (p. 675); respiratory system (p. 676); nervous system (p. 678); spinal cord, cranial and spinal nerves (p. 682); autonomic nervous system (p. 682); organs of special sense (p. 684); urinogenital organs (p. 684); development (p. 686).	
Sub-class Prototheria	687
Order Triconodonta	687
Order Symmetrodonta	687
Order Monotremata	688
Skeleton (p. 696); visceral anatomy (p. 700).	
Sub-class Allotheria	701
Order Multituberculata	701
Sub-class Theria	703
Infra-class Pantotheria (Trituberculata)	703
Order Dryolestoides	703
Order Docodonta	703
Infra-class Metatheria (Marsupialia, Didelphia)	704
Super-families Didelphoidea (p. 711); Borhyaenoidea (p. 712); Dasyuroidea (p. 712); Perameloidea (p. 715); Cænolestoides (p. 715); Phalangeroidea (p. 716). Endoskeleton of Metatheria (p. 719).	
Infra-class Eutheria (Placentalia, Monodelphia)	723
Introduction (p. 723).	
Cohort Unguiculata	726
Order Lipotyphla	726
Skeleton of the Lipotyphla (p. 730).	
Order Menotyphla	731
Order Primates	734
Sub-order Prosimii	736
Infra-orders Lemuriformes (p. 737); Lorisiformes (p. 738); Tarsiiformes (p. 739).	
Sub-order Anthropeoidea	741
Super-family Ceboidea (Platyrrhina)	743
Super-family Cercopithecoidea	743
Super-family Hominoidea	744
Families Pongidæ (p. 744); Hominidæ (p. 745). Skeleton of the Primates (p. 748).	
Order Dermoptera	753
Order Chiroptera	754
Sub-orders Megachiroptera (p. 756); Microchiroptera (p. 756). Skeleton of the Chiroptera	758
Order Tæniodontia	760
Order Tillodontia	760
Order Edentata	760
Sub-orders Palæanodontia (p. 761); Xenarthra (p. 761). Skeleton of Xenarthra (p. 765).	

Order Pholidota	772
Cohort Glires	774
Order Rodentia	774
Skeleton of the Rodentia (p. 778).	
Order Lagomorpha	779
Cohort Mutica	779
Order Cetacea	780
Sub-orders Archæoceti (p. 782); Odontoceti (p. 783); Mysticeti (p. 784).	
Skeleton of Cetacea (p. 785); internal adaptations of Cetacea (p. 789).	
Cohort Ferungulata	791
Super-order Feræ	791
Order Carnivora	791
Sub-orders Creodonta (p. 792); Fissipedia (p. 793); Pinnipedia (p. 797).	
Skeleton of the Carnivora (p. 798).	
Super-order Protoungulata	801
Order Condylarthra	802
Order Tubulidentata	804
South American 'Ungulata'	805
Order Litopterna	805
Order Notoungulata	806
Order Astrapotheria	807
Super-order Pænungulata (Subungulata)	807
Order Hyracoidea	808
Order Embrithopoda	810
Order Proboscidea	811
Orders Pantodonta (Amblypoda) and Dinocerata	818
Order Pyrotheria	820
Order Sirenia	820
Skeleton of Sirenia (p. 823).	
Super-order Mesaxonia	824
Order Perissodactyla	825
Families Equidæ (p. 826); Brontotheriidae (Titanotheriidae) (p. 831); Chalicotheriidae (p. 832).	
Sub-order Ceratomorpha	833
Super-family Tapiroidea	833
Super-family Rhinocerotioidea	834
Super-order Paraxonia	836
Order Artiodactyla	836
Sub-orders Suiformes (p. 837); Tylopoda (p. 838); Ruminantia (p. 839).	
General Organisation of Mammalia: Integument and general external features (p. 841); endoskeleton (p. 850); alimentary canal and associated structures (p. 862); blood-vascular system (p. 881); lymphatic system (p. 883); respiratory system (p. 884); nervous system (p. 884); organs of special sense (p. 887); endocrine organs (p. 888); urinogenital system (p. 888); breeding seasons (p. 896); development and associated phenomena (p. 898); post-natal behaviour in marsupials (p. 907).	

REFERENCES	909
General (p. 909); Protochordates (p. 914); Agnathans (p. 915); Placoderms and Fishes (p. 917); Amphibians (p. 920); Reptiles (p. 921); Birds (p. 921); Mammals (p. 923).	

INDEX	927
-----------------	-----

TABLES

I. Ages (in million of years) of geological periods	3
II. Relationship of somites, muscles, nerves and visceral arches	134
III. Divisions of hypophysis	151
IV. Composition of milk	849

ILLUSTRATIONS

SUB-PHYLUM ADELOCHORDATA

CLASS ENTEROPNEUSTA *Balanoglossus*, Whole animal, Fig. 1 (p. 6); in burrow, Fig. 2 (p. 7); median sagittal section, Fig. 3 (p. 8); branchial region, Fig. 4 (p. 9); development, Fig. 5 (p. 12); Fig. 6 (p. 13); tornaria larva, Fig. 7 (p. 14).

CLASS PTEROBRANCHIA *Cephalodiscus*, Colony, Fig. 8 (p. 15); zooid, Fig. 9 (p. 16); visceral relationships, Fig. 10 (p. 17); *Atubaria*, Fig. 11 (p. 18); *Rhabdopleura*, Colony, Fig. 12 (p. 18); male zooid and cœnœcium, Fig. 13 (p. 19).

SUB-PHYLUM TUNICATA (UROCHORDATA)

CLASS ASCIDIACEA *Ascidia*, whole animal, Fig. 15 (p. 23); branchial sac, Fig. 16 (p. 24); pharyngeal region, Fig. 17 (p. 24); circulation and reproduction, Fig. 18 (p. 25); nerve ganglion and associated parts, Fig. 19 (p. 27); nervous system, Fig. 20 (p. 28). Ascidian development: Figs. 21–25 (pp. 29–33). Composite ascidians: *Botryllus*, Fig. 26 (p. 35); a colonial zooid, Fig. 27 (p. 35).

CLASS THALIACEA *Doliolum*, whole animal, Fig. 28 (p. 37); development, Figs. 29–30 (p. 38); asexual zooid, Fig. 31 (p. 39). *Pyrosoma*, Luminous colony, Fig. 32 (p. 40); colonial individuals, Fig. 33 (p. 40). *Salpa*, Asexual form, Fig. 34 (p. 41); oozoid, Fig. 35 (p. 42); nervous system, Fig. 36 (p. 43); development, Fig. 37 (p. 43).

CLASS LARVACEA *Oikopleura*, Animal in 'house', Fig. 38 (p. 45); *Appendicularia*, visceral relationships, Fig. 39 (p. 45).

SUB-PHYLUM ACRANIA

CLASS CEPHALOCHORDATA *Branchiostoma* (*Amphioxus*), Whole animal in sand, Fig. 40 (p. 48); ventral and lateral veins, Fig. 41 (p. 48); visceral relationships, Fig. 42 (p. 50); general anatomy, Fig. 43 (p. 51); pharyngeal region, Fig. 44 (p. 54); blood vascular system, Fig. 45 (p. 55); nephridium, Fig. 46 (p. 56); solenocytes, Fig. 47 (p. 57); nervous system, Fig. 48 (p. 58); development Figs. 49–56 (pp. 60–66).

SUB-PHYLUM CRANIATA (VERTEBRATA)

Introductory Examples: General and Skeletal. 'Ideal' vertebrate, external view, Fig. 57 (p. 79); epidermis, Fig. 58 (p. 82); placoid scale, Fig. 59 (p. 83); cosmoid scale, Fig. 60 (p. 83); ganoid scale, Fig. 61 (p. 84); myomere patterns, Fig. 62 (p. 86); visceral relationships, Fig. 63 (p. 87); vertebral column in embryo, Fig. 64 (p. 89); segmentation, Fig. 65 (p. 89); cranial elements in embryonic *Salmo*, Fig. 66 (p. 90); cartilaginous skull, Fig. 67 (p. 92); bony skull, Fig. 68 (p. 95); pelvic fins, Fig. 69 (p. 98); pectoral fins, Fig. 70 (p. 99); tetrapod girdles, manus, and pes, Fig. 71 (p. 100); dermal and buccal armour; placoid scale and teeth, Fig. 72 (p. 103).

Respiratory System. Pharyngeal region, including branchiogenic organs, Fig. 73 (p. 106); pulmonary lobule, Fig. 74 (p. 107); lungs and swim-bladder, Fig. 75 (p. 108).

Blood-vascular System. In fishes, Fig. 76 (p. 109); course of circulation, Fig. 77 (p. 110); circulation in embryonic air-breather, Fig. 78 (p. 113); cardiac circulation in amphibian and crocodile, Fig. 79 (p. 114); relationship of blood and lymph capillaries, Fig. 80 (p. 116).

Nervous System and Special Senses. Brain and proximal part of cord during development and maturity, Fig. 81 (p. 118); autonomic system, Fig. 82 (p. 119); sympathetic and spinal nerves and cord, Fig. 83 (p. 120); fibre, motor end-plate and muscle, Fig. 84 (p. 121); cord and constituent structures, Fig. 85 (p. 123); cranial and anterior spinal nerves, Fig. 86 (p. 129); lateral-line system in bony, Fig. 87 (p. 136) and cartilaginous fishes, Fig. 88 (p. 137); nasal and buccal cavity in various vertebrates, Fig. 89 (p. 138); eye (macroscopic structures), Fig. 90 (p. 139); retinal structures, Fig. 91 (p. 140); orbit, Fig. 92 (p. 143); audio-equilibration: evolutionary stages, Fig. 93 (p. 144); semi-circular canals and lagena, Fig. 94 (p. 145).

Endocrine System. Hypothalamico-hypophysial relationships, Fig. 95 (p. 149); hypophysis arrangements, Fig. 96 (p. 150); secretory apparatus of testis, Fig. 97 (p. 152).

Urinogenital System. Development and relationships, Fig. 98 (p. 153); kidney circulation and disposal, Fig. 99 (p. 154); nephron, Fig. 100 (p. 155); and probable stages of evolution, Fig. 101 (p. 157); development, Fig. 102 (p. 161).

SUPER-CLASS AGNATHA

CLASS EUPHANERIDA *Jamoytius*, Fig. 103 (p. 167).

CLASS HETEROSTRACI *Pteraspis*, Fig. 104 (p. 168); *Drepanaspis*, Fig. 105 (p. 170).

CLASS ANASPIDA *Rhyncholepis*, Fig. 106 (p. 171).

CLASS OSTEOSTRACI *Hemicyclaspis*, Fig. 107 (p. 171); *Kiaeraspis*, cranial anatomy, Fig. 108 (p. 173); *Lanarkia*, Fig. 109 (p. 174).

CLASSES PETROMYZONTIA and MYXINOIDEA *Petromyzon*, whole animal, Fig. 110 (p. 175); buccal funnel, Fig. 111 (p. 177); integumental and epidermal cells of *Myxine* and fishes, Figs. 112–113 (pp. 178–179); *Petromyzon*, trunk region, Fig. 114 (p. 180); skull, Fig. 115 (p. 181); skull and branchial basket in adult, Fig. 116 (p. 182); skull during metamorphosis, Fig. 117 (p. 182); head region, Fig. 118 (p. 183); *Ammocete* (*Geotria*), head region, Fig. 119 (p. 186); *Petromyzon*, brain, Fig. 120 (p. 188); olfactory and naso-hypophysial ducts, Fig. 121 (p. 189); development, Fig. 122 (p. 190); *Lampetra*, eye and adjacent structures, Fig. 123 (p. 191); *Petromyzon*, cloacal region, Fig. 125 (p. 193); development, Fig. 126 (p. 195); *Lampetra*: (development, Fig. 127 (p. 196); *Myxine* and *Eptatretus*, anterior regions, Fig. 128 (p. 198); *Myxine* skeletal and visceral structures, Fig. 129 (p. 199); skull of adult, Fig. 130 (p. 200); of embryo, Fig. 131 (p. 201); membranous labyrinth, Fig. 132 (p. 202).

SUPER-CLASS GNATHOSTOMATA

CLASS PLACODERMI *Dinichthys*, whole animal, Fig. 133 (p. 206); *Climatius*, head, Fig. 134 (p. 207); *Diplacanthus*, pectoral girdle, Fig. 135 (p. 208); *Acanthodes*, skull, Fig. 136 (p. 209); head region, Fig. 137 (p. 210); *Coccosteus*, skeleton, Fig. 138 (p. 211); *Macropetalichthys*, skeleton, Fig. 139 (p. 212); head and pectoral shield of *Lunaspis* and *Macropetalichthys*, Fig. 140 (p. 213); *Macropetalichthys*, pectoral girdle, Fig. 141 (p. 213); *Pterichthyodes*, skeleton, Fig. 142 (p. 214); *Bothriolepis*, whole animal, Fig. 143 (p. 215); *Gemuendina*, whole animal, and trunk, Figs. 144–145 (pp. 216–217); *Palaeospondylus*, skeleton, Fig. 146 (p. 218).

CLASS ELASMOBRANCHII *Carcharodon*, jaws and teeth, Fig. 147 (p. 220); *Cladoseleache*, whole animal, Fig. 148 (p. 223); *Ctenacanthus*, whole animal, Fig. 149 (p. 223); *Pleuranacanthus*, skeleton, Fig. 150 (p. 224); *Heterodontus*, whole animal, Fig. 151 (p. 225); dentition, Fig. 152 (p. 226); *Chlamydoselachus*, Fig. 153 (p. 227); *Pristis* and *Aetobatis* illustrating adaptation and radiation, Fig. 154 (p. 228).

Example of Class: Dogfish. Whole animal, Fig. 155 (p. 229); vertebral column, Fig. 156 (p. 231); skull and visceral arches, Fig. 157 (p. 232); visceral arches, Fig. 158 (p. 234); pectoral arch and fin, Fig. 159 (p. 235); pelvic arch and fin, Fig. 160 (p. 236); visceral structures, Fig. 161 (p. 238); Respiration and circulation: branchial sac, Fig. 162 (p. 239); head and branchial circulation, Fig. 163 (p. 240); general circulation, Fig. 164 (p. 242); Nervous system: brain, Fig. 165 (p. 244); brain ventricles, Fig. 166 (p. 246); brain and spinal nerves, Fig. 167 (p. 247); dorsal root cranial nerves, Fig. 168 (p. 248). Endocrine and urinogenital elements: adrenal, Fig. 169 (p. 252); male and female urinogenitalia, Fig. 170 (p. 253); kidney and urinary sinus of female, Fig. 171 (p. 254); egg-case of dogfish, Fig. 172 (p. 254).

SUB-CLASS BRADYODONTI Representative whole animals, Fig. 173 (p. 257); *Chimæra*, vertebral column, Fig. 174 (p. 259); skull, Fig. 175 (p. 260). *Callorhynchus*, skull, Fig. 176 (p. 261); brain, Fig. 177 (p. 262); urinogenital system, Fig. 178 (p. 263); embryo within egg, Fig. 179 (p. 264).

General Organisation of Elasmobranchii. Skeletal structures: *Scymnorhinus*, spinal column, Fig. 180 (p. 267); *Urolophus*, endoskeleton, Fig. 181 (p. 268); *Heptranchias*, skull, Fig. 182 (p. 269). *Torpedo*, Electric organs, Fig. 183 (p. 271); *Carcharodon*, Mechanism of eye accommodation, Fig. 184 (p. 273); *Cetorhinus*, Urinogenital system, Fig. 185 (p. 277); development, Figs. 186–190 (pp. 279–281).

CLASS OSTEICHTHYES SUPER-ORDER CHONDOSTREI *Palæoniscus*, whole animal and skeleton, Fig. 191 (p. 286); *Helichthys*, whole animal and skeleton, Fig. 192 (p. 287); *Acipenser*, whole animal, Fig. 193 (p. 288); pre-caudal vertebræ, Fig. 194 (p. 289). *Chondrosteus*, skull and pectoral girdle, Fig. 195 (p. 289); *Polypterus*, whole animal, Fig. 196 (p. 290); **SUPER-ORDER HOLOSTEI**, *Dapedius*, whole animal, Fig. 197 (p. 291); *Lepisosteus*, whole animal, Fig. 198 (p. 292); *Amia*, whole animal, Fig. 199 (p. 293). **SUPER-ORDER TELEOSTEI**, *Leptolepis*, whole animal and skull, Fig. 200 (p. 294).

Example of Sub-class: *Salmo fario*, Trout. Whole animal, Fig. 201 (p. 296); *Tilapia*, whole animal and parental care, Fig. 202 (p. 297); *Salmo*, vertebræ, Fig. 203 (p. 301). **Teleostei**, caudal skeleton, Fig. 204 (p. 302); skull and relationship with pectoral girdle, Fig. 205 (p. 304); *Salmo*, skull of juvenile, Fig. 206 (p. 307); dermal fin-ray and its supports, Fig. 207 (p. 307); pectoral girdle and fin, Fig. 208 (p. 308); pelvic fin and skeleton, Fig. 209 (p. 309); skeletal and visceral relationships, Fig. 210 (p. 310); brain and cranial nerves, Fig. 211 (p. 313); brain and adjacent structures, Fig. 212 (p. 314); **Teleostei**, eye and optic nerve, Fig. 213 (p. 316); *Salmo*, audio-equilibration, Fig. 214 (p. 317); kidney and adjacent structures, Fig. 215 (p. 318); development, Figs. 216–217 (p. 319).

General Organisation of Actinopterygians.—**Teleostei**, optic adaptation, Fig. 218 (p. 322); exoskeletal and reproductive adaptations, Fig. 219 (p. 323); bioluminescence and angling mechanisms, Fig. 220 (p. 324); morphogenetic adaptation to bottom-living, Fig. 221 (p. 326); bioluminescence in *Argyrops*, Fig. 222 (p. 328); tactile and other specialisations, Fig. 223 (p. 328); Actinopterygian scales (cycloid and ctenoid), Fig. 224 (p. 329). **Acipenseroidæ**, skull, Fig. 225 (p. 330); **Polypterini**, skull, Fig. 226 (p. 331). **Teleostei**, deglutition specialisations, Fig. 227 (p. 332); **Polypterini**, fins, Figs. 228–229 (p. 333); **Teleostei**, electric organs, Fig. 230 (p. 334); dental specialisation in *Sargus*, Fig. 231 (p. 336); extra-branchial respiration in *Clarias*, Fig. 232 (p. 338); in *Anabas*, Fig. 233 (p. 339). **Holostei**, swim-bladder in *Lepisosteus*, Fig. 234 (p. 340). **Teleostei**, swim-bladder and audio-equilibration in a cypriinoid, Fig. 235 (p. 341); gas gland, oval and blood supply, Fig. 236 (p. 342). **Holostei**, brain, Fig. 237 (p. 344); **Teleostei**: brain and labyrinth, Fig. 238 (p. 345). **Holostei**: urinogenital systems, Figs. 239–240 (p. 348); **Teleostei**: ectoparasitism and reproduction, Fig. 241 (p. 350); viviparity, Fig. 242 (p. 351); **Holostei**, development, Fig. 243 (p. 353).

SUB-CLASS CROSSOPTERYGII

Order Rhipidistia: *Osteolepis*, whole animal and skull, Fig. 244 (p. 355); *Undina*, whole animal, Fig. 245 (p. 357); *Latimeria*, whole animal, Fig. 246 (p. 358); head and sensory organs, Figs. 247–248 (p. 359).

Order Dipnoi: Lung-fishes, whole animals and distribution, Fig. 249 (p. 361); *Dipterus*, whole animal, Fig. 250 (p. 363).

Example of Sub-class.—*Protopterus æthiopicus*, whole animal, Fig. 251 (p. 365); skull, shoulder girdle, and fin, Fig. 252 (p. 367); Dentition, Fig. 253 (p. 368); alimentary canal and associated structures, Fig. 254 (p. 369); lung cf. that of *Polypterus*, Fig. 255 (p. 371); blood vascular and respiratory systems, Fig. 256 (p. 372); visceral relationships, Fig. 257 (p. 373); brain, Fig. 258 (p. 374); æstivation, Fig. 259 (p. 375); urinogenital system, Fig. 260 (p. 376); development, Figs. 261–262 (pp. 377–378).

CLASS AMPHIBIA Fish/amphibian transition: shoulder girdle and fore-limb, Fig. 263 (p. 382).

Example of Class: *Rana*, endoskeleton, Fig. 264 (p. 392); bony skull, Fig. 265 (p. 393); cartilaginous skull of tadpole, Fig. 266 (p. 394); pectoral girdle, Fig. 267 (p. 397); pelvic girdle, Fig. 268 (p. 398); musculature, Fig. 269 (p. 399); visceral relationships, Fig. 270 (p. 401); abdominal viscera, Fig. 271 (p. 402); cardiac anatomy and circulation, Fig. 272 (p. 405); arterial system, Fig. 273 (p. 406); venous system, Fig. 274 (p. 408); brain, pituitary and cranial nerves, Fig. 275 (p. 411); eye and optic nerve, Fig. 276 (p. 413); urinogenital system, Figs. 277–278 (pp. 416–417); development, Fig. 279 (p. 419).

SUB-CLASS APSIDOSPONDYLI

Labyrinthodontia: tooth, Fig. 280 (p. 422); skulls and vertebræ, Fig. 281 (p. 422); *Ichthyostega*, whole animal, Fig. 282 (p. 423); cranium, Fig. 283 (p. 424); *Paracyclotosaurus*, whole animal, Fig. 284 (p. 425); skulls of *Palæogyrinus* and the fish *Osteolepis* showing evidence of fish/amphibian transition, Fig. 285 (p. 427); *Seymouria*, illustrating mosaic evolution, Fig. 286 (p. 429).

SUB-CLASS LEOSPONDYLI

'Horned' skull, and characteristic vertebra, Fig. 287 (p. 431); *Ophiderpeton*, skeleton, Fig. 288 (p. 431); **Order Urodela:** *Andrias*, Witness of the Deluge, Fig. 289 (p. 432); permanent larvæ, Fig. 290 (p. 433); *Salamandra salamandra*, Fig. 291 (p. 434). **Order Apoda:** Cephalic and genital structures of *Cæcilia*, Fig. 292 (p. 434).

General Organisation of the Amphibia.—**Urodela:** chondrocranium of *Proteus*, Fig. 293 (p. 438); skull of *Salamandra atra*, Fig. 294 (p. 439); **Apoda:** skull of *Ichthyophis*, Fig. 295 (p. 440); **Urodela:** shoulder girdle and sternum, Fig. 296 (p. 441); pelvic girdle, Fig. 297 (p. 442); **Anura:** vocal apparatus, Fig. 298 (p. 443); **Urodela:** circulation and respiration, Fig.

299 (p. 445); venous system, Fig. 300 (p. 446); vibration exteroception, Fig. 301 (p. 448); **Anura**: lateral line in *Xenopus*, Fig. 302 (p. 449); melanophores in *Xenopus*, Fig. 303 (p. 450); **Urodela**: urinogenitalia, Figs. 304–305 (pp. 450–451); **Anura**: amplexus and parental care in *Gastrotheca marsupiat*, Fig. 306 (p. 453); ovoviviparity and pseudo-placentation in *Pipa dorsigera*, Figs. 307–308 (pp. 454–455); **Apoda**: reproduction and development in *Ichthyophis*, Fig. 309 (p. 456).

CLASS REPTILIA

Skeleton of anapsid 'stem-reptile', *Labidosaurus*, Fig. 310 (p. 458).

Example of Class: Lacerta, vertebrae, Fig. 311 (p. 463); pectoral arch and sternum, Fig. 312 (p. 465); skull, Fig. 313 (p. 466); carpus, Fig. 314 (p. 469); pelvis, Fig. 315 (p. 470); tarsus, Fig. 316 (p. 471); visceral relationships, Figs. 317–318 (pp. 472–473); heart and large vessels, Fig. 319 (p. 475); cardiac structure and circulation, Fig. 320 (p. 476); brain and cranial nerves, Fig. 321 (p. 479); brain and hypophysis, Fig. 322 (p. 480); audio-equilibration, Fig. 323 (p. 481); urinogenitalia, Figs. 324–325 (pp. 482–483); skull structure of reptiles, Fig. 326 (p. 484),

SUB-CLASS ANAPSIDA

Chelonia, whole animal, Fig. 327 (p. 487).

SUB-CLASS ICHTHYOPTERYGIA *Ophthalmosaurus*, whole animal, Fig. 328 (p. 490); locomotory specialisations, Fig. 329 (p. 491).

SUB-CLASS SYNAPTOSAURIA *Elasmosaurus*, whole animal, Fig. 330 (p. 493).

SUB-CLASS LEPIDOSAURIA *Rhynchocephalia*, *Sphenodon*, whole animal, Fig. 331 (p. 496); *Squamata*, *Crotalus*, whole animal, Fig. 332 (p. 498).

SUB-CLASS ARCHOSAURIA *Crocodylus*, Fig. 333 (p. 500); *Rhamphorhynchoidea*, skull and teeth, Fig. 335 (p. 504); *Saurischia* and *Ornithischia*: Representative dinosaurs, Fig. 336 (p. 505); pelvic structure, Fig. 337 (p. 506); *Saurischia*, axial skeleton of *Tyrannosaurus*, Fig. 338 (p. 506); respiratory adaptations in dinosaurs, Fig. 339 (p. 508); *Ornithischia*, endoskeleton of *Iguanodon*, Fig. 340 (p. 509); dentition in unrelated herbivores, Fig. 341 (p. 510); brain, hypophysis and spinal enlargements in *Stegosaurus*, Fig. 342 (p. 511).

SUB-CLASS SYNAPSIDA *Pelycosauria*; *Dimetrodon*, whole animal, Fig. 343 (p. 512); *Edaphosaurus*, vertebral spine, Fig. 344 (p. 513); *Therapsida*, *Kannemeyeria*, skeleton, Fig. 345 (p. 514); pectoral girdle, Fig. 346 (p. 515). *Cynodontia*, skull, Fig. 347 (p. 516); *Gorgonopsia*, skull, Fig. 348 (p. 516); *Lycæonops*, whole animal, Fig. 349 (p. 517); *Ictidosauria*, skull, Fig. 350 (p. 517).

General Organisation of Recent Reptilia: Lacertilia, arboreal specialisation in *Chamaeleon*; Fig. 351 (p. 519); Fossorial specialisations in legless lizard, *Pygopus*, Fig. 352 (p. 520); *Sphenodon*, vertebrae, Fig. 353 (p. 524); *Python*, vertebrae, Fig. 354 (p. 524); *Crocodylus* and *Sphenodon*, endoskeleton, Figs. 355–356 (p. 525); *Crocodylus*, vertebrae, Fig. 357 (p. 526); *Chelonia*, exo- and endo-skeleton, Figs. 358–359 (p. 527); *Natrix*, skull, Fig. 360 (p. 528); *Sphenodon*, skull, Fig. 361 (p. 529); *Chelonia*, skull, Fig. 362 (p. 530); *Crocodylus*, skull of adult and juvenile, Figs. 363–364 (p. 531); *Chelonia*, tarsus, Fig. 365 (p. 532); *Alligator*, carpus, Fig. 366 (p. 532); pelvis, Fig. 367 (p. 533); *Crocodylus*, Fig. 368 (p. 533); *Crotalus*, rattlesnake fangs, poison apparatus, Figs. 369–370 (pp. 534–535); Elapid dental specialisations, Fig. 371 (p. 536); *Crocodylus*, heart and great vessels, Fig. 372 (p. 542); *Alligator*, brain and cranial nerves, Fig. 373 (p. 543); *Lacertilia*, vomeronasal organ, Fig. 374 (p. 544); *Squamata*, comparison of lacertilian and ophidian eye, Fig. 375 (p. 545); *Sphenodon* pineal apparatus, Fig. 376 (p. 546); *Crotalinae*, special senses, Fig. 377 (p. 547); *Ophidia*, bilateral copulatory organs, Fig. 378 (p. 550); *Alligator*, development, Fig. 379 (p. 552); *Lacerta*, development, Fig. 380 (p. 553).

CLASS AVES.

Epigamic plumage, Fig. 381 (p. 561); sexual and aggression display, Fig. 382 (p. 562).

Example of Class: Columba livia, Fig. 383 (p. 566); contour feather, Fig. 384 (p. 568); down-feather and its formation, Fig. 385 (p. 570); pterylosis, Fig. 386 (p. 571); internal strutting and pneumatism in bone, Fig. 387 (p. 572); endoskeleton, Fig. 388 (p. 572); cervical vertebra, Fig. 389 (p. 573); synsacrum, Fig. 390 (p. 574); skull, Fig. 391 (p. 575); *Neornithes*, skull, Fig. 392 (p. 576); hyoid apparatus, Fig. 393 (p. 576); columella, Fig. 394 (p. 577); fore-limb, Fig. 395 (p. 577); manus, Fig. 396 (p. 578); pelvic girdle, Fig. 397 (p. 578); hind-limb, Fig. 398 (p. 579); pes, Fig. 399 (p. 580); musculature, Fig. 400 (p. 581); visceral relationships, Fig. 401 (p. 584); trachea, bronchi, and lungs, Fig. 402 (p. 586); air-sac relationships, Fig. 403 (p. 587); lungs and air-sacs, Fig. 404 (p. 588); heart and large vessels, Fig. 405 (p. 589); blood vascular system; Fig. 406 (p. 590); brain and cranial nerves, Fig. 407 (p. 592); brain, Fig. 408 (p. 593); eye, Fig. 409 (p. 594); audio-equilibration, Fig. 410 (p. 595); male urinogenitalia, Fig. 411 (p. 596); female urinogenitalia, Fig. 412 (p. 596); cloacal region, Fig. 413 (p. 597).

SUB-CLASS ARCHÆORNITHES

Archæopteryx (and the reptile **Rhamphorhynchus**, Fig. 414 (p. 599); **Archæopteryx** (Berlin specimen *in situ*), Fig. 415 (p. 600); skull, Fig. 416 (p. 601); manus, Fig. 417 (p. 601).

SUB-CLASS NEORNITHES

Hesperornis and **Ichthyornis**, Fig. 418 (p. 603); skeletons, Figs. 419–420 (pp. 604–605); **Dinornis** and **Apteryx**, Fig. 421 (p. 608); **Eudromia (Tinamu)**, Fig. 422 (p. 609); **Aptenodytes** and parental care, Fig. 423 (p. 611); **Map**: adaptation and geographical distribution, Fig. 424 (p. 612); **Diatryma**, Fig. 425 (p. 616).

General Organisation of Neornithes: Rostral and associated specialisations, Fig. 426 (p. 621); clawed wings of, **Opisthocomus**, Fig. 427 (p. 622); pedal specialisations, Fig. 428 (p. 623); adaptation to lily lagoons, Fig. 429 (p. 624); toilet comb (of bittern, **Botaurus**, Fig. 430 (p. 625); **Casuaris**, feather construction, Fig. 431 (p. 627); avian sternum and pectoral articulation, Fig. 432 (p. 629); **Apteryx**, skull, Figs. 433–434 (pp. 630–631); **Anas**, skull, Fig. 435 (p. 631); **Psittacidae**, cranio-facial hinge, Fig. 436 (p. 632); **Apteryx**, shoulder girdle, Fig. 437 (p. 633); avian embryonic fore-limb showing digits, Fig. 438 (p. 634); **Apteryx**, pelvic girdle, Fig. 439 (p. 634); hind-limb and pes, Fig. 440 (p. 635); **Cecinus** (woodpecker), feeding specialisations, Fig. 441 (p. 636); **Collocalia** (swiftlet), salivary glands, Fig. 442 (p. 637); **Grus** (crane), vocal specialisations, Fig. 443 (p. 638); **Cyanochen** (goose), vocal specialisations, Fig. 444 (p. 639); **Botaurus** (bittern), adaptation to reed-beds, Fig. 445 (p. 641); avian tactile bristles, Fig. 446 (p. 643); development, Figs. 447–451 (pp. 646–650).

CLASS MAMMALIA

Example of Class: Oryctolagus cuniculus (Rabbit). Vertebrae, Fig. 452 (p. 656); skull, Figs. 453–454 (pp. 658–659); jaw musculature, Fig. 455 (p. 662); fore-arm and carpus, Fig. 456 (p. 664); sacrum, Fig. 457 (p. 665); pes, Fig. 458 (p. 666); head, neck, and thorax, Fig. 459 (p. 667); nasal region, Fig. 460 (p. 668); visceral relationships, Fig. 461 (p. 669); heart and great vessels, Fig. 462 (p. 671); blood vascular system, Fig. 463 (p. 673); cardiac region and thoracic duct (of *Homo*), Fig. 464 (p. 676); larynx, Fig. 465 (p. 677); thoracic relationships, Fig. 466 (p. 677); brain, Figs. 467–468 (pp. 678–679); urinogenitalia, Fig. 469 (p. 684); internal genitalia of female, Fig. 470 (p. 685); placentation, Fig. 471 (p. 686).

SUB-CLASS PROTOTHERIA

Priacodon, skull, Fig. 472 (p. 687); **Ornithorhynchus**, whole animal in habitat, Fig. 473 (p. 689); **Tachyglossus**, whole animal in habitat, Fig. 474 (p. 690); **Ornithorhynchus**, cephalic anatomy, Fig. 475 (p. 691); buccal apparatus, Fig. 476 (p. 691); manus and pes, Fig. 477 (p. 692); poison apparatus, Fig. 478 (p. 693); eggs, Fig. 479 (p. 693); mammary glands, cloaca and tail, Fig. 480 (p. 694); dentition, Fig. 481 (p. 695); **Tachyglossus**, cephalic anatomy, Fig. 482 (p. 695); manus and pes, Fig. 483 (p. 696); **Ornithorhynchus**, endoskeleton, Fig. 484 (p. 697); **Tachyglossus**, skull, Fig. 485 (p. 698); **Ornithorhynchus**, scapula, Fig. 486 (p. 699); **Tachyglossus**, female urinogenitalia, Fig. 487 (p. 700); reptilian and monotreme affinities shown in male urinogenitalia, Fig. 488 (p. 701).

SUB-CLASS ALLOTHERIA

Order Multituberculata: Tæniolabidiæ, skull, Fig. 489 (p. 702).

SUB-CLASS THERIA

Order Dryolestoidæ: Amphitherium, lower jaw, Fig. 490 (p. 703).

Map: animal distribution, continental connections, Fig. 491 (p. 705); **Map**: faunal regions, Fig. 492 (p. 706).

Order Marsupialia: Tree-kangaroo, **Dendrolagus**, Fig. 493 (p. 707); Water Opossum, **Chironectes**, Fig. 494 (p. 711); Tiger 'Cat', **Dasyurops**, Fig. 495 (p. 713); Tasmanian 'Wolf', **Thylacinus**, Fig. 496 (p. 714); Marsupial 'Mole', **Notoryctes**, Fig. 497 (p. 715); Pygmy Glider, **Acrobates**, Fig. 498 (p. 717); **Diprotodon**, skeleton, Fig. 499 (p. 718); **Nototherium**, skull and dentition, Fig. 500 (p. 718); **Thylacoleo**, skull, Fig. 501 (p. 718); Kangaroo, **Macropus**, atlas, Fig. 502 (p. 719); skeleton, Fig. 503 (p. 720); Wombat, **Phascodomys**, skull, Fig. 504 (p. 720); Marsupial 'Cat', **Dasyurus**, skull, Fig. 505 (p. 721); Rock Wallaby, **Petrogale**, skull, Fig. 506 (p. 721); Possum, **Trichosurus**, hind-leg and pes, Fig. 507 (p. 722); **Macropus**, pes, Fig. 508 (p. 723).

Order Lipotyphla: **Solenodon**, Fig. 509 (p. 728); Golden Mole, **Chrysochloris**, Fig. 510 (p. 729); Tenrec, **Centetes**, skull, Fig. 511 (p. 730); Mole, **Talpa**, sternum and girdle, Fig. 512 (p. 730); fore-arm and manus, Fig. 513 (p. 731).

Order Menotyphla: Tree-shrews, **Tupaia**, Fig. 514 (p. 732); **Ptilocercus**, skull, Fig. 515 (p. 733).

Order Primates: *Lemur*, Fig. 516 (p. 737); skull and dentition, Fig. 517 (p. 738); *Pero-dicticus*, Fig. 518 (p. 739); *Tarsius*, Fig. 519 (p. 740); skull and dentition, Fig. 520 (p. 741); faces of Platyrrhine (*Ateles*) and Catarrhine (*Macaca*) compared, Fig. 521 (p. 742); *Gorilla*, Fig. 522 (p. 745); jaw structures of *Gorilla* and *Homo* compared, Fig. 523 (p. 746); *Homo*, skull, Fig. 524 (p. 749); skeletons of *Homo* and *Hylobates* compared, Fig. 525 (p. 751); *Papio*, carpus, Fig. 526 (p. 752); Feet of *Homo*, *Gorilla* and *Pongo*, Fig. 527 (p. 753).

Order Dermoptera: *Colugo*, *Cynocephalus*, Fig. 528 (p. 754).

Order Chiroptera. Fruit-bats, *Pteropus* Fig. 529 (p. 755); feeding adaptations of *Microchiroptera*, Fig. 530 (p. 757); *Pteropus*, skeleton, Fig. 531 (p. 759).

Order Edentata: Fairy Armadillo, *Chlamyphorus*, Fig. 532 (p. 762); Three-banded armadillo *Tolypeutes*, Fig. 533 (p. 763); defensive mechanism, Fig. 534 (p. 764); *Glyptodon*, skeleton, Fig. 535 (p. 764); *Mylodon*, skeleton, Fig. 536 (p. 765); Anteaters, *Myrmecophaga*, Fig. 537 (p. 766) and *Tamandua*, Fig. 538 (p. 767); Three-toed sloth, *Bradypus*, Fig. 539 (p. 768); whole skeleton, shoulder girdle, manus, pes, skull, Figs. 540-544 (pp. 769-770); *Dasypus*, skull, Fig. 545 (p. 770); *Myrmecophaga*, skull, Figs. 546-547 (pp. 770-771); *Dasypus*, shoulder-girdle, Fig. 548 (p. 771); pelvis and sacrum, Fig. 549 (p. 772).

Order Pholidota. Pangolin, *Manis*, Figs. 550-551 (p. 773).

Order Rodentia. *Rattus*, jaw musculature, Fig. 552 (p. 776); Molar dentition in rodents, Fig. 553 (p. 777).

Order Cetacea: Whale-bone (*Balenoptera*) and toothed (*Orcinus*) whales, Fig. 554 (p. 780); skull of *Protocetus*, Fig. 555 (p. 782); *Basilosaurus*, skeleton; Fig. 556 (p. 782); evolution of blow-hole, Fig. 557 (p. 782); *Neobalæna*, baleen and tongue sieve mechanism, Figs. 558-559 (pp. 785-786); *Balæna*, vertebrae, Fig. 560 (p. 786); porpoise, *Phocæna*, skeleton, Fig. 561 (p. 787); *Balenoptera*, sternum, Fig. 562 (p. 788); dolphin, *Globicephala*, skull, Fig. 563 (p. 788); hyperphalangy, Fig. 564 (p. 789).

Order Carnivora: Creodont and fissipede dentition compared, Fig. 565 (p. 793); Dentition in modern carnivores, Fig. 566 (p. 794); Cheetah, *Acinonyx*, Fig. 567 (p. 795); dentition of Sabre-tooth (*Smilodon*) and true Cat (*Metailurus*) compared, Fig. 568 (p. 796); Seal, *Phoca*, skeleton, Fig. 569 (p. 798); *Canis*, dog, skull, Fig. 570 (p. 799); *Felis*, tiger, auditory bulla, Fig. 571 (p. 799); *Ursus*, bear, auditory bulla, Fig. 572 (p. 800); carpus, Fig. 573 (p. 800).

Order Condylarthra: *Hyopsodus*, skull, Fig. 574 (p. 802); *Phenacodus*, Fig. 575 (p. 803).

Order Tubulidentata: Aardvark, *Orycteropus*, Fig. 576 (p. 804).

Order Litopterna: *Diadiaphorus*, Fig. 577 (p. 805).

Order Notoungulata: *Toxodon*, jaw, Fig. 578 (p. 806); *Homalodotherium*, skeleton, Fig. 579 (p. 806).

Order Astrapotheria: *Astrapotherium*, skeleton, Fig. 580 (p. 807).

Order Hyracoidea: Rock Hyrax, *Procavia*, Fig. 581 (p. 809); skull and dentition, Fig. 582 (p. 810).

Order Embrithopoda: *Arsinoitherium*, Fig. 583 (p. 811).

Order Proboscidea: Hairy mammoth, *Elephas*, Fig. 584 (p. 812); *Loxodonta*, African elephant, skull pneumatization, Fig. 585 (p. 813); *Moeritherium*, skull and dentition, Fig. 586 (p. 814); *Palaemastodon*, skull and dentition, Fig. 587 (p. 815); *Deinotherium*, Fig. 588 (p. 816); skull and dentition, Fig. 589 (p. 817); Dentition of African (*Loxodonta*) and Indian (*Elephas*) elephants compared, Fig. 590 (p. 817).

Order Pantodonta. *Coryphodon*, Fig. 591 (p. 818).

Order Dinocerata. *Uintatherium*, Fig. 592 (p. 819); dentition of *Uintatheriida*, Fig. 593 (p. 820).

Order Sirenia: Manatee, *Trichechus*, Fig. 594 (p. 821); *Dugong*, skeleton, Fig. 595 (p. 822); *Trichechus*, skull and dentition, Fig. 596 (p. 823).

Order Perissodactyla: Phylogeny and distribution of horses (Equidae), Fig. 597 (p. 827); limbs and teeth, Fig. 598 (p. 828); *Brontops*, Fig. 599 (p. 832); *Tapirus*, and young, Fig. 600 (p. 834); *Baluchitherium*, Fig. 601 (p. 835).

Order Artiodactyla: *Hippopotamus*, Fig. 602 (p. 838).

General Organisation of Mammalia: Hair, Fig. 603 (p. 841); tactile vibrissæ, Fig. 604 (p. 842); hair-tracts, Fig. 605 (p. 843); claw structure, Fig. 606 (p. 845); pes of *Tarsius*, Fig. 607 (p. 846); cutaneous glands and olfactory signals, Fig. 608 (p. 847); territory establishment in Ungulata, Fig. 609 (p. 848); mammalian skull (relationships of principal bones), Fig. 610 (p. 852); tympanic structure, Fig. 611 (p. 853); hyoid apparatus, Fig. 612 (p. 854); cranial modifications, Fig. 613 (p. 855); primitive plantigrade limbs of Carnivora, Fig. 614 (p. 857); convergence in cursorial adaptations, Fig. 615 (p. 858); paraxonic carpus and tarsus, Fig. 616

(p. 859); mammalian limb structures, Figs. 617–624 (pp. 860–861); tooth structure, Fig. 625 (p. 863); deciduous and permanent dentition of *Canis*, Fig. 626 (p. 864); Homodont dentition of dolphins (in *Lagenorhynchus*), Fig. 627 (p. 865); marsupial dentition (in *Perameles*), Fig. 628 (p. 865); Cope–Osborn theory of tritubercular origin of molars, Fig. 629 (p. 866); skull and dentition of *Phascolarctos*, Fig. 630 (p. 869); dentition of *Macropus*, Fig. 631 (p. 870); skull and dentition in *Sarcophilus*, Fig. 632 (p. 870); dentition of *Didelphys*, Fig. 633 (p. 871); dentition of *Orycteropus*, Aardvark, Fig. 634 (p. 871); dentition of *Sus*, pig, Fig. 635 (p. 872); skull and dentition of *Equus*, horse, Fig. 636 (p. 873); foetal jaw and teeth of Right Whale, *Balenoptera*, Fig. 637 (p. 874); pharyngeal region of Man, Fig. 638 (p. 877); rumination and digestion in Eutherians, Fig. 639 (p. 878); marsupial digestion, Fig. 640 (p. 879); brain of *Tachyglossus* (Spiny Ant-eater), *Petrogale* (Rock Wallaby), *Ornithorhynchus*, (Platypus) *Macropus* (Kangaroo), *Kogia* (Whale), Figs. 641–646 (pp. 885–886); female external genitalia of *Lemur*, Fig. 647 (p. 890); female urinogenitalia in Metatheria, Fig. 648 (p. 891); in Eutheria, Fig. 649 (p. 892); external genitalia in *Crocota crocuta*, Spotted Hyæna, Fig. 650; (p. 893) internal genitalia in *C. crocuta*, Fig. 651 (p. 894); development and placentation in Eutheria, Figs. 652–656 (pp. 900–904), Metatheria, Figs. 657–658 (p. 906); relative size of blastula in eutherian and monotreme, Fig. 659 (p. 907); mammary foetus of *Macropus*, Fig. 660 (p. 907).