

**Editors-in-Chief:**

J.-M. Morel, Cachan

B. Teissier, Paris

**Advisory Board:**

Camillo De Lellis, Zurich

Mario di Bernardo, Bristol

Alessio Figalli, Austin

Davar Khoshnevisan, Salt Lake City

Ioannis Kontoyiannis, Athens

Gabor Lugosi, Barcelona

Mark Podolskij, Aarhus

Sylvia Serfaty, Paris and New York

Catharina Stroppel, Bonn

Anna Wienhard, Heidelberg

More information about this series at <http://www.springer.com/series/304>

Fritz Gesztesy • Marcus Waurick

# The Callias Index Formula Revisited

 Springer

Fritz Gesztesy  
Dept of Mathematics  
University of Missouri  
Missouri  
Columbia, USA

Marcus Waurick  
Institut für Analysis  
TU Dresden  
Sachsen  
Dresden, Germany

ISSN 0075-8434

ISSN 1617-9692 (electronic)

Lecture Notes in Mathematics

ISBN 978-3-319-29976-1

ISBN 978-3-319-29977-8 (eBook)

DOI 10.1007/978-3-319-29977-8

Library of Congress Control Number: 2016942549

Mathematics Subject Classification (2010): 47A53, 47F05, 47B25

© Springer International Publishing Switzerland 2016

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

This Springer imprint is published by Springer Nature  
The registered company is Springer International Publishing AG Switzerland

# Preface

We revisit the Callias index formula in connection with supersymmetric Dirac-type operators  $H$  of the form

$$H = \begin{pmatrix} 0 & L^* \\ L & 0 \end{pmatrix}$$

in odd space dimensions  $n$ , originally derived in 1978, and prove that

$$\begin{aligned} \text{ind}(L) &= \left(\frac{i}{8\pi}\right)^{(n-1)/2} \frac{1}{[(n-1)/2]!} \lim_{\Lambda \rightarrow \infty} \frac{1}{2\Lambda} \sum_{i_1, \dots, i_n=1}^n \varepsilon_{i_1 \dots i_n} \\ &\quad \times \int_{\Lambda S^{n-1}} \text{tr}_{\mathbb{C}^d} (U(x) (\partial_{i_1} U)(x) \dots (\partial_{i_{n-1}} U)(x) x_{i_n}) d^{n-1} \sigma(x), \end{aligned} \quad (1)$$

where

$$U(x) := |\Phi(x)|^{-1} \Phi(x) = \text{sgn}(\Phi(x)), \quad x \in \mathbb{R}^n.$$

Here the closed operator  $L$  in  $L^2(\mathbb{R}^n)^{2\hat{d}}$  is of the form

$$L = \mathcal{Q} + \Phi,$$

where

$$\mathcal{Q} := Q \otimes I_d = \left( \sum_{j=1}^n \gamma_{j,n} \partial_j \right) I_d,$$

with  $\gamma_{j,n}, j \in \{1, \dots, n\}$ , elements of the Euclidean Dirac algebra, such that  $n = 2\hat{n}$  or  $n = 2\hat{n} + 1$ . Here  $\Phi$  is identified with  $I \otimes \Phi$ , satisfying

$$\begin{aligned}\Phi &\in C_b^2(\mathbb{R}^n; \mathbb{C}^{d \times d}), \quad d \in \mathbb{N}, \\ \Phi(x) &= \Phi(x)^*, \quad x \in \mathbb{R}^n,\end{aligned}$$

there exists  $c > 0, R \geq 0$  such that

$$|\Phi(x)| \geq cI_d, \quad x \in \mathbb{R}^n \setminus B(0, R),$$

and there exists  $\varepsilon > 1/2$  such that, for all  $\alpha \in \mathbb{N}_0^n, |\alpha| < 3$ , there is  $\kappa > 0$  such that

$$\|(\partial^\alpha \Phi)(x)\| \leq \begin{cases} \kappa(1 + |x|)^{-1}, & |\alpha| = 1, \\ \kappa(1 + |x|)^{-1-\varepsilon}, & |\alpha| = 2, \end{cases} \quad x \in \mathbb{R}^n.$$

These conditions on  $\Phi$  render  $L$  a Fredholm operator, and to the best of our knowledge, they represent the most general conditions known to date for which the Callias index formula (1) has been derived.

We also consider a generalization of the index formula (1) to certain classes of non-Fredholm operators  $L$  for which (1) represents its (generalized) Witten index (based on a resolvent regularization scheme).

Columbia, MO, USA  
Dresden, Saxony, Germany  
October 2015

Fritz Gesztesy  
Marcus Waurick

# Acknowledgments

We thank Bernhelm Booß-Bavnbek, Alan Carey, Stamatis Dostoglou, Yuri Latushkin, Marius Mitrea, and Andreas Wipf for valuable discussions and/or correspondence. We are especially grateful to Alan Carey for extensive correspondence and hints to the literature and to Stamatis Dostoglou for a critical reading of a substantial part of our manuscript and for many constructive comments that helped in improving our exposition. We also thank Ralph Chill, Martin Körber, Rainer Picard, and Sascha Trostorff for the useful discussions, particularly, concerning the counterexample discussed in Appendix B. In addition, we are indebted to Hendrik Vogt for kindly communicating the example in Remark 3.5.

M.W. gratefully acknowledges the hospitality of the Department of Mathematics at the University of Missouri, USA, extended to him during two one-month visits in the spring of 2014 and 2015. Moreover, he is particularly indebted to Yuri Tomilov in connection with the support received from the EU grant “AOS,” FP7-PEOPLE-2012-IRSES, No. 318910.

We gratefully acknowledge assistance from the staff of Springer-Verlag, especially from Ute McCrory.





# Contents

|           |   |     |
|-----------|---|-----|
| <b>1</b>  | <b>Introduction</b> .....   | 1   |
| <b>2</b>  | <b>Notational Conventions</b> .....   | 9   |
| <b>3</b>  | <b>Functional Analytic Preliminaries</b> .....  | 13  |
| <b>4</b>  | <b>On Schatten–von Neumann Classes and Trace Class Estimates</b> .....  | 23  |
| <b>5</b>  | <b>Pointwise Estimates for Integral Kernels</b> .....   | 35  |
| <b>6</b>  | <b>Dirac-Type Operators</b> .....   | 55  |
| <b>7</b>  | <b>Derivation of the Trace Formula: The Trace Class Result</b> .....  | 65  |
| <b>8</b>  | <b>Derivation of the Trace Formula: Diagonal Estimates</b> .....  | 77  |
| <b>9</b>  | <b>The Case <math>n = 3</math></b> .....  | 101 |
| <b>10</b> | <b>The Index Theorem and Some Consequences</b> .....  | 107 |
| <b>11</b> | <b>Perturbation Theory for the Helmholtz Equation</b> .....   | 119 |
| <b>12</b> | <b>The Proof of Theorem 10.2: The Smooth Case</b> .....   | 131 |
| <b>13</b> | <b>The Proof of Theorem 10.2: The General Case</b> .....  | 151 |
| <b>14</b> | <b>A Particular Class of Non-Fredholm Operators<br/>    <i>L</i> and Their Generalized Witten Index</b> ..... | 157 |
| <b>A</b>  | <b>Construction of the Euclidean Dirac Algebra</b> .....  | 167 |
| <b>B</b>  | <b>A Counterexample to [22, Lemma 5]</b> .....  | 175 |
|           | <b>References</b> .....   | 185 |
|           | <b>Index</b> .....  | 191 |