

# Lecture Notes in Mathematics

1972

**Editors:**

J.-M. Morel, Cachan

F. Takens, Groningen

B. Teissier, Paris

Takuro Mochizuki

# Donaldson Type Invariants for Algebraic Surfaces

Transition of Moduli Stacks

 Springer

Takuro Mochizuki  
Research Institute for Mathematical Sciences  
Kyoto University  
Kyoto 606-8502  
Japan  
takuro@kurims.kyoto-u.ac.jp

ISBN: 978-3-540-93912-2      e-ISBN: 978-3-540-93913-9  
DOI: 10.1007/978-3-540-93913-9

Lecture Notes in Mathematics ISSN print edition: 0075-8434  
ISSN electronic edition: 1617-9692

Library of Congress Control Number: 2008943979

Mathematics Subject Classification (2000): 14D20, 14J60, 14J80

© 2009 Springer-Verlag Berlin Heidelberg

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer. Violations are liable to prosecution under the German Copyright Law.

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

*Cover design:* SPi Publisher Services

Printed on acid-free paper

9 8 7 6 5 4 3 2 1

springer.com

*To my friends Yoshinobu Akahori  
and Shu Kawaguchi for memory  
of our informal student seminar*

# Preface

In this monograph, we define and investigate an algebro-geometric analogue of Donaldson invariants by using moduli spaces of semistable sheaves with arbitrary ranks on a polarized projective surface. We may expect the existence of interesting “universal relations among invariants”, which would be a natural generalization of the “wall-crossing formula” and the “Witten conjecture” for classical Donaldson invariants. Our goal is to obtain a weaker version of such relations, in other brief words, to describe a relation as the sum of integrals over the products of moduli spaces of objects with lower ranks. Fortunately, according to a recent excellent work of L. Göttsche, H. Nakajima and K. Yoshioka, [53], a wall-crossing formula for Donaldson invariants of projective surfaces can be deduced from such a weaker result in the rank two case.

We hope that our work in this monograph would, at least tentatively, provides a part of foundation for the further study on such universal relations.

In the rest of this preface, we would like to explain our motivation and some of important ingredients of this study. See *Introduction* for our actual problems and results.

## Donaldson Invariants

Let us briefly recall Donaldson invariants. We refer to [22] for more details and precise. We also refer to [37], [39], [51] and [53]. Let  $X$  be a compact simply connected oriented real 4-dimensional  $C^\infty$ -manifold with a Riemannian metric  $g$ . Let  $P$  be a principal  $SO(3)$ -bundle on  $X$ . A connection  $\nabla$  of  $P$  is called an anti-self-dual (ASD) connection if the curvature  $F(\nabla)$  satisfies  $*_g F(\nabla) = -F(\nabla)$ , where  $*_g$  denotes the Hodge star operator associated to  $g$ . For simplicity, let us restrict ourselves to the case that  $P$  comes from a principal  $SU(2)$ -bundle. Note that  $P$  is determined by its second Chern class  $c_2 := c_2(P)$ . Let  $b^+(X)$  be the number of positive eigenvalues of the intersection form on  $H^2(X)$ . Let  $M(c_2, g)$  denote the moduli space of ASD connections of  $P$ . If  $g$  is sufficiently general, it is known that  $M(c_2, g)$  is naturally a  $C^\infty$ -manifold with  $\dim_{\mathbb{R}} M(c_2, g) = 8c_2 - 3(1 + b^+(X))$ .

Let  $d := \dim_{\mathbb{R}} M(c_2, g)/2$  and write  $d = l + 2m$ . We consider integrals

$$\Phi_d^{X,g}(\alpha_1 \cdots \alpha_l \cdot p^m) := \int_{M(c_2,g)} \mu(\alpha_1) \cup \cdots \cup \mu(\alpha_l) \cdot \mu(P)^m \quad (0.1)$$

for  $\alpha_i \in H_2(X, \mathbb{Q})$  ( $i = 1, \dots, l$ ) and  $p \in H_0(X, \mathbb{Q})$ . The map  $\mu : H_*(X, \mathbb{Q}) \rightarrow H^*(M(c_2, g), \mathbb{Q})$  is formally given by  $\mu(\alpha) = c_2(\mathcal{P})/\alpha$ , where  $\mathcal{P}$  is the universal principal bundle on  $X \times M(c_2, g)$ . Although  $M(c_2, g)$  is not compact in general,  $\mu(\alpha_i)$  are naturally extended to the cohomology classes on the Uhlenbeck compactification  $\overline{M}(c_2, g)$ .

Moreover, let  $\mathfrak{A}_*(X)$  be the symmetric algebra generated by  $H_2(X) \oplus H_0(X)$ , which is graded by giving degree  $2 - i/2$  to elements of  $H_i(X)$ . Then,  $\Phi_d^{X,g}$  gives a linear map  $\mathfrak{A}_d(X) \rightarrow \mathbb{Q}$ . Thus, we obtain a map

$$\Phi^{X,g} := \sum_{d \geq 0} \Phi_d^{X,g} : \mathfrak{A}_*(X) \rightarrow \mathbb{Q}. \quad (0.2)$$

It is called a Donaldson invariant of  $X$ . If  $b^+(X) > 1$ , (0.1) are shown to be independent of the choice of general  $g$ .

They were successfully applied in the study of low-dimensional differential topology, until the appearance of Seiberg-Witten invariants which are defined more easily and believed to contain equivalent information in most cases. Nowadays, the attention to Donaldson invariants has been rather limited. So the author should explain why he would like to study a generalization of their algebro-geometric analogue in this comparatively long monograph. Although they might be less interesting in terms of topological application, there would exist attractive problems of “universal relations among invariants” which are natural generalizations of the “wall-crossing formula” (“Kotschick-Morgan conjecture”) and the “Witten conjecture”.

We remark that P. Kronheimer [76] studied such a generalization of Donaldson invariants for real 4-dimensional manifolds by using the moduli of higher rank objects from a viewpoint of differential geometry. It was also investigated in mathematical physics. (See [84], for example.)

### Kotschick-Morgan Conjecture and Witten Conjecture

Let us recall the conjectures for an explanation of the motivation of our study, although they are beyond the scope of this monograph. We should remark that they have been studied intensively from mathematical viewpoints, by V. Y. Pidstrigach–A. N. Tyurin, B. Chen, and in particular P. Feehan–T. Leness.

In the case  $b^+ = 1$ , the integrals (0.1) depend on the choice of the metric  $g$ . Let us recall the result of D. Kotschick–J. Morgan in [74], following [51]. Let  $H^2(X, \mathbb{R})^+$  denote the set of  $\alpha \in H^2(X, \mathbb{R})$  such that  $\alpha^2 > 0$ . Then,  $H^2(X, \mathbb{R})^+/\mathbb{R}^+$  has two connected components. Let us fix one component  $\Omega^+$ . The period point  $\omega(g)$  is the point in  $\Omega^+$  defined by the closed two form which is harmonic with respect to  $g$  and satisfies  $*_g \omega(g) = \omega(g)$ . An element  $\xi \in H^2(X, \mathbb{Z})$  is called of type  $d$ , if

$(d+3)/4 + \xi^2 \in \mathbb{Z}_{\geq 0}$ . For such  $\xi$ , let  $W^\xi := \{L \in \Omega^+ \mid L \cdot \xi = 0\}$  be called a wall of type  $d$ . The chambers of type  $d$  are the connected components in the complement of the walls of type  $d$  in  $\Omega^+$ . Kotschick and Morgan proved that  $\Phi_d^{X,g}$  depends only on the chamber in which  $\omega(g)$  is contained. It is very interesting to ask what happens when  $\omega(g)$  crosses a wall of type  $d$ , that is called a wall-crossing phenomenon. We obtain the linear map  $\delta_{\xi,d}^X : \mathfrak{A}_d(X) \rightarrow \mathbb{Q}$  given by  $\Phi_d^{X,g_1} - \Phi_d^{X,g_2} = \delta_{\xi,d}^X$ , where the chambers of  $\omega(g_1)$  and  $\omega(g_2)$  are divided with  $W^\xi$ . They conjectured that  $\delta_{\xi,d}^X(\alpha^d)$  ( $\alpha \in H_2(X)$ ) is the polynomial in the intersection numbers  $\alpha^2$  and  $\alpha \cdot \xi$  whose coefficients depend only on  $\xi^2$ ,  $d$  and the homotopy type of  $X$ . Very interestingly, assuming the conjecture, Göttsche showed that  $\delta_{\xi,d}^X$  can be expressed by using modular forms, which is called a *wall-crossing formula*. ([50]. See also [51].)

Since the appearance of Seiberg-Witten invariants, it has been expected that Seiberg-Witten invariants and Donaldson invariants give equivalent information in most cases. Let  $\chi$  and  $\tau$  denote the Euler number and the signature of  $X$ , respectively. Let  $m(X) = 2 + (7\chi + \tau)/4$ . Let  $\Lambda$  denote the set of isomorphism classes of line bundles  $L$  on  $X$  such that  $c_1(L)^2 = 2\chi + 3\tau$ . For  $L \in \Lambda$ , let  $n_L$  denote the Seiberg-Witten invariant, i.e., the number of the solutions of the Seiberg-Witten equation associated to a  $\text{Spin}^c$ -structure whose determinant bundle is  $L$ . Then, Donaldson invariants and Seiberg-Witten invariants are expected to be related by the following impressive formula:

$$\Phi^X(\alpha) = 2^{m(X)} e^{\alpha \cdot \alpha / 2} \sum_{L \in \Lambda} n_L \cdot e^{c_1(L) \cdot \alpha}$$

(See [130] and [21] for more precise and details.)

Both conjectures claim the existence of universal relations which are expressed in impressive ways. It might be desirable to understand them as a part of larger universal relations by extending the framework to the higher rank cases, and moreover we might dream to understand some geometry behind them.

### Donaldson Invariants in Algebraic Geometry

We recall how Donaldson invariants were discussed in algebraic geometry. Let  $X$  be a smooth projective surface with an ample line bundle  $\mathcal{O}_X(1)$ . Let  $\omega$  be a Kahler metric which represents the first Chern class of  $\mathcal{O}_X(1)$ .

Let  $E$  be a holomorphic vector bundle on  $X$  such that  $\det(E) \simeq \mathcal{O}_X$ . A hermitian metric  $h$  of  $E$  is called Hermitian-Einstein with respect to  $\omega$ , if  $\Lambda_\omega F(h) = 0$  is satisfied, where  $F(h)$  is the curvature of the canonical unitary connection  $\nabla_h$  of  $(E, h)$ , and  $\Lambda_\omega$  denotes the adjoint of the multiplication of  $\omega$ . Let  $P$  be the principal  $SU(2)$ -bundle associated to  $E$  with a hermitian metric  $h$ . Then, the induced connection  $\nabla_h$  on  $P$  is ASD, if and only if  $h$  is Hermitian-Einstein. Therefore, considering ASD-connections is equivalent to considering holomorphic vector bundles with Hermitian-Einstein metrics.

Recall a very deep correspondence between objects in differential geometry and algebraic geometry, so called the Kobayashi-Hitchin correspondence. (See [18], [19], [128].) Let  $E$  be a holomorphic vector bundle on  $X$  such that  $\det(E) \simeq \mathcal{O}_X$ . Then,  $E$  is called  $\mu$ -stable if and only if  $\deg_\omega(F) := \int_X c_1(F) \cdot \omega < 0$  for any coherent subsheaf  $F$  of  $E$ . According to the Kobayashi-Hitchin correspondence,  $E$  is  $\mu$ -stable if and only if  $E$  is indecomposable and has a Hermitian-Einstein metric. Moreover, such a metric is unique up to obvious ambiguity. Thus, we obtain the bijective correspondence between  $\mu$ -stable bundles and holomorphic indecomposable vector bundles with Hermitian-Einstein metrics, more strongly the homeomorphism of the moduli spaces.

Stable bundles and more general stable sheaves have been studied in algebraic geometry. A moduli space  $\mathcal{M}(c_2)$  of semistable sheaves was constructed as the projective variety by D. Gieseker and M. Maruyama in the framework of geometric invariant theory, which gives a compactification of moduli of stable vector bundles, called the Gieseker-Maruyama compactification. It was known that there exists the projective morphism of the Gieseker-Maruyama compactification to the Uhlenbeck compactification. It is natural to expect that the integral (0.1) can be defined as the evaluation over  $\mathcal{M}(c_2)$ . It doesn't work in general, since a Kahler metric is not necessarily generic and the moduli space  $\mathcal{M}(c_2)$  does not have the expected dimension. But, if  $c_2$  is sufficiently large, it is known that  $\mathcal{M}(c_2)$  has the expected dimension, and the evaluation over  $\mathcal{M}(c_2)$  gives the Donaldson invariants according to J. Li and Morgan. (See [81] and [94].) By using the blow up formulas due to R. Fintushel and R. J. Stern [34], which relate the Donaldson invariants of  $X$  and the blow up  $\hat{X}$  of  $X$  in a point, the full Donaldson invariants can be defined in a purely algebro-geometric way. (See [53] for more details.) Note that we will later give a different construction of invariants in an algebro-geometric way. (They are the same, if the moduli has the expected dimension.) Each method has its own advantage.

## Reduction to Sum of Integrals over the Products of Hilbert Schemes

G. Ellingsrud-L. Göttsche and R. Friedman-Z. Qin studied the wall crossing phenomena of Donaldson invariants in algebraic geometry. (See [26] and [36].) Although we omit the details here, their results briefly say that if  $X$  is a smooth projective surface,  $\delta_{\xi,d}^X$  can be described as the sum of integrals of naturally induced cohomology classes over the products of Hilbert schemes of points on  $X$ , under the assumption that *the wall  $W^\xi$  is good*. Such a formula is called a *weak wall-crossing formula* in this monograph, and actually one of our main goals is to show it without any assumption on the walls by using *intrinsic smoothness* of the moduli spaces.

## Hilbert Schemes of Points on a Surface

The author thinks that it is a proper goal to describe some relations among invariants as the sum of integrals over the products of Hilbert schemes, although not final.



For an explanation, we recall a brief history of the study of Hilbert schemes  $X^{[n]}$  of  $n$ -points on a connected projective surface  $X$ . We refer to [49] and [98] for more details. The first important result is the irreducibility and smoothness of  $X^{[n]}$ . Among several important and pioneering works, one of the most impressive and famous results is a formula of Göttsche. It is irresistible to reproduce it here. Let  $b_i(X)$  denote the  $i$ -th Betti numbers of  $X$ . Let  $P_{X^{[n]}}(z)$  denote the Poincaré polynomial of  $X^{[n]}$  in a variable  $z$ . Then, Göttsche's formula is

$$\sum_{n \geq 0} P_{X^{[n]}}(z)t^n = \prod_{m \geq 1} \frac{(1 + z^{2m-1}t^m)^{b_1(X)}(1 + z^{2m+1}t^m)^{b_3(X)}}{(1 - z^{2m-2}t^m)(1 - z^{2m}t^m)^{b_2(X)}(1 - z^{2m+2}t^m)}.$$

Many mathematicians were attracted by the formula for deeper understanding. For example, it has been shown to hold in the level of Grothendieck group of varieties ([11], [52], [16]). In the other direction, a deep observation was given by Nakajima and I. Grojnowski ([56] and [97]), who constructed geometrically a representation of Heisenberg algebra on  $\bigoplus_n H^*(X^{[n]})$ .

Since then, a considerable number of studies have been conducted on Hilbert schemes of points, and hence they have been much more familiar than moduli of sheaves with higher ranks. It could be one reason why we try to express some quantity as the sum of integrals over the products of Hilbert schemes. But moreover, in principle, we may expect that some information of Hilbert schemes  $X^{[n]}$  are comprehensible from that of the original surface  $X$  with universal and lucid procedures which are independent of  $X$ . The formula of Göttsche and the Nakajima-Grojnowski construction are typical. The principle was enforced by the results on cobordism classes, some integrals and others. (For example, see [27] and [80].) That is the main reason why the author believes it worthwhile to express something in terms of Hilbert schemes, like a weak wall-crossing formula in the rank two case.

Fortunately for the author, after the solution of Nekrasov's conjecture ([99], [100], [101], [102] [106]), Göttsche, Nakajima and Yoshioka successfully showed that the wall-crossing formula for Donaldson invariants of projective surface can be deduced from the weak wall-crossing formula! (See [53].) We refer to the lecture notes and the survey by Nakajima and Yoshioka for this story. ([99] and [102]).

### Virtual Fundamental Classes

We should emphasize that our moderate goal, such as a weak wall-crossing formula, is already related with important, interesting and rather recent development in mathematics, so called derived algebraic geometry. In their study of the wall-crossing phenomena, Ellingsrud-Göttsche and Friedman-Qin used a sequence of flips to relate the moduli spaces corresponding to two polarizations. Their argument can work if flips are neat, which is related with the assumption that the wall is good. But, singularity may appear in general.

Singularity of moduli spaces causes difficult but attractive problems in many aspects of mathematics. Since 1990's, several mathematicians, such as P. Deligne,

V. Drinfeld and M. Kontsevich, have proposed significance of *intrinsic smoothness of moduli spaces*. (See [71].) This revolutionary idea was employed most efficiently in the Gromov-Witten theory. In constructing Gromov-Witten invariants, we need to evaluate some cohomology classes over fundamental classes of moduli stacks of stable maps. However, the moduli stacks may not be smooth and of expected dimension, and hence they do not have fundamental classes in the naive sense. As a solution of this difficulty, L. Behrend-B. Fantechi, K. Fukaya-K. Ono and J. Li-G. Tian gave constructions of *virtual fundamental classes* by using intrinsic smoothness called perfect obstruction theory or Kuranishi structure ([6], [40] and [82]). Rather recently, virtual fundamental classes have also been used in the study of Donaldson-Thomas invariants [121] for three dimensional Calabi-Yau varieties.

As for our problems, we will define our algebro-geometric analogue of Donaldson invariants by using virtual fundamental classes, which allows us to utilize intrinsic smoothness effectively in the study of transitions of moduli stacks. More specifically, we can use a technique of localization with respect to torus actions to obtain weak wall crossing formulas and similar formulas, instead of flips.

## Master Space

Another important ingredient is the excellent and beautiful idea of *master space* due to M. Thaddeus [120]. Let  $G$  be a linear reductive group. Let  $U$  be a projective variety with a  $G$ -action. Let  $\mathcal{L}_i$  ( $i = 1, 2$ ) be  $G$ -polarizations of  $U$ . Then, we have the open subset  $U^{ss}(\mathcal{L}_i)$  of semistable points of  $U$  with respect to  $\mathcal{L}_i$ . It is interesting and significant to compare the quotient stacks  $\mathcal{M}_i := U^{ss}(\mathcal{L}_i)/G$  ( $i = 1, 2$ ). (Since the moduli stacks of semistable sheaves have such descriptions, it is clearly related with our problem.)

For that purpose, Thaddeus introduced the idea of master space. Let us consider the  $G$ -variety  $\text{TH} := \mathbb{P}(\mathcal{L}_1^{-1} \oplus \mathcal{L}_2^{-1})$  on  $U$ , the projectivization of the vector bundle  $\mathcal{L}_1 \oplus \mathcal{L}_2$ . We have the canonical polarization  $\mathcal{O}_{\mathbb{P}}(1)$  on  $\text{TH}$ . We have the canonically defined  $G$ -action on  $\text{TH}$ , and  $\mathcal{O}_{\mathbb{P}}(1)$  gives the  $G$ -polarization. The set of the semistable points is denoted by  $\text{TH}^{ss}$ . Then we obtain the quotient stack  $M := \text{TH}^{ss}/G$ . Let  $\pi : \text{TH}^{ss} \rightarrow M$  denote the projection.

Let  $G_m$  denote a one dimensional torus  $\text{Spec } \mathbb{C}[t, t^{-1}]$ . We have the  $G_m$ -action on  $\text{TH}$  given by  $\rho(t) \cdot [x : y] = [t \cdot x : y]$ , where  $[x : y]$  denotes the homogeneous coordinate of  $\text{TH}$  along the fiber direction. Since  $\rho$  commutes with the action of  $G$ , it induces a  $G_m$ -action  $\bar{\rho}$  on  $M$ .

We have the natural inclusions  $\text{TH}_i = \mathbb{P}(\mathcal{L}_i^{-1}) \rightarrow \text{TH}$ . Because  $\mathcal{O}_{\mathbb{P}}(1)|_{\text{TH}_i} = \mathcal{L}_i$ , we have  $\text{TH}_i^{ss} = U(\mathcal{L}_i)^{ss}$  and inclusions  $\mathcal{M}_i \rightarrow M$ . Since  $\text{TH}_i$  are components of the fixed point set of the action  $\rho$ , the stacks  $\mathcal{M}_i$  are the components of the fixed point set of the action  $\bar{\rho}$ .

The action  $\bar{\rho}$  may have fixed points not contained in  $\mathcal{M}_1 \cup \mathcal{M}_2$ . Let  $x \in \text{TH}^{ss}$ . In general,  $\pi(x)$  is fixed by the action  $\bar{\rho}$  if and only if  $G_m \cdot x \subset G \cdot x$ . So we may have the components of the fixed points  $\pi(x)$  such that  $x$  is not fixed by  $\rho$ . Such a component is called exceptional. In a sense, the information on the difference of

the quotient stacks  $\mathcal{M}_i$  ( $i = 1, 2$ ) is contained in the exceptional fixed point sets. As already mentioned, we will extract this information by using a  $G_m$ -localization technique in our situation.

We have one more technical remark. In the study of the transition of moduli stacks of objects with higher ranks, one of the main difficulties for the author is the appearance of objects which are decomposable into more than three components. In our argument, it causes that the master space is not Deligne-Mumford. To avoid such difficulty, we introduce enhanced master spaces, which requires a detailed and interesting investigation in controlling fixed point sets.

## Further Study

This monograph is tentative and experimental, partially because of rapid and intensive development of the theory of stacks. For example, J. Lurie, B. Toën and their collaborators are rewriting the foundation of algebraic geometry. (See [124] for overview.) Their theory, so called *derived algebraic geometry* or *homotopical algebraic geometry*, seems to provide us with a powerful tool in constructing perfect obstruction theories, and hence virtual fundamental classes. But, unfortunately, it is beyond the author's ability to use their theory at this moment. He believes that it should be one of the themes in the study of stacks to make it easy to deal with the objects and the formalisms in our work. He hopes to study these problems from the viewpoint of derived algebraic geometry once more in future. (However, he also believes that we will obtain the same invariants even if we adopt another method in constructing perfect obstruction theory.)

One of the important results missing in our theory is the blow-up formula, i.e., a comparison of invariants for  $X$  and a blow up of  $X$ . The author originally intended to develop a theory "without blow up". But it seems to contain interesting problems for such a comparison. For example, see the recent attractive work of Nakajima and Yoshioka, [103] and [104].

From a perfect obstruction theories, we obtain not only virtual fundamental classes but also *virtual structure sheaves*, which seem very attractive subjects. For example, it would be very interesting to find a formula to express the Euler numbers of the tensor products of line bundles and the virtual structure sheaves. (See [28], [54] and [101] for related topics.)

As mentioned in the beginning, we hope that our weak relations would lead to more impressive formulas. Moreover, it would be desirable to understand such formulas in a geometric way, as in the case of Göttsche's formula and the Nakajima-Grojnowski construction.

The study of wall-crossing phenomena for Donaldson-Thomas invariants has been developing intensively. (See [4], [73], [114], [123], for example.) The author expects that we will have wall crossing phenomena in many other situations. It would be very interesting if some of the techniques used in this monograph could be applied.

## Acknowledgements

I started this project in 1998, my last year at the graduate school in Kyoto University. I continued the study in Osaka City University and the Institute for Advanced Study. I resumed it in 2006 at the Max-Planck Institute for Mathematics, and continued the study at the Institut des Hautes Études Scientifiques. Then, the final version was written at Kyoto University, Department of Mathematics, and Research Institute for Mathematical Sciences. During the course of the study, I visited the Tata Institute of Fundamental Research and the International Centre for Theoretical Physics. I am heartily grateful to the institutions and the colleagues for the excellent mathematical environment, and I would like to express my deep gratitude to those who kindly supported me. The final manuscript is completely different from the first version after several major revisions, and it took very long time for publication partially because of my poor ability in writing, which I apologize to those who kindly showed their interest in this study. Anyway, I feel relief that the manuscript is published at last.

My mathematical interest has been clearly and deeply influenced by Kenji Fukaya. I learned many things through his talks and lectures. I was also very encouraged by his interest in my study. It is my great pleasure to express my gratitude to him.

I thank Hiraku Nakajima for his stimulating questions, which made me resume this study. And, I am grateful to Lothar Göttsche, H. Nakajima and Kōta Yoshioka for their amazing work and their interest in this study. I also thank Barbara Fantechi for her valuable comments and suggestions.

I express my gratitude to K. Fukaya, Yoshihiro Fukumoto, Mikio Furuta, Ryushi Goto, Yoshitake Hashimoto, Yukio Kametani, Hiroshi Konno, Norihiko Minami, Hiroshi Ohta, Kaoru Ono, Tatsuru Takakura, who kindly listened to a series of my bad talks on this study in Gero. It is impossible to express how I was glad and encouraged and how I appreciate them.

I thank Yoshifumi Tsuchimoto and Akira Ishii for their constant encouragement and advice since my undergraduate student days. All what I do in mathematics is based on what I learned from them.

Special thanks go to Mark de Cataldo, Pierre Deligne, William Fulton, David Gieseker, Masaki Kashiwara, Akira Kono, Mikiya Masuda, Claude Sabbah, Carlos Simpson, Tomohide Terasoma, Michael Thaddeus and Kari Vilonen for their help. I am willing to express my gratitude to Takeshi Abe, Kürsat Aker, Andreas Bender, Ryan Budney, Ron Donagi, Simon Donaldson, Michihiko Fujii, Koji Fujiwara, Rajendra Vasant Gurjar, Claus Hertling, Michiaki Inaba, Masaharu Kaneda, Atushi Kasue, Tsuyoshi Kato, Hiroshige Kajiura, Nariya Kawazumi, Yoshikata Kida, Maxim Kontsevich, Kong Liang, Kefeng Liu, Pierre Matsumi, Atsushi Matsuo, Vikram B. Mehta, Luca Migliorini, Hiroaki Narita, Shin-ichi Ohta, So Okada, Tony Pantev, Ramadas T. Ramakrishnan, Utkir Rozikov, Masa-Hiko Saito, Christian Sevenheck, Szilard Szabo, Alexander Usnich, Kimiko Yamada, Satoshi Yamaguchi for their kindness and some discussion. I still remember vigorous and powerful

lectures by Masaki Maruyama on semistable sheaves. I thank Fumiharu Kato for some valuable information on stacks and his kindness.

I would like to express my deep thanks to the anonymous referee for his careful and patient reading. I also appreciate his useful and practical advice in writing.

I thank the partial supports by the Ministry of Education, Culture, Sports, Science and Technology, the Japan Society for the Promotion of Science, the Sumitomo Foundation, and the Sasagawa Foundation. I wish to acknowledge the National Scientific Foundation for a grant DMS 9729992, although any opinions, findings and conclusions or recommendations expressed in this material do not necessarily reflect the views of the National Scientific Foundation.

Kyoto, June 2008

*Takuro Mochizuki*

# Contents

<b>1</b>	<b>Introduction</b> .....	1
1.1	Problems .....	1
1.2	Construction of Invariants .....	3
1.2.1	Orientation and Reduced $L$ -Bradlow Pairs .....	3
1.2.2	Virtual Fundamental Classes .....	5
1.3	Transition Formulas in a Simple Case .....	6
1.4	Rank Two Case .....	9
1.4.1	Dependence on Polarizations .....	9
1.4.2	Reduction to Integrals over Hilbert Schemes .....	11
1.5	Higher Rank Case .....	12
1.5.1	$p_g > 0$ .....	12
1.5.2	Transition Formula in the Case $p_g = 0$ .....	13
1.5.3	Weak Intersection Rounding Formula .....	15
1.6	Master Space .....	16
1.6.1	Master Space in the Case that a 2-Stability Condition is Satisfied .....	17
1.6.2	$G_m$ -Localization Method (I) .....	19
1.6.3	Enhanced Master Space .....	20
1.6.4	$G_m$ -Localization Method (II) .....	22
<b>2</b>	<b>Preliminaries</b> .....	25
2.1	Some Convention .....	25
2.1.1	Product and Projection .....	25
2.1.2	Vector Bundles .....	26
2.1.3	Coherent Sheaves on a Product .....	26
2.1.4	Quotient Stacks .....	26
2.1.5	Signature in Complexes .....	26
2.1.6	Filtrations and Complexes on a Curve .....	27
2.1.7	Virtual Vector Bundle .....	28
2.1.8	Compatible Diagrams .....	29

- 2.2 Geometric Invariant Theory . . . . . 29
  - 2.2.1 GIT Quotient and Algebraic Stacks . . . . . 29
  - 2.2.2 Mumford-Hilbert Criterion and Some Elementary Examples 30
- 2.3 Cotangent Complex . . . . . 32
  - 2.3.1 Basic Facts . . . . . 32
  - 2.3.2 Quotient Stacks . . . . . 34
  - 2.3.3 Some More Examples . . . . . 39
- 2.4 Obstruction Theory . . . . . 44
  - 2.4.1 Definition and Fundamental Theorems . . . . . 44
  - 2.4.2 Easy Example . . . . . 46
  - 2.4.3 Locally Free Subsheaves . . . . . 48
  - 2.4.4 Filtrations of a Vector Bundle on a Curve . . . . . 51
- 2.5 Equivariant Complexes on Deligne-Mumford Stacks  
with GIT Construction . . . . . 55
  - 2.5.1 Locally Free Resolution . . . . . 55
  - 2.5.2 Equivariant Representative . . . . . 56
- 2.6 Elementary Remarks on Some Extremal Sets . . . . . 57
  - 2.6.1 Preparation for a Proof of Proposition 4.3.3 . . . . . 57
  - 2.6.2 Preparation for a Proof of Proposition 4.4.4 . . . . . 59
- 2.7 Twist of Line Bundles . . . . . 61
  - 2.7.1 Construction . . . . . 61
  - 2.7.2 Weight of the Induced Action . . . . . 61
- 3 Parabolic  $L$ -Bradlow Pairs . . . . . 63**
  - 3.1 Sheaves with Some Structure and their Moduli Stacks . . . . . 64
    - 3.1.1 Orientation . . . . . 64
    - 3.1.2 Parabolic Structure . . . . . 65
    - 3.1.3  $L$ -Bradlow Pairs and Reduced  $L$ -Bradlow Pairs . . . . . 66
    - 3.1.4 Type and Moduli Stacks . . . . . 68
    - 3.1.5 Tautological Line Bundle and Relations  
Among Some Moduli Stacks . . . . . 69
  - 3.2 Hilbert Polynomials . . . . . 71
    - 3.2.1 Hilbert Polynomials of Coherent Sheaves . . . . . 71
    - 3.2.2 Hilbert Polynomials of Parabolic Sheaves . . . . . 71
    - 3.2.3 Hilbert Polynomial for Parabolic  $L$ -Bradlow Pairs . . . . . 72
    - 3.2.4 Hilbert Polynomials Associated to a Type . . . . . 72
  - 3.3 Semistability . . . . . 73
    - 3.3.1 Semistability Conditions and the Associated Moduli Stacks . 73
    - 3.3.2 Harder-Narasimhan Filtration and Partial Jordan-Hölder  
Filtration . . . . . 76
    - 3.3.3  $(\delta, \ell)$ -Semistability . . . . . 77
  - 3.4 Some Boundedness . . . . . 78
    - 3.4.1 Fundamental Theorems . . . . . 78
    - 3.4.2 Boundedness of Semistable  $L$ -Bradlow Pairs . . . . . 79
    - 3.4.3 Boundedness of Yokogawa Family . . . . . 80

- 3.5 1-Stability Condition and 2-Stability Condition ..... 84
  - 3.5.1 Parabolic Sheaf ..... 84
  - 3.5.2 Parabolic  $L$ -Bradlow Pair ..... 85
  - 3.5.3 Parabolic  $L$ -Bradlow Pair ..... 88
- 3.6 Quot Schemes ..... 89
  - 3.6.1 Preliminary ..... 89
  - 3.6.2 Quotient Sheaves ..... 89
  - 3.6.3 Quotient Quasi-Parabolic Sheaves and Maruyama-Yokogawa Construction ..... 90
  - 3.6.4 Quotient  $L$ -Bradlow Pair ..... 91
  - 3.6.5 Quotient Reduced  $L$ -Bradlow Pair ..... 93
  - 3.6.6 Quotient Reduced  $L$ -Bradlow Pair ..... 94
  - 3.6.7 Oriented Objects ..... 94
  - 3.6.8 Quotient Stacks ..... 95
- 4 Geometric Invariant Theory and Enhanced Master Space ..... 97**
  - 4.1 Semistability Condition and Mumford-Hilbert Criterion ..... 98
    - 4.1.1 Statement ..... 98
    - 4.1.2 Mumford-Hilbert Criterion ..... 101
    - 4.1.3 A Lemma ..... 102
    - 4.1.4 Proof of the Claim 1 in Proposition 4.1.2 ..... 103
    - 4.1.5 Proof of the Claim 2 of Proposition 4.1.2 ..... 104
    - 4.1.6 Complement ..... 105
  - 4.2 Perturbation of Semistability Condition ..... 106
    - 4.2.1 Preliminary ..... 106
    - 4.2.2  $\delta_+$ -Semistability and  $\delta_-$ -Semistability ..... 107
    - 4.2.3  $(\delta, \ell)$ -Semistability ..... 109
  - 4.3 Enhanced Master Space ..... 113
    - 4.3.1 Construction ..... 113
    - 4.3.2 Proof of Lemma 4.3.4 ..... 114
    - 4.3.3 Proof of Lemma 4.3.5 ..... 116
    - 4.3.4 Proof of Lemma 4.3.6, Step 1 ..... 117
    - 4.3.5 Proof of Lemma 4.3.6, Step 2 ..... 118
    - 4.3.6 Proof of Lemma 4.3.6, Step 3 ..... 119
  - 4.4 Fixed Point Set of Torus Action on Enhanced Master Space ..... 120
    - 4.4.1 Preliminary ..... 120
    - 4.4.2 Statements ..... 122
    - 4.4.3 Step 1 ..... 122
    - 4.4.4 Step 2 ..... 124
    - 4.4.5 Step 3 ..... 125
    - 4.4.6 Step 4 ..... 126
    - 4.4.7 End of the Proof of Proposition 4.4.4 ..... 127
  - 4.5 Enhanced Master Space in Oriented Case ..... 128
    - 4.5.1 Construction ..... 128
    - 4.5.2 Obvious Fixed Point Sets ..... 129



4.5.3	Fixed Point Sets Associated to Decomposition Types	130
4.5.4	Statement	130
4.5.5	Ambient Stack	131
4.5.6	Fixed Point Set of the Ambient Space	131
4.5.7	Proof of Proposition 4.5.3	132
4.6	Decomposition of Exceptional Fixed Point Sets	134
4.6.1	Statement	134
4.6.2	Preliminary	136
4.6.3	Construction of the Stack $\mathcal{S}$	136
4.6.4	Universal Sheaf	138
4.7	Simpler Cases	138
4.7.1	Case in Which a 2-Stability Condition is Satisfied	138
4.7.2	Oriented Reduced $L$ -Bradlow Pairs	140
<b>5</b>	<b>Obstruction Theories of Moduli Stacks and Master Spaces</b>	<b>145</b>
5.1	Deformation of Torsion-Free Sheaves	146
5.1.1	Construction of a Basic Complex	146
5.1.2	The Trace-Free Part and the Diagonal Part	148
5.1.3	Preparation for Master space	150
5.1.4	Basic Complex on the Moduli Stack $\mathcal{M}(m, y)$	151
5.1.5	Line Bundles	153
5.2	Relative Obstruction Theory for Orientations	154
5.2.1	Construction of a Complex	154
5.2.2	Relative Obstruction Property	155
5.3	Relative Obstruction Theory for $L$ -Sections	156
5.3.1	Construction of a Complex	156
5.3.2	Relative Obstruction Property	158
5.3.3	Preparation for Obstruction Theory of Master Space	160
5.3.4	Preparation for Proposition 6.2.1	161
5.3.5	Another Construction	162
5.4	Relative Obstruction Theory for Reduced $L$ -Sections	164
5.4.1	Construction of a Complex	164
5.4.2	Relative Obstruction Property	166
5.4.3	Preparation for Obstruction Theory of Master Space	168
5.4.4	Preparation for Proposition 6.2.1	170
5.5	Relative Obstruction Theory for Parabolic Structures	170
5.5.1	Construction of a Complex	170
5.5.2	Relative Obstruction Property	173
5.5.3	Decomposition into the Trace-Free Part and the Diagonal Part	173
5.6	Obstruction Theory for Moduli Stacks of Stable Objects	174
5.6.1	Relative Complexes	174
5.6.2	Construction of Complexes and Morphisms	175
5.6.3	Obstruction Theories of Quot Schemes and Moduli Stacks	176
5.6.4	Obstruction Theories of Moduli Stacks of Stable Objects	177

- 5.7 Obstruction Theory for Enhanced Master Spaces and Related Stacks ..... 178
  - 5.7.1 Enhanced Master Space ..... 178
  - 5.7.2 Substack  $\widehat{M}^*$  ..... 179
  - 5.7.3 Moduli Stack  $\widetilde{\mathcal{M}}(m, \widehat{y}, [L])$  ..... 181
  - 5.7.4 Moduli Stack  $\widetilde{\mathcal{M}}(m, y, L)$  ..... 184
  - 5.7.5 Moduli Stack  $\widetilde{\mathcal{M}}^s(\widehat{y}, \alpha_*, +)$  ..... 184
  - 5.7.6 Case in Which a 2-Stability Condition is Satisfied ..... 185
  - 5.7.7 Oriented Reduced  $L$ -Bradlow Pairs ..... 186
- 5.8 Moduli Theoretic Obstruction Theory of Fixed Point Set ..... 187
  - 5.8.1 Statement ..... 187
  - 5.8.2 Moduli Stack of Split Objects with Orientations ..... 188
  - 5.8.3 Embedding into Moduli Stack of Non-Split Objects ..... 190
  - 5.8.4 Some Compatibility ..... 194
  - 5.8.5 Deformation ..... 196
  - 5.8.6 Proof of Proposition 5.8.1 ..... 198
  - 5.8.7 Case in Which a 2-Stability Condition is Satisfied ..... 199
  - 5.8.8 Oriented Reduced  $L$ -Bradlow Pairs ..... 200
- 5.9 Equivariant Obstruction Theory of Master Space ..... 201
  - 5.9.1 Statements ..... 201
  - 5.9.2  $G_m$ -Equivariant Lift of  $\text{Ob}(\widehat{M})$  and  $\text{ob}(\widehat{M})$  ..... 204
  - 5.9.3 Comparison of  $G_m$ -Equivariant Structures of  $\tau_{\geq -1}L_{\widehat{M}}$  ..... 205
  - 5.9.4  $G_m$ -Equivariant structure of  $\text{Ob}(\widehat{M}^*)$  ..... 207
  - 5.9.5 Proof of Proposition 5.9.2 ..... 208
  - 5.9.6 Proof of Proposition 5.9.3 ..... 208
  - 5.9.7 Case in Which a 2-Stability Condition is Satisfied ..... 209
  - 5.9.8 Oriented Reduced  $L$ -Bradlow Pairs ..... 210
- 6 Virtual Fundamental Classes ..... 213**
  - 6.1 Perfectness of Obstruction Theories ..... 213
    - 6.1.1 Moduli Stacks of Semistable Objects ..... 213
    - 6.1.2 Master Spaces and Some Related Stacks ..... 214
    - 6.1.3 Vanishing of Some Cohomology Groups ..... 216
    - 6.1.4 Proof of the Propositions in Subsection 6.1.1 ..... 219
    - 6.1.5 Expected Dimension ..... 220
  - 6.2 Comparison of Oriented Reduced Case and Unoriented Unreduced Case ..... 221
    - 6.2.1 Statements ..... 221
    - 6.2.2 Proof of Proposition 6.2.1 ..... 223
  - 6.3 Rank One Case ..... 224
    - 6.3.1 Moduli of  $L$ -Abelian Pairs ..... 224
    - 6.3.2 Involutivity and Relation with Seiberg-Witten Invariants ..... 227
    - 6.3.3 Parabolic Hilbert Schemes ..... 228
    - 6.3.4 Splitting into Moduli of Abelian Pairs and Parabolic Hilbert Schemes ..... 231

6.3.5	Morphism to Moduli of Abelian Pairs	232
6.3.6	Morphism to Parabolic Hilbert Scheme	237
6.3.7	Mixed Case	239
6.3.8	Proof of Proposition 6.3.8	240
6.4	Bradlow Perturbation	242
6.4.1	Statements	242
6.4.2	Construction of $\mathfrak{B}$	244
6.4.3	Compatibility of the Obstruction Theories	244
6.4.4	Ambient Smooth Stack	248
6.5	Comparison with Full Flag Bundles	250
6.6	Parabolic Perturbation	252
6.6.1	Statement	252
6.6.2	Construction of a Stack $\mathfrak{B}$ with an Obstruction Theory	254
6.6.3	Compatibility of the Obstruction Theories	257
6.6.4	Smooth Ambient Stack	260
6.6.5	Proof of Proposition 6.6.1	261
<b>7</b>	<b>Invariants</b>	<b>263</b>
7.1	Preliminary	263
7.1.1	Ring $\mathcal{R}_l$	264
7.1.2	Homomorphisms	266
7.1.3	Evaluation	267
7.1.4	Equivariant Case	271
7.1.5	Ring $\mathcal{R}_{\text{CH}}$	273
7.1.6	Equivariant Euler Class	274
7.1.7	Twist by Line Bundle	275
7.2	Transition Formulas in Simpler Cases	276
7.2.1	Basic Case	276
7.2.2	Twist with the Euler Class of the Relative Tangent Bundle	279
7.2.3	Oriented Reduced $L$ -Bradlow Pairs	282
7.3	Invariants	286
7.3.1	Construction	286
7.3.2	Easy Property	289
7.3.3	Integrals over $\widetilde{\mathcal{M}}^{ss}(\widehat{y}, \alpha_*, +)$	290
7.3.4	Another Expression	290
7.3.5	Deformation Invariance	291
7.4	Rank Two Case	293
7.4.1	Reduction to the Sum of Integrals over the Products of Hilbert Schemes	293
7.4.2	Dependence on Polarizations	296
7.4.3	Proof of Theorem 7.4.3 (I)	298
7.4.4	Proof of Theorem 7.4.3 (II)	300
7.5	Higher Rank Case ( $p_g > 0$ )	303
7.5.1	Transition Formula in the Case $p_g > 0$	303

7.5.2	Reduction to the Sum of Integrals over the Products of Hilbert Schemes . . . . .	306
7.5.3	Independence from Polarizations in the Case $p_g > 0$ . . . . .	309
7.6	Transition Formula ( $p_g = 0$ ) . . . . .	311
7.6.1	Statement . . . . .	311
7.6.2	Step 1 . . . . .	313
7.6.3	Step 2 . . . . .	315
7.6.4	Step 3 . . . . .	318
7.7	Weak Wall Crossing Formula . . . . .	319
7.7.1	Statement . . . . .	319
7.7.2	Proof of Theorem 7.7.1 . . . . .	321
7.7.3	Weak Wall Crossing Formula in the Rank 3 Case . . . . .	323
7.7.4	Weak Intersection Rounding Formula in the Rank 3 Case . . . . .	325
7.7.5	Transition for a Critical Parabolic Weight . . . . .	330
7.8	Weak Intersection Rounding Formula . . . . .	331
7.8.1	Preliminary . . . . .	331
7.8.2	Statement . . . . .	335
7.8.3	Preparation from Combinatorics . . . . .	335
7.8.4	Proof of Theorem 7.8.2 . . . . .	337
	<b>References</b> . . . . .	341
	<b>Glossary</b> . . . . .	347
	<b>Index</b> . . . . .	379