SpringerBriefs in Earth System Sciences

South America and the Southern Hemisphere

Series Editors

Gerrit Lohmann

Lawrence A. Mysak

Justus Notholt

Jorge Rabassa

Vikram Unnithan

For further volumes:

<http://www.springer.com/series/10032>

Marcelo Reguero · Francisco Goin
Carolina Acosta Hospitaleche · Tania Dutra
Sergio Marensi

Late Cretaceous/ Paleogene West Antarctica Terrestrial Biota and Its Intercontinental Affinities

Marcelo Reguero
División Paleontología de Vertebrados
Museo de La Plata
La Plata
Argentina

Tania Dutra
SINOS-UNISINOS
Universidade do Vale do Rio dos
Sao Leopoldo
Rio Grande do Sul
Brazil

Francisco Goin
División Paleontología de Vertebrados
Museo de La Plata
La Plata
Argentina

Sergio Marenssi
Instituto Antártico Argentino
Buenos Aires
Argentina

Carolina Acosta Hospitaleche
División Paleontología de Vertebrados
Museo de La Plata
La Plata
Argentina

ISSN 2191-589X
ISBN 978-94-007-5490-4
DOI 10.1007/978-94-007-5491-1
Springer Dordrecht Heidelberg New York London

ISSN 2191-5903 (electronic)
ISBN 978-94-007-5491-1 (eBook)

Library of Congress Control Number: 2012945740

© The Author(s) 2013

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media (www.springer.com)

Preface

During the past two decades, geological explorations of the James Ross Basin, Weddell Sea, have revealed that this basin, located off the northeast tip of the Antarctic Peninsula (West Antarctica), contains one of the most important records of Late Cretaceous and Early Paleogene life in the Southern Hemisphere. The early explorer and scientist Otto Nordenskjöld (1905, p. 252), leader of the Swedish South Polar Expedition, envisioned the paleontologic and biogeographic importance of this basin “... *where, maybe, many animals and plants were first developed that afterwards found their way as far as to northern lands*”. These discoveries have not only provided new insights into the geologic history of Antarctica, but they have also provided answers to questions about life in Southern Hemisphere that have puzzled naturalists since Charles Darwin’s voyage on HMS *Beagle*.

The sedimentary sequence exposed in the James Ross Basin comprises a thick section of Coniacian, Turonian, Campanian, Maastrichtian, Paleocene, Eocene, and probably earliest Oligocene. They form the only marine sequence of this age interval that crops out in Antarctica. The high-latitude biota contained in both the Cretaceous and Paleogene beds is unusually rich and diverse and rivaled only by that from New Zealand and southeastern Australia.

After the Swedish South Polar Expedition (1901–1903), more than 40 years passed before the basin was scientifically visited again, this time by members of the Falkland Islands Dependencies Survey (now the British Antarctic Survey). The establishment of the Argentine station Marambio on Seymour Island in 1969 initiated the modern phase of geologic and paleontologic studies in the James Ross Basin.

The view that Antarctica and South America were connected by a long causeway between the West Antarctica (WANT) and southern South America in the Late Cretaceous through the late Paleocene, and that terrestrial vertebrates were able to colonize new frontiers using this physiographical feature, is almost certainly correct. One of the most intriguing palaeobiogeographical phenomena involving the last phase of the breakup of the supercontinent Gondwana concerns the close similarities and, in most cases, inferred sister-group relationships of a number of terrestrial vertebrate taxa recovered from uppermost Cretaceous and Paleogene deposits of Antarctic Peninsula (West Antarctica) and southern South

America (Magellanic Region and Patagonia). Dispersion of extinct vertebrates between South America and Australia across Late Cretaceous land bridges involving WAGG predicts their presence during that interval on the latter. However, although knowledge of Late Cretaceous vertebrates from West Antarctica (and the Antarctic Peninsula) is still severely limited, it has expanded substantially over the last two decades and includes discoveries of avian and non-avian dinosaurs. Plants and small- to medium-sized, obligate terrestrial mammals (e.g., marsupials and meridiungulates) gained broad distribution across West Antarctica land mass prior to fragmentation and were isolated on the Antarctic Peninsula before the end of the Paleocene.

From an evolutionary perspective the close relationships between terrestrial taxa found in the James Ross Basin are difficult to accommodate because, by the beginning of the Late Cretaceous (99.6 Ma), the major Gondwanan continental blocks (South America, Africa, Antarctica, Madagascar, the Indian subcontinent, and New Zealand–Australia) were well into the process of breakup and dispersion.

This paper deals with the hypothesis that the Antarctic Peninsula linked South America and West Antarctica in the Late Cretaceous until the beginning of the Paleogene and we review here through the plant and vertebrate fossil taxa the paleogeographical evolution of the southern continents and oceans for both earlier and later time intervals. This communication aims to explain the geological/geophysical problems inherent in the hypothesis and to explore the consequences for the biogeographical distributions of marine and terrestrial vertebrate faunas in West Antarctica.

Contents

1	Introduction	1
	References	5
2	West Antarctica: Tectonics and Paleogeography	9
2.1	East Antarctica/West Antarctica and Gondwanan Paleobiogeography	14
	References	15
3	Late Cretaceous/Paleogene Stratigraphy in the James Ross Basin . . .	19
3.1	Late Cretaceous	19
3.2	Paleogene	21
3.2.1	Cross Valley Formation	21
3.2.2	La Meseta Formation	22
	References	23
4	South America/West Antarctica: Pacific Affinities of the Weddellian Marine/Coastal Vertebrates	27
4.1	Late Cretaceous/Paleogene Marine Fossil Vertebrates of the James Ross Basin	29
4.1.1	Neoselachii and Teleostei Fossil Fishes	29
4.1.2	Marine Reptiles	34
4.1.3	Whales	35
4.2	Weddellian Sphenisciformes: Systematics, Stratigraphy, Biogeography and Phylogeny	36
	References	49
5	The Terrestrial Biotic Dimension of West Antarctica	55
5.1	West Antarctica Paleoflora	55
5.2	Late Cretaceous Terrestrial Vertebrates of the James Ross Basin . . .	65
5.2.1	Non Avian Dinosaurs	67
5.2.2	Avian Dinosaurs	72

5.3	Paleogene Terrestrial Vertebrates of the James Ross Basin	74
5.3.1	Gondwanatheres	77
5.3.2	Metatherians	77
5.3.3	Xenarthrans	84
5.3.4	“Insectivora”	85
5.3.5	Litopterns	86
5.3.6	Astrapotheres	88
5.4	Paleogene Environmental Reconstruction of the Cucullaea I (Ypresian) and Submeseta (Priabonian) Biotas	89
5.5	Correlation of the Cucullaea I Terrestrial Fauna with Early Paleogene Patagonian Faunas.	89
5.6	West Antarctic Terrestrial Biota and Its Intercontinental Relationships	97
	References	99
	Acknowledgments	111
	Appendix	113