Monographs in Mathematics
Vol. 88

Managing Editors:
H. Amann
Universität Zürich, Switzerland
K. Grove
University of Maryland, College Park
H. Kraft
Universität Basel, Switzerland
P.-L. Lions
Université de Paris-Dauphine, France

Associate Editors:
H. Araki, Kyoto University
J. Ball, Heriot-Watt University, Edinburgh
F. Brezzi, Università di Pavia
K.C. Chang, Peking University
N. Hitchin, University of Warwick
H. Hofer, ETH Zürich
H. Knörrer, ETH Zürich
K. Masuda, University of Tokyo
D. Zagier, Max-Planck-Institut Bonn

Roelof W. Bruggeman

Families of Automorphic Forms

1994 Springer Basel AG

Author:
Roelof W. Bruggeman
Mathematisch Instituut
Universiteit Utrecht
Postbus 80.010
3508 TA Utrecht
The Netherlands

A CIP catalogue record for this book is available from the Library of Congress,
Washington D.C., USA

Deutsche Bibliothek Cataloging-in-Publication Data
Bruggeman, Roelof W.:
Families of automorphic forms / Roelof W. Bruggeman.
(Monographs in mathematics ; Vol. 88)
ISBN 978-3-0346-0335-5 ISBN 978-3-0346-0336-2 (eBook)
DOI 10.1007/978-3-0346-0336-2
NE: GT

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, re-use of illustrations, broadcasting, reproduction on microfilms or in other ways, and storage in data banks. For any kind of use permission of the copyright owner must be obtained.

© 1994 Springer Basel AG

Originally published by Birkhäuser Verlag Basel, P.O. Box 133, CH-4010 Basel, Switzerland 1994
Softcover reprint of the hardcover 1st edition 1994

Camera-ready copy prepared by the author
Printed on acid-free paper produced of chlorine-free pulp

ISBN 978-3-0346-0335-5

9 8 7 6 5 4 3 2 1

Contents

Preface IX

1 Modular introduction

1.1 The modular group 1
1.2 Maass forms 3
1.3 Holomorphic modular forms 5
1.4 Fourier expansion 7
1.5 More modular forms 9
1.6 Truncation and perturbation 14
1.7 Further remarks 18

I General theory

2 Universal covering group

2.1 Upper half plane 25
2.2 Universal covering group 27
2.3 Automorphic forms 31

3 Discrete subgroups

3.1 Cofinite groups 33
3.2 The quotient 35
3.3 Canonical generators 37
3.4 Characters 42
3.5 Notations 44

4 Automorphic forms

4.1 Fourier expansion 48
4.2 Spaces of Fourier terms 51
4.3 Growth condition 58
4.4 Differentiation of Fourier terms 61
4.5 Differentiation of automorphic forms 65
4.6 Maass-Selberg relation 66

5 Poincaré series

5.1 Construction of Poincaré series 71
5.2 Fourier coefficients 78

6 Selfadjoint extension
6.1 Hilbert space 86
6.2 Energy subspace 88
6.3 Fourier coefficients 91
6.4 Compactness 94
6.5 Extension of the Casimir operator 97
6.6 Relation to automorphic forms 99
6.7 The discrete spectrum 101

7 Families of automorphic forms
7.1 Parameter spaces 107
7.2 Holomorphic families 109
7.3 Families of eigenfunctions 113
7.4 Automorphic transformation behavior 117
7.5 Families of automorphic forms 123
7.6 Families of Fourier terms 124
7.7 Differentiation 130

8 Transformation and truncation
8.1 Parameter space 133
8.2 Transformation 135
8.3 Truncation 138
8.4 Energy subspace 141
8.5 Families of automorphic forms 147

9 Pseudo Casimir operator
9.1 Sesquilinear form 151
9.2 Pseudo Casimir operator 156
9.3 Meromorphy of the resolvent 160
9.4 Meromorphic families 167
9.5 Dimension 174

10 Meromorphic continuation
10.1 Cells of continuation 177
10.2 Meromorphic continuation 180
10.3 Functional equations 185

11 Poincaré families along vertical lines
11.1 General results 191
11.2 Eisenstein families 198
11.3 Other Poincaré families 203

12 Singularities of Poincaré families
12.1 Local curves 213
12.2 Value sets 217
12.3 General results 221
12.4 General parameter spaces 227
12.5 Restricted parameter spaces 230

II Examples

13 Modular group
13.1 The covering group 239
13.2 Fourier expansions 244
13.3 The modular spectrum 247
13.4 Families of modular forms 249
13.5 Derivatives 252
13.6 Distribution results 258

14 Theta group
14.1 Theta group 265
14.2 The covering group 266
14.3 Fourier expansions 268
14.4 Eisenstein series 271
14.5 More than one parameter 274

15 Commutator subgroup
15.1 Commutator subgroup 275
15.2 Automorphic forms 278
15.3 The period map 280
15.4 Poincaré series 284
15.5 Eisenstein family of weight 0 286
15.6 Harmonic automorphic forms 290
15.7 Maass forms 302

References 307

Index 311

Preface

Automorphic forms on the upper half plane have been studied for a long time. Most attention has gone to the holomorphic automorphic forms, with numerous applications to number theory. Maass, [34], started a systematic study of real analytic automorphic forms. He extended Hecke's relation between automorphic forms and Dirichlet series to real analytic automorphic forms. The names Selberg and Roelcke are connected to the spectral theory of real analytic automorphic forms, see, e.g., [50], [51]. This culminates in the trace formula of Selberg, see, e.g., Hejhal, [21].

Automorphic forms are functions on the upper half plane with a special transformation behavior under a discontinuous group of non-euclidean motions in the upper half plane. One may ask how automorphic forms change if one perturbs this group of motions. This question is discussed by, e.g., Hejhal, [22], and Phillips and Sarnak, [46]. Hejhal also discusses the effect of variation of the multiplier system (a function on the discontinuous group that occurs in the description of the transformation behavior of automorphic forms). In [5]–[7] I considered variation of automorphic forms for the full modular group under perturbation of the multiplier system. A method based on ideas of Colin de Verdière, [11], [12], gave the meromorphic continuation of Eisenstein and Poincaré series as functions of the eigenvalue and the multiplier system jointly. The present study arose from a plan to extend these results to much more general groups (discrete cofinite subgroups of $\mathrm{SL}_2(\mathbb{R})$).

To carry this out I look at more general families of automorphic forms than one usually considers. In particular, I admit singularities inside the upper half plane, and relax the usual condition of polynomial growth at the cusps. This led me to reconsider a fairly large part of the theory of real analytic automorphic forms on the upper half plane.

This is done in Part I for arbitrary cofinite discrete groups. Chapters 2–6 discuss real analytic automorphic forms of a rather general type. Most results are known, or are easy extensions of known results. Chapters 7–12 consider families of these automorphic forms, with the eigenvalue and the multiplier system as the parameters. The ideas of Colin de Verdière are worked out in Chapters 8–10. The central result is Theorem 10.2.1; it gives the meromorphic continuation of Eisenstein and Poincaré series. The meromorphic continuation in the eigenvalue is well known; the meromorphy in all parameters jointly is new. Chapters 11 and 12 study singularities of the resulting families of automorphic forms. In Chapter 11 the eigenvalue is the sole variable. I summarize known results, and prepare for the study in Chapter 12 of the singularities in more than one variable. Table 1.1 on p. 18 gives a more detailed description of Part I.

The treatment in Part I is complicated. This is due to the fact that I consider general cofinite discrete groups, without restriction on the dimension of the group of multiplier systems. Chapter 1 is meant as an introduction. It explains the main ideas and results in the context of the full modular group. In the three chapters of Part II I consider three examples of cofinite discrete groups. Chapter 13 extends the discussion for the full modular group in Chapter 1. The other groups considered are the theta group and the commutator subgroup of the modular group. Although the first objective of Part II is to give the reader examples of the concepts, I have included some discussions that did not fit into the general context of Part I; see 1.7.7–1.7.10.

The reader I have had in mind has seen automorphic forms before, holomorphic as well as real analytic ones. For the latter I would suggest to have a look at Chapters IV and V of Maass's lecture notes [35], or at §3.5–7 of Terras's book [57]. The reader should also be prepared to look up facts concerning analytic functions in more than one complex variable, and be not afraid of a modest use of sheaf language when dealing with this subject.

The ideas of Colin de Verdière employed in Part I concern unbounded operators in Hilbert spaces. Kato's book, [25], is consulted for many results from functional analysis. I restrict the discussion of the spectral theory of automorphic forms to those results I need. In particular I do not mention the continuous part of the spectral decomposition.

R. Matthes visited Utrecht in 1989. The discussions we had gave me the stimulus to start this work. I am very grateful for this contribution, and also for the many comments he gave on early versions of this book. At a later stage the interest of D. Zagier has been a great encouragement. I also thank F. Beukers, J. Elstrodt, and B. van Geemen for corrections, comments, and suggestions.