

Editors-in-Chief:

J.-M. Morel, Cachan

B. Teissier, Paris

Advisory Board:

Camillo De Lellis (Zurich)

Mario Di Bernardo (Bristol)

Alessio Figalli (Pisa/Austin)

Davar Khoshnevisan (Salt Lake City)

Ioannis Kontoyiannis (Athens)

Gabor Lugosi (Barcelona)

Mark Podolskij (Heidelberg)

Sylvia Serfaty (Paris and NY)

Catharina Stroppel (Bonn)

Anna Wienhard (Heidelberg)

For further volumes:

<http://www.springer.com/series/304>

Tatsuo Nishitani

Hyperbolic Systems with Analytic Coefficients

Well-posedness of the Cauchy Problem



Springer

Tatsuo Nishitani
Department of Mathematics
Graduate School of Science
Osaka University
Toyonaka, Osaka, Japan

ISBN 978-3-319-02272-7 ISBN 978-3-319-02273-4 (eBook)
DOI 10.1007/978-3-319-02273-4
Springer Cham Heidelberg New York Dordrecht London

Lecture Notes in Mathematics ISSN print edition: 0075-8434
ISSN electronic edition: 1617-9692

Library of Congress Control Number: 2013955050

Mathematics Subject Classification (2010): 35L45, 35L40, 35L55

© Springer International Publishing Switzerland 2014

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media (www.springer.com)

Preface

In this monograph we discuss the C^∞ well-posedness of the Cauchy problem for hyperbolic systems. We are mainly concerned with the following two questions for differential operators of order q with smooth $m \times m$ matrix coefficients:

- (A) Under which conditions on lower order terms is the Cauchy problem C^∞ well posed?
- (B) When is the Cauchy problem C^∞ well posed for any lower order term?

For scalar case, that is $m = 1$, the question (B) has been answered. As for the question (A), in particular for second order scalar equations, that is $m = 1$ and $q = 2$, so many works are devoted to this question and the situation is fairly well understood. Contrary to the scalar case, for systems that is if $m \geq 2$ we have no satisfactory result.

Even for differential operators with characteristics of constant multiplicity with real analytic matrix coefficients, the question (A) has been solved very recently.

So in this monograph, assuming that the coefficients are real analytic in a neighborhood of the origin, we study these two questions. Of course this analyticity assumption is rather restrictive but which allows us to make detailed studies on the Cauchy problem. We hope that this study can throw light on the studies of the Cauchy problem for hyperbolic systems with less regular, in particular C^∞ coefficients.

The contents are organized as follows. In Chap. 1 after giving the definition of C^∞ well-posedness of the Cauchy problem we show that the Cauchy problem for symmetric hyperbolic systems is C^∞ well posed for any lower order term. Then we give an example of first order 2×2 system which is not symmetrizable but for which the Cauchy problem is C^∞ well posed for any lower order term. Actually there is a class of non-symmetrizable systems for which the Cauchy problem is C^∞ well posed for any lower order term. This is a main objection when we try to answer to the problem (B). We prove the Lax-Mizohata theorem exhibiting naive ideas which are used in Chaps. 2 and 3. For first order systems with characteristics of constant multiplicities, the necessity of the Levi condition for the C^∞ well-posedness is proved which is used in Chap. 2. In Chap. 2, we study necessary conditions about the

problem (B) for $m \times m$ first order systems with real analytic coefficients. We prove rather general necessary conditions in terms of minors of the principal symbols. Contrary to the scalar case the multiplicity of characteristics is irrelevant for the problem (B) since for symmetric or symmetrizable hyperbolic systems (of first order) the Cauchy problem is always C^∞ well posed for any lower order term. Here the maximal size of the Jordan blocks, which is supposed to measure the distance from diagonal matrices, plays an important role in the problem (B).

In Chap. 3, we study two questions (A) and (B) for first order 2×2 systems with two independent variables with real analytic coefficients. For this special case we can give a necessary and sufficient condition for the questions (A) and (B), that is, in this case we have complete answer for (A) and (B). The results provide many instructive examples. For instance, we can exhibit a first order 2×2 system with analytic coefficients which is strictly hyperbolic outside the initial line for which no lower order term could be taken so that the Cauchy problem is C^∞ well posed. This cannot happen for second order hyperbolic scalar operators with two independent variables with analytic coefficients.

In Chap. 4, we introduce a new class of hyperbolic systems, that is hyperbolic systems with nondegenerate characteristics which generalizes strictly hyperbolic systems. Strictly hyperbolic systems are hyperbolic systems with nondegenerate characteristics of order one. The theory of strictly hyperbolic systems is rich, but first order strictly hyperbolic system hardly exists. We prove that the Cauchy problem for hyperbolic systems with nondegenerate characteristics is C^∞ well posed for any lower order term. We also show that nondegenerate characteristics are stable, that is any hyperbolic system which is close to a hyperbolic system with a nondegenerate characteristic of order r has a nondegenerate characteristic of the same order nearby. This shows, in particular, that near any hyperbolic system with a nondegenerate characteristic of order $r \geq 2$ there is no strictly hyperbolic system, which gives a great difference from the scalar case and shows a complexity of hyperbolic systems.

We also discuss hyperbolic systems which are perturbations of symmetric systems and prove that if the dimension of the linear space that the symbol of the symmetric system spans is large enough, then generically such hyperbolic system is similar to a symmetric system.

Osaka, Japan
March 2013

Tatsuo Nishitani

Contents

1	Introduction	1
1.1	Well-Posedness of the Cauchy Problem	1
1.2	Symmetric Hyperbolic Systems	5
1.3	Systems Which Are Not Symmetrizable	12
1.4	Lax-Mizohata Theorem	18
1.5	Levi Condition	22
1.6	A Lemma on Hyperbolic Polynomials	26
2	Necessary Conditions for Strong Hyperbolicity	31
2.1	Necessary Conditions for Strong Hyperbolicity	31
2.2	Key Propositions	33
2.3	Proof of Theorem 2.1 (Simplest Case)	37
2.4	Proof of Theorem 2.1 (General Case)	44
2.5	Proofs of Propositions 2.4 and 2.5	50
2.6	Proof of Key Proposition	55
2.7	Proof of Key Proposition, Asymptotic Diagonalization	63
2.8	Involutive Characteristics	71
2.9	Localization at Involutive Characteristics	75
2.10	Concluding Remarks	82
3	Two by Two Systems with Two Independent Variables	85
3.1	Reduction to Almost Diagonal Systems	85
3.2	Nonnegative Real Analytic Functions	90
3.3	Well-Posedness and Pseudo-Characteristic Curves	92
3.4	Strongly Hyperbolic 2×2 Systems	96
3.5	Nonnegative Functions and Newton Polygons	98
3.6	Behavior Around Pseudo-Characteristic Curves	104
3.7	Proof of Proposition 3.2	108
3.8	Energy Estimates Near Pseudo-Characteristic Curves	112
3.9	Energy Estimates of Higher Order Derivatives	121
3.10	Weighted Energy Estimates	125
3.11	Conditions for Well-Posedness	136

3.12	Construction of Asymptotic Solutions	142
3.13	Proof of Necessity	150
3.14	Equivalence of Conditions	153
3.15	Concluding Remarks	160
4	Systems with Nondegenerate Characteristics	161
4.1	Nondegenerate Characteristics	161
4.2	Nondegenerate Double Characteristics	169
4.3	Symmetrizability (Special Case)	174
4.4	Stability and Smoothness of Nondegenerate Characteristics	179
4.5	Symmetrizability (General Case)	194
4.6	Well Posed Cauchy Problem	209
4.7	Nondegenerate Characteristics of Symmetric Systems	213
4.8	Hyperbolic Perturbations of Symmetric Systems	216
4.9	Stability of Symmetric Systems Under Hyperbolic Perturbations	221
4.10	Some Special Cases	225
4.11	Concluding Remarks	229
	References	231
	Index	235