

Editors-in-Chief:

J.-M. Morel, Cachan

B. Teissier, Paris

Advisory Board:

Camillo De Lellis, Zurich

Mario di Bernardo, Bristol

Alessio Figalli, Zurich

Davar Khoshnevisan, Salt Lake City

Ioannis Kontoyiannis, Athens

Gabor Lugosi, Barcelona

Mark Podolskij, Aarhus

Sylvia Serfaty, Paris and New York

Catharina Stroppel, Bonn

Anna Wienhard, Heidelberg

Fondazione C.I.M.E., Firenze



C.I.M.E. stands for *Centro Internazionale Matematico Estivo*, that is, International Mathematical Summer Centre. Conceived in the early fifties, it was born in 1954 in Florence, Italy, and welcomed by the world mathematical community: it continues successfully, year for year, to this day.

Many mathematicians from all over the world have been involved in a way or another in C.I.M.E.'s activities over the years. The main purpose and mode of functioning of the Centre may be summarised as follows: every year, during the summer, sessions on different themes from pure and applied mathematics are offered by application to mathematicians from all countries. A Session is generally based on three or four main courses given by specialists of international renown, plus a certain number of seminars, and is held in an attractive rural location in Italy.

The aim of a C.I.M.E. session is to bring to the attention of younger researchers the origins, development, and perspectives of some very active branch of mathematical research. The topics of the courses are generally of international resonance. The full immersion atmosphere of the courses and the daily exchange among participants are thus an initiation to international collaboration in mathematical research.

C.I.M.E. Director (2002 – 2014)

Pietro Zecca
Dipartimento di Energetica “S. Stecco”
Università di Firenze
Via S. Marta, 3
50139 Florence
Italy
e-mail: zecca@unifi.it

C.I.M.E. Director (2015 –)

Elvira Mascolo
Dipartimento di Matematica “U. Dini”
Università di Firenze
viale G.B. Morgagni 67/A
50134 Florence
Italy
e-mail: mascolo@math.unifi.it

C.I.M.E. Secretary

Paolo Salani
Dipartimento di Matematica “U. Dini”
Università di Firenze
viale G.B. Morgagni 67/A
50134 Florence
Italy
e-mail: salani@math.unifi.it

CIME activity is carried out with the collaboration and financial support of INdAM (Istituto Nazionale di Alta Matematica)
For more information see CIME's homepage:
<http://www.cime.unifi.it>

Michel Boileau • Gerard Besson •
Carlo Sinestrari • Gang Tian

Ricci Flow and Geometric Applications

Cetraro, Italy 2010

Riccardo Benedetti, Carlo Mantegazza
Editors



Authors

Michel Boileau
Aix-Marseille Université, CNRS, Central
Marseille
Institut de Mathématiques de Marseille
Marseille, France

Gerard Besson
Institut Fourier
Université Grenoble Alpes
Grenoble, France

Carlo Sinestrari
Dip. di Ingegneria Civile e Ingegneria
Informatica
Università di Roma “Tor Vergata”
Rome, Italy

Gang Tian
Princeton University
Princeton, NJ
USA

Editors

Riccardo Benedetti
Department of Mathematics
University of Pisa
Pisa, Italy

Carlo Mantegazza
Department of Mathematics
University of Naples
Naples, Italy

ISSN 0075-8434

ISSN 1617-9692 (electronic)

Lecture Notes in Mathematics

ISBN 978-3-319-42350-0

ISBN 978-3-319-42351-7 (eBook)

DOI 10.1007/978-3-319-42351-7

Library of Congress Control Number: 2016951889

Mathematics Subject Classification (2010): 53C44, 57M50, 57M40

© Springer International Publishing Switzerland 2016

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

This Springer imprint is published by Springer Nature

The registered company is Springer International Publishing AG Switzerland

Preface

Our aim in organizing this CIME course was to present to young students and researchers the impressive recent achievements in differential geometry and topology obtained by means of techniques based on the Ricci flow. We then invited some of the leading researchers in the field of geometric analysis and low-dimensional geometry/topology to introduce some of the central ideas in their work. Here is the list of speakers together with the titles of their lectures:

- Gérard Besson (Grenoble) – *The differentiable sphere theorem (after S. Brendle and R. Schoen)*
- Michel Boileau (Toulouse) – *Thick/thin decomposition of three-manifolds and the geometrization conjecture*
- Carlo Sinestrari (Roma “Tor Vergata”) – *Singularities of three-dimensional Ricci flows*
- Gang Tian (Princeton) – *Kähler–Ricci flow and geometric applications.*

The summer school had around 50 international attendees (mostly PhD students and postdocs). Even though they were sometimes technically heavy, the lectures were followed by all the students with interest. The participants were very satisfied by the high quality of the courses. The not-so-intense scheduling of the lectures gave the students many opportunities to interact with the speakers, who were always very friendly and available for discussion. It should be mentioned that the wonderful location and the careful CIME organization were also greatly appreciated.

We think that the fast-growing field of geometric flows and more generally of geometric analysis, which has always received great attention in the international community, but which is still relatively “young” in Italy, will benefit from its diffusion by this CIME course.

We briefly describe the contents of the lectures collected in this volume.

Gérard Besson presented the impact of the Ricci flow technique on the theory of positively curved manifolds, the central result being the *differentiable 1/4-pinched sphere theorem*, proved by Brendle and Schoen. It says that a complete, simply

connected Riemannian manifold whose sectional curvature varies in $(1/4, 1]$ is diffeomorphic to the standard sphere.

The problem was first proposed by H. Hopf, and then in 1951, H.E. Rauch showed that a complete Riemannian manifold whose sectional curvature is positive and varies between two numbers whose ratio is close to 1 has a universal cover homeomorphic to a sphere. In the 1960s, M. Berger and W. Klingenberg obtained the optimal result: a simply connected Riemannian manifold which is *strictly* $1/4$ -pinched is *homeomorphic* to the sphere. The analogous *diffeomorphic* conclusion remained open until S. Brendle and R. Schoen proved the following:

Theorem (S. Brendle and R. Schoen, 2008) *Let M be a pointwise strictly $1/4$ -pinched Riemannian manifold of positive sectional curvature. Then M carries a metric of constant sectional curvature. Hence, it is diffeomorphic to the quotient of a sphere by a finite subgroup of $O(n)$.*

The proof relies on the use of the Ricci flow introduced by R. Hamilton and culminating in the work of G. Perelman. The idea is to construct a deformation of the Riemannian metric, evolving it by means of the Ricci flow toward a constant curvature metric. We recall that this was the method that R. Hamilton used in his seminal paper, proving the following theorem:

Theorem (R. Hamilton, 1982) *Let M be a closed 3-dimensional Riemannian manifold which carries a metric of positive Ricci curvature; then it also carries a metric of positive constant curvature.*

The lectures also focus on the extension to higher dimensions of the following result, due to C. Böhm and B. Wilking. Recall that a curvature operator is 2-positive if the sum of its two smallest eigenvalues is positive.

Theorem (C. Böhm and B. Wilking, 2008) *Let M be a closed Riemannian manifold whose curvature operator is 2-positive; then M carries a constant curvature metric.*

In the lectures, the connection between this method and the algebraic properties of the Riemann curvature operator is stressed, the main focus being the identification of those properties of the curvature operator which are preserved under the Ricci flow.

In his lectures, Michel Boileau gave an introduction to the geometrization of 3-manifolds. Sections 2.1 and 2.2 cover Thurston's classification of the eight 3-dimensional geometries and the characterization of *geometric* (and *Seifert*) closed 3-manifolds in terms of basic topological properties. This follows by combining Thurston's hyperbolization theorems (in particular the characterization of hyperbolic 3-manifolds that are fibered over \mathbb{S}^1), Perelman's general geometrization theorem, and Agol's recent (2013) proof of a deep conjecture of Thurston that closed hyperbolic 3-manifolds are "virtually fibered."

Section 2.3 discusses the following: (1) A central result of classical 3-dimensional geometric topology, that is, the *canonical decomposition* of a

3-manifold by splitting it along spheres and tori. (2) *Thurston's geometrization conjecture*. This roughly says that every piece of a canonical decomposition is geometric together with a prediction on the geometry carried by the piece in terms of basic topological properties. It includes as a particular case the celebrated *Poincaré conjecture*. (3) Thurston's fundamental *hyperbolization theorem for Haken manifolds*.

Perelman's proof of the general *geometrization theorem* deals with all of these topics and also allows us to recover, as a by-product, the canonical decomposition itself. This is done by completing the program based on the *Ricci flow with surgeries*, first proposed by R. Hamilton. This is the subject of Boileau's notes from Sect. 2.4.

Since the appearance of Perelman's three celebrated preprints, several simplifications and variants of the original proofs have been developed by various authors. At the end of the day, we can say that the *Poincaré conjecture* (i.e., the case when the Ricci flow with surgery becomes extinct in finite time) is in a sense the "simplest" case. The general case (when the Ricci flow with surgery exists at all times, which includes the complete hyperbolization theorem) requires nontrivial extra arguments, in particular, to obtain a key *non-collapsing theorem*. In Perelman's original work, these come from the theory of *Alexandrov spaces*. Bessi eres, Besson, Boileau, Maillot, and Porti developed instead an alternative approach where the basic tools are Thurston's hyperbolization theorem for Haken manifolds and some well-established properties of Gromov's *simplicial volume*, allowing one to bypass the need for the (somewhat more exotic) theory of Alexandrov spaces. Boileau's notes are largely based on the monograph by

L. Bessi eres, G. Besson, M. Boileau, S. Maillot, and J. Porti, *Geometrisation of 3-Manifolds*, EMS Tracts in Mathematics **13**, 2010.

In this tract, the authors developed a slightly different notion of surgery by defining the so-called Ricci flow with bubbling-off. Actually, one might roughly say that the Ricci flow with bubbling-off reduces the general hyperbolization theorem to Thurston's hyperbolization theorem for Haken manifolds.

Carlo Sinestrari provided an extensive introduction to the Ricci flow by first giving a survey of the basic results and examples, then concentrating on the analysis of the singularities of the flow in the three-dimensional case, which is needed in Hamilton and Perelman's surgery construction. After reviewing the properties of the Ricci flow and the fundamental estimates of the theory, such as Hamilton's Harnack differential inequality, the Hamilton–Ivey pinching estimate, and Perelman's no collapsing result, he presented Perelman's analysis of kappa-solutions and the canonical neighborhood property which gives a full description of the singular behavior of the solutions in dimension 3. All these results are central to the proof of the Poincar e and geometrization conjectures.

The exposition is quite accessible to nonexperts. Indeed, the presentation is often informal, and the proofs are omitted except in some simple and significant cases, focusing more on the description of the results and their applications and consequences. A final detailed bibliographical section gives to the interested reader all the references needed for an advanced study of these topics.

Gang Tian's expository notes, based on his lectures, discuss some aspects of the Analytic Minimal Model Program through the Kähler–Ricci flow, developed in collaboration with other authors, particularly, J. Song and Z. Zhang. Very stimulating open problems and conjectures are also presented.

Section 4.2 contains a detailed account of the sharp version of the Hamilton–DeTurck local existence theorem, which holds in the Kähler case: the maximal time $T_{\max} \in (0, +\infty]$ such that the flow exists on the interval $[0, T_{\max})$ is precisely determined in terms of a cohomological property of the initial Kähler metric. As a corollary, one deduces that $T_{\max} = +\infty$ for every initial metric on a compact Kähler manifold with a numerically positive canonical bundle.

In Sect. 4.3, the limit singularities that can arise when $t \rightarrow T_{\max} < +\infty$ are analyzed. After having established a general convergence theorem (Theorem 4.3.1), one faces questions concerning regularity and the geometric properties of the limit. A combination of partial results (particularly in the case of projective varieties and when one can apply deep results of algebraic geometry) and well-motivated conjectures outlines a pregnant scenario.

Sections 4.4 and 4.5 discuss the construction of a *Kähler–Ricci flow with surgery* (assuming the truth of a conjecture stated in Sect. 4.3) and its asymptotic behavior. Numerous conjectures arise throughout this discussion such as: the characterization (up to birational isomorphism) of “Fano-like” manifolds as those whose flow becomes extinct at a finite time; the characterization of uniruled manifolds (up to birational isomorphism) as those whose flow collapses in finite time; and the existence of a flow with surgery globally defined in time and with only finitely many surgery times.

In Sect. 4.6, algebraic surfaces are considered, showing how most of the program is carried out in this case.

We are pleased to express our thanks to the speakers for their excellent lectures and to the participants for contributing with their enthusiasm to the success of the Summer School.

The speakers, the participants, and the CIME organizers collectively created a stimulating, rich, pleasant, and friendly atmosphere at Cetraro. For this reason, we would finally like to thank the Scientific Committee of CIME and, in particular, Pietro Zecca and Elvira Mascolo.

Pisa, Italy
Naples, Italy

Riccardo Benedetti
Carlo Mantegazza

Acknowledgements

CIME activity is carried out with the collaboration and financial support of: INdAM (Istituto Nazionale di Alta Matematica) and MIUR (Ministero dell'Istruzione, dell'Università e della Ricerca).

Contents

1	The Differentiable Sphere Theorem (After S. Brendle and R. Schoen).....	1
	G�rard Besson	
2	Thick/Thin Decomposition of Three-Manifolds and the Geometrisation Conjecture.....	21
	Michel Boileau	
3	Singularities of Three-Dimensional Ricci Flows.....	71
	Carlo Sinestrari	
4	Notes on K�hler-Ricci Flow.....	105
	Gang Tian	