

Editors:

J.-M. Morel, Cachan

F. Takens, Groningen

B. Teissier, Paris

K. Habermann · L. Habermann

Introduction to Symplectic Dirac Operators

 Springer

Authors

Katharina Habermann
State and University Library Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen
Germany
e-mail: habermann@sub.uni-goettingen.de

Lutz Habermann
Department of Mathematics
University of Hannover
Welfengarten 1
30167 Hannover
Germany
e-mail: habermann@math.uni-hannover.de

Library of Congress Control Number: 2006924123

Mathematics Subject Classification (2000): 53-02, 53Dxx, 58-02, 58Jxx

ISSN print edition: 0075-8434

ISSN electronic edition: 1617-9692

ISBN-10 3-540-33420-3 Springer Berlin Heidelberg New York

ISBN-13 978-3-540-33420-0 Springer Berlin Heidelberg New York

DOI 10.1007/b138212

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer. Violations are liable for prosecution under the German Copyright Law.

Springer is a part of Springer Science+Business Media
springer.com

© Springer-Verlag Berlin Heidelberg 2006
Printed in The Netherlands

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

Typesetting: by the authors and SPI Publisher Services using a Springer L^AT_EX package
Cover design: *design & production* GmbH, Heidelberg

Printed on acid-free paper SPIN: 11736684 VA41/3100/SPI 543210

To Karen

Preface

This book aims to give a systematic and self-contained introduction to the theory of symplectic Dirac operators and to reflect the current state of the subject. At the same time, it is intended to establish the idea that symplectic spin geometry and symplectic Dirac operators may give valuable tools in symplectic geometry and symplectic topology, which have become important fields and very active areas of mathematical research.

The basic idea of symplectic spin geometry goes back to the early 1970s, when Bertram Kostant introduced symplectic spinors in order to give the construction of the half-form bundle and the half-form pairings in the context of geometric quantization [37]. During the next two decades, however, almost no attention has been given to a closer study of symplectic spin geometry itself.

In 1995, the first author introduced symplectic Dirac operators [24] and started a systematical investigation [25, 26, 27]. These *symplectic Dirac operators* are called *Dirac operators*, since they are defined in an analogous way as the classical Riemannian Dirac operator known from Riemannian spin geometry (cf. e.g. [21]). They are called *symplectic* because they are constructed by use of the symplectic setting of the underlying symplectic manifold. All tools which were necessary for that construction have already been known and accepted, mainly in mathematical physics. These are the symplectic Clifford algebra (also known as Weyl algebra), the metaplectic group, the metaplectic representation (Segal–Shale–Weil representation) acting on $L^2(\mathbb{R}^n)$, metaplectic structures, and symplectic connections.

One of the basic ideas in differential geometry is that the study of analytic properties of certain differential operators acting on sections of vector bundles yields geometric and topological properties of the underlying base manifold. There are several classical results in that direction. An example is Hodge–de Rham theory. Here, one considers the Hodge–Laplace–Beltrami operator Δ

acting on differential forms. This operator is one of the most studied operators in global Riemannian geometry and his spectrum gives important topological invariants. In particular, the dimension of the kernel of Δ on p -forms over a closed Riemannian manifold is the p -th Betti number. Other well known and well studied operators are the Kodaira–Hodge–Laplace operator on differential forms with values in a holomorphic vector bundle or the classical Dirac operator on Riemannian manifolds. Now, symplectic spinor fields are sections in an $L^2(\mathbb{R}^n)$ -Hilbert space bundle over a symplectic manifold and symplectic Dirac operators, acting on symplectic spinor fields, are associated to the symplectic manifold in a very natural way. Hence they may be expected to give interesting applications in symplectic geometry and symplectic topology.

It is our opinion that, besides the already stated, there are further close relations to mathematical physics. Some steps towards this direction have been made by the first author and Andreas Klein in [28, 29, 30].

Another perspective could be the extension of Clifford analysis and spin geometry to super differential geometry. According to the most used version of super geometry developed by Bertram Kostant, geometrical structures over a supermanifold consist of \mathbb{Z}_2 -graded objects and thus have an even as well as an odd part. Then one can imagine that a metric has to satisfy some kind of graded symmetry which, roughly speaking, corresponds to a symmetric object on the even part as well as to a skew symmetric object on the odd part of the supermanifold. For the first one, we have the classical Riemannian spin geometry, whereas the second one is basically given by symplectic spin geometry. The main aspects of that idea are treated in a paper by Frank Sommen [45].

Although the construction of symplectic Dirac operators follows the same procedure as for the classical Riemannian Dirac operator, using the symplectic structure of the underlying manifold instead of the Riemannian metric, there are essential differences to the Riemannian case. These are caused by the fact that the algebraic structure of the symplectic Clifford algebra is completely different from that of Riemannian spin geometry. For the classical Clifford algebra, we have the relation $\mathbf{v}^2 = -\|\mathbf{v}\|^2$, whereas the algebraic structure of the symplectic Clifford algebra is given by $\mathbf{v} \cdot \mathbf{w} - \mathbf{w} \cdot \mathbf{v} = -\omega_0(\mathbf{v}, \mathbf{w})$. This implies essentially different properties for the Clifford multiplications, which enter into the definition of the Dirac operators.

Moreover, the non-compactness of the symplectic group leads to analytic difficulties. Namely, since the typical fiber of the symplectic spinor bundle is the Hilbert space $L^2(\mathbb{R}^n)$, we deal with operators acting on sections of a vector bundle of infinite rank. For elliptic formally self-adjoint pseudo-differential operators with positive definite leading symbol acting on sections in a vector bundle of finite rank, one has a completely developed theory. So, in order to be able to apply these techniques, we are interested in equivariance properties

of our operators with respect to a certain decomposition of the symplectic spinor bundle into a series of subbundles of finite rank. It turns out that an associated second order operator respects this decomposition provided that a technical assumption, which always can be realized, holds true.

Let us now briefly describe the content of each chapter of this book. The first chapter is introductory. It contains preliminaries and basic material needed for our considerations. Chapter 2 is devoted to symplectic connections. In particular, we introduce a further Ricci tensor, which we will call symplectic Ricci tensor. To our knowledge, no attention has been given to this tensor in previous studies. In fact, this symplectic Ricci tensor is a new object in the case of non-vanishing torsion. To date, mostly only torsion-free symplectic connections have been considered. It turns out that, in our context, it is convenient to work also with symplectic connections with torsion and that the symplectic Ricci tensor is more suitable for our purposes. The next chapter introduces the symplectic spinor bundle and the spinor derivative and analyzes a splitting property of the spinor bundle. In Chapter 4, we give the definition of symplectic Dirac operators and describe in detail how these operators depend on the objects from which they are built. Chapter 5 is concerned with an associated second order operator of Laplace type and addresses properties of this operator. The objective of Chapter 6 is the situation for a special class of symplectic manifolds, namely Kähler manifolds. Here, we also investigate the example of $\mathbb{C}P^1$. The aim of Chapter 7 is to construct a Fourier transform for symplectic spinor fields and to derive consequences for the symplectic Dirac operators. The last chapter focuses on relations to mathematical physics, in particular to quantization. This closes the circle to the beginnings by Bertram Kostant.

The present text is originated in research ideas of the first author and provides an extended version of her “Habilitationsschrift”, which was never published separately. Starting with helpful discussions from the beginning of the investigations and proposing many improvements in the selection and presentation of the material, the second author became more and more involved into the subject. We decided to write this book three years ago and, for this book, he made a thorough revision of the material, in particular, to improve the strictness of the presentation. Then it took time to compose it in a natural and organic way. Furthermore, our working places are no longer as close to each other as before and it became difficult to keep our discussions at its intensive level. Now we consider the text ready for publication and hope to present the reader a mature work.

During the work on the book, we received financial support from DFG, the German Research Foundation, contract HA 3056/1-1,2. Many thanks are due to our students Paul Rosenthal and Steffen Rudnick for proof-reading the L^AT_EX-type-written manuscript.

We dedicate this book to our daughter Karen. She accepts our mathematical family circle and enriches it with her interest. We are grateful to being able to give her an understanding of the fascination of mathematics. Karen, you are a wonderful teenager which makes us enjoying the challenge of teaching mathematics.

Göttingen and Greifswald,
November 2004

Katharina & Lutz Habermann

Contents

1	Background on Symplectic Spinors	1
1.1	Symplectic Group and Clifford Algebra	1
1.2	The Stone–von Neumann Theorem	5
1.3	Metaplectic Representation	7
1.4	Symplectic Clifford Multiplication	11
1.5	Hermite Functions	16
2	Symplectic Connections	21
2.1	Symplectic Manifolds	21
2.2	Constructions and Torsion	25
2.3	Symplectic Curvature and Ricci Tensors	29
3	Symplectic Spinor Fields	35
3.1	Metaplectic Structures	35
3.2	Symplectic Spinor Bundle	37
3.3	Splitting of the Spinor Bundle	43
4	Symplectic Dirac Operators	49
4.1	Definition of the Operators	49
4.2	Dependence on the Symplectic Connection	52
4.3	Dependence on the Metaplectic Structure	57
4.4	Dependence on the Almost Complex Structure	62
4.5	Formal Self-Adjointness	64

5	An Associated Second Order Operator	67
5.1	Definition and Ellipticity	67
5.2	A Weitzenböck Formula	68
5.3	Splitting of the Operator	74
6	The Kähler Case	81
6.1	The Operator \mathcal{P} on Kähler Manifolds	81
6.2	Lower Bound Estimates	86
6.3	The Spectrum of \mathcal{P} on $\mathbb{C}P^1$	87
7	Fourier Transform for Symplectic Spinors	97
7.1	Definition of the Transform	97
7.2	Basic Properties	98
7.3	Symmetry of the Spectra of \mathcal{D} and $\tilde{\mathcal{D}}$	99
8	Lie Derivative and Quantization	101
8.1	Lie Derivative of Symplectic Spinor Fields	101
8.2	Schrödinger Equation for Quadratic Hamiltonians	109
8.3	Lie Derivative as Quantization	111
	References	115
	Index	119