

Lecture Notes in Mathematics

2016

Editors:

J.-M. Morel, Cachan

B. Teissier, Paris

For further volumes:

<http://www.springer.com/series/304>

Jan Lang • David Edmunds

Eigenvalues, Embeddings and Generalised Trigonometric Functions

Prof. Jan Lang
Ohio State University
Department of Mathematics
231 West 18th Avenue
Columbus, Ohio 43210
USA
lang@math.ohio-state.edu

Prof. David Edmunds
University of Sussex
Department of Mathematics
Pevensey 2, North-South Road
Brighton BN1 9QH
United Kingdom
davideedmunds@aol.com

ISBN 978-3-642-18267-9 e-ISBN 978-3-642-18429-1
DOI 10.1007/978-3-642-18429-1
Springer Heidelberg Dordrecht London New York

Lecture Notes in Mathematics ISSN print edition: 0075-8434
ISSN electronic edition: 1617-9692

Library of Congress Control Number: 2011924532

Mathematics Subject Classification (2011): 41A35, 41A46, 47B06, 33E30, 47G10, 35P05, 47A75,
35P15, 46E35, 47B05

© Springer-Verlag Berlin Heidelberg 2011

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer. Violations are liable to prosecution under the German Copyright Law.

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

Cover design: deblik, Berlin

Printed on acid-free paper

Springer is part of Springer Science+Business Media (www.springer.com)

Preface

The main theme of these notes is the study, from the standpoint of s -numbers, of operators of Hardy type and related Sobolev embeddings. More precisely, let $p, q \in (1, \infty)$ and suppose that I is the interval (a, b) , where $-\infty < a < b < \infty$. Maps $T : L_p(I) \rightarrow L_q(I)$ of the form

$$(Tf)(x) = v(x) \int_a^x u(t)f(t)dt, \quad (1)$$

where u and v are prescribed functions satisfying some integrability conditions, are said to be of Hardy type. They are of importance in connection with ‘small ball’ problems in probability theory [87] and also in the theory of embeddings of Sobolev spaces when the underlying subset Ω of \mathbb{R}^n is a generalised ridged domain, which means crudely that Ω has a central axis (the generalised ridge) that is the image of a tree under a Lipschitz map [42]. In addition, the literature on such maps T has grown to such an extent that the topic has acquired an independent life. Our object is, so far as we are able, to give an account of the present state of knowledge in this area in the hope that it will stimulate further work. In addition to the main theme, topics that arise naturally include the geometry of Banach spaces, generalised trigonometric functions and the p -Laplacian, and we have not hesitated to develop these subsidiary melodies beyond the strict requirements of Hardy operators when the intrinsic interest warranted it. We hope that the resulting contrapuntal effect will appeal to the reader.

Chapter 1 supplies basic information about bases of Banach spaces and such geometric concepts as strict and uniform convexity, uniform smoothness and super-reflexivity. It also gives an account of very recent work (see [44]) on the representation of compact linear operators $S : X \rightarrow Y$, where X and Y are reflexive Banach spaces with strictly convex duals. What emerges is the existence of a sequence (x_n) in the unit sphere of X and a sequence (λ_n) of positive numbers in terms of which the action of S can be described and points $x \in X$ represented, under suitable conditions; the λ_n are norms of the restrictions of S to certain subspaces. These results provide an analogue in Banach spaces of the celebrated Hilbert space results of Erhard Schmidt. As a byproduct we have (in Chap. 3) a proof of the existence of an infinite sequence of ‘eigenvectors’ of the Dirichlet problem for the p -Laplacian in an arbitrary bounded domain in \mathbb{R}^n .

The next chapter gives an account of generalised trigonometric functions. To explain what is involved here, let $p \in (1, \infty)$, put

$$\pi_p = \frac{2\pi}{p \sin(\pi/p)}$$

and let $F_p : [0, 1] \rightarrow \mathbb{R}$ be given by

$$F_p(x) = \int_0^x (1 - t^p)^{-1/p} dt.$$

Then the generalised sine function \sin_p is the function defined on $[0, \pi_p/2]$ to be the inverse of F_p and extended to the whole of \mathbb{R} in a natural way so as to be $2\pi_p$ -periodic. Plainly $\sin_2 = \sin$. Moreover, p -analogues of the other trigonometric functions may easily be given: for example, \cos_p is defined to be the derivative of \sin_p , from which it follows quickly that

$$|\sin_p x|^p + |\cos_p x|^p = 1 \text{ for all } x \in \mathbb{R}.$$

After establishing the main properties of these p -functions and some of the identities obtainable by their use, such as a new representation of the Catalan constant, the chapter finishes with a proof of the fact (first given in [9]) that if p is not too close to 1, then the functions $\sin_p(n\pi_p t)$ form a basis in $L_q(0, 1)$ for all $q \in (1, \infty)$. The usefulness of such p -functions is underlined in Chap. 3, where it is shown how \sin_p and \cos_p arise naturally in the study of initial- and boundary-value problems for the one-dimensional p -Laplacian on an interval.

Chapter 4 provides necessary and sufficient conditions for the boundedness and compactness of the Hardy operator T of (1) acting between Lebesgue spaces. The norm of T_0 , the particular form of T when $u = v = 1$, is determined explicitly and is shown to be attained at functions expressible in terms of generalised trigonometric functions. After this preparation, Chap. 5 is devoted to the s -numbers of T_0 , together with the calculation of s -numbers of the basic Sobolev embedding on intervals. We remind the reader that in the theory of s -numbers, to every bounded linear map $S : X \rightarrow Y$, where X and Y are Banach spaces, is attached a non-increasing sequence $(s_n(S))_{n \in \mathbb{N}}$ of non-negative numbers with a view to classifying operators according to the behaviour of $s_n(S)$ as $n \rightarrow \infty$. The approximation numbers are particularly important examples: the n th approximation number of S is defined to be

$$a_n(S) = \inf \|S - F\|,$$

where the infimum is taken over all linear maps $F : X \rightarrow Y$ with rank less than n . These are special cases of the so-called “strict” s -numbers, further examples of which are provided by the Bernstein, Gelfand, Kolmogorov and Mityagin numbers. As might be expected, the results obtained regarding T_0 are especially sharp when $p = q$. In fact, it then turns out that all the strict s -numbers of T_0 coincide, the n th such number $s_n(T_0)$ being given by the formula

$$s_n(T_0) = \frac{(b-a)\gamma_p}{n+1/2}, \text{ where } \gamma_p = \frac{1}{2\pi} p^{1/p'} (p')^{1/p} \sin(\pi/p).$$

Chapter 6 deals with the general case of the operator T given by (1), in which u and v are merely required to satisfy certain integrability conditions. The precision of the results for T_0 is not obtainable for $T : L_p(I) \rightarrow L_p(I)$, but it emerges that if $1 < p < \infty$, then again all the strict s -numbers of T coincide, and that this time the asymptotic formula

$$\lim_{n \rightarrow \infty} ns_n(T) = \gamma_p \int_a^b |u(t)v(t)| dt$$

holds, where $s_n(T)$ denote the common value of the n th strict s -number of T . The cases $p = 1$ and ∞ present particular difficulties, but even then upper and lower estimates for the approximation numbers of T are obtained. The next chapter develops the theme of Chap. 6: it includes the derivation of more precise asymptotic information about the strict s -numbers of T , given additional restrictions on u and v .

So far, knowledge of the behaviour of the s -numbers of T has been obtained only for the case in which T acts from $L_p(I)$ to itself. When T is viewed as a map from $L_p(I)$ to $L_q(I)$ and $p \neq q$, special problems arise and new techniques are required. Chapter 8 deals with this situation and obtains results by consideration of the variational problem of determining

$$\sup_{g \in T(B)} \|g\|_q,$$

where B is the closed unit ball in $L_p(I)$. When $1 < q < p < \infty$, the asymptotic behaviour of the approximation numbers and the Kolmogorov numbers is established: thus

$$\lim_{n \rightarrow \infty} na_n(T) = C(p, q) \left(\int_a^b |u(t)v(t)|^r dt \right)^{1/r},$$

where $C(p, q)$ is an explicitly known function of p and q , and $r = 1/q + 1/p'$. Moreover, when $1 < p < q < \infty$, a corresponding formula is shown to hold for the Bernstein numbers of T . In both cases connections are made between the s -numbers of T and ‘eigenvalues’ of the variational problem mentioned above. We stress the key rôle played in the arguments presented in Chaps. 5–8 by the generalised trigonometric functions; Chap. 8 also uses more topological ideas, such as the Borsuk antipodal theorem.

The final chapter extends the discussion of the Hardy operator to the situation in which it acts on spaces with variable exponent, the $L_{p(\cdot)}$ spaces. Here p is a given function with values in $(1, \infty)$: if p is a constant function the space coincides with the usual L_p space. Such spaces have attracted a good deal of interest lately because they occur naturally in various physical contexts and in variational problems involving integrands with non-standard growth properties.

Contents

1	Basic Material	1
1.1	Bases and Trigonometric Functions	1
1.2	Strict and Uniform Convexity	3
1.3	Representations of Compact Linear Operators	20
2	Trigonometric Generalisations	33
2.1	The Functions \sin_p and \cos_p	33
2.2	Basis Properties	43
3	The Laplacian and Some Natural Variants	49
3.1	The Laplacian	49
3.2	The p -Laplacian	50
3.3	Eigenfunctions of the p -Laplacian, $n = 1$	55
3.4	Eigenfunctions of the p -Laplacian, $n > 1$	61
4	Hardy Operators	65
4.1	Conditions for Boundedness and Compactness	65
4.2	The Norm of the Hardy Operator	67
5	s-Numbers and Generalised Trigonometric Functions	73
5.1	s -Numbers and n -Widths	73
5.2	The s -Numbers of Hardy Operators	83
5.3	s -Numbers of the Sobolev Embedding on Intervals	95
6	Estimates of s-Numbers of Weighted Hardy Operators	105
6.1	Introduction and Basic Notation	105
6.2	Properties of \mathcal{A}	107
6.3	Equivalence of Strict s -Numbers for T	112
6.4	The First Asymptotic Term when $1 < p < \infty$	115
6.5	The Cases $p = \infty$ and $p = 1$	118

7	More Refined Estimates	129
7.1	Remainder Estimates	129
7.2	The Second Asymptotic Term	138
7.3	l_q and Weak- l_q Estimates when $1 < p < \infty$	141
7.4	l_q and Weak- l_q Estimates, Cases $p = 1$ and ∞	149
8	A Non-Linear Integral System	153
8.1	Upper and Lower Estimates for a Non-Linear Integral System	153
8.2	The Case $q \leq p$	165
8.3	The Case $p \leq q$	175
8.4	The Case $p = q$	182
9	Hardy Operators on Variable Exponent Spaces	183
9.1	Spaces with Variable Exponent	183
9.2	Hardy Operators	192
9.2.1	The Case when p is a Step-Function	196
9.2.2	The Case when p is Strongly Log-Hölder-Continuous	198
9.3	A Version of the p -Laplacian	207
	References	211
	Index	217

Basic Notation

\mathbb{N} : Set of all natural numbers

$\mathbb{N}_0 = \mathbb{N} \cup \{0\}$

\mathbb{Z} : Set of all integers

\mathbb{R} : Set of all real numbers

\mathbb{R}_+ : Set of all non-negative real numbers

\mathbb{R}^n : n -dimensional Euclidean space

If $\Omega \subset \mathbb{R}^n$, then $\overline{\Omega}$ is the closure of Ω , $\overset{o}{\Omega}$ is the interior of Ω , $\partial\Omega$ is the boundary of Ω

$F \sim G$: G is bounded above and below by positive multiples of F independent of any variables occurring in F, G