

Editors:

J.-M. Morel, Cachan

B. Teissier, Paris

For further volumes:

<http://www.springer.com/series/304>

Fabien Morel

\mathbb{A}^1 -Algebraic Topology over a Field

Fabien Morel
Mathematisches Institut der LMU
München, Germany

ISBN 978-3-642-29513-3 ISBN 978-3-642-29514-0 (eBook)
DOI 10.1007/978-3-642-29514-0
Springer Heidelberg New York Dordrecht London

Lecture Notes in Mathematics ISSN print edition: 0075-8434
ISSN electronic edition: 1617-9692

Library of Congress Control Number: 2012942079

Mathematics Subject Classification (2010): 14-XX, 19-XX, 55-XX

© Springer-Verlag Berlin Heidelberg 2012

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media (www.springer.com)

Preface

This work should be considered as a natural sequel to the foundational paper [59] where the \mathbb{A}^1 -homotopy category of smooth schemes over a base scheme was defined and its first properties studied. In this text the base scheme will always be the spectrum of a perfect field k .

One of our first motivations is to emphasize that, contrary to the first impression, the relationship between the \mathbb{A}^1 -homotopy theory over k and the category Sm_k of smooth k -schemes is of the same nature as the relationship between the classical homotopy theory and the category of differentiable manifolds. This explains the title of this work; we hope to convince the reader in this matter. This slogan was already discussed in [55], see also [4].

This text is the result of the compilation of two preprints “ \mathbb{A}^1 -algebraic topology over a field” and “ \mathbb{A}^1 -homotopy classification of vector bundles over smooth affine schemes” to which we added some recent new stuff, consisting of the two sections: “Geometric *versus* canonical transfers” (Chap. 4) and “The Rost–Schmid complex of a strongly \mathbb{A}^1 -invariant sheaf” (Chap. 5).

The main objective of these new sections was primarily to correctly establish the equivalence between the notions of “strongly \mathbb{A}^1 -invariant” and “strictly \mathbb{A}^1 -invariant” for sheaves of abelian groups, see Theorem 1.16. These new sections appear also to be interesting in their own. As the reader will notice, the introduction of the notion of *strictly \mathbb{A}^1 -invariant sheaves with generalized transfers* in our work on the Friedlander–Milnor conjecture [56–58] is directly influenced from these.

Our treatment of transfers in Chap. 4, which is an adaptation of the original one of Bass and Tate for Milnor K -theory [9], is, we think, clarifying. Besides reaching the structure of strictly \mathbb{A}^1 -invariant sheaves with generalized transfers, we obtain on the way a new proof of the fact [38] that the transfers in Milnor K -theory do not depend on choices of a sequence of generators, see Theorem 4.27 and Remark 4.32. Our proof is in spirit different from the one of Kato [38], as we prove directly by geometric means the independence in the case of two generators.

The construction in Chap. 5 for a general strongly \mathbb{A}^1 -invariant sheaf of abelian groups of its “Rost–Schmid” complex is directly influenced by the work of Rost [68] and its adaptation to Witt groups in [70]. Our philosophy on transfers in this work was to use them as less as possible, and to only use them when they really show up by themselves. For instance we will define in Chap. 3 the sheaves of unramified Milnor K-theory (as well as Milnor–Witt K-theory) without using any transfers. The proof of Lemma 5.24 contains the geometric explanation of the formula for the differential of the “Gersten–Rost–Schmid” complex (Corollary 5.44) and the appearance of transfers in it.

The results and ideas in this work have been used and extended in several different directions. In [16] some very concrete computations on \mathbb{A}^1 -homotopy classes of rational fractions give a nice interpretation of our sheaf theoretic computations of $\pi_1^{\mathbb{A}^1}(\mathbb{P}^1)$. The structure and property of the \mathbb{A}^1 -fundamental group sheaves as well as the associated theory of \mathbb{A}^1 -coverings have been used in [2, 3, 86]. It is also the starting point of [4]. Our result (Chap. 9) concerning the Suslin–Voevodsky construction $Sing_{\bullet}^{\mathbb{A}^1}(SL_n)$, $n \geq 3$, has been generalized to a general split semi-simple group of type not SL_2 in [87] and to the case of SL_2 in [60].

The present work plays also a central role in our approach to the Friedlander–Milnor conjecture [56–58].

Conventions and notations. Everywhere in this work, k denotes a fixed perfect field and Sm_k denotes the category of smooth finite type k -schemes.

We denote by \mathcal{F}_k the category of field extensions $k \subset F$ of k such that F is of finite transcendence degree over k . By a discrete valuation v on $F \in \mathcal{F}_k$ we will always mean one which is trivial on k . We let $\mathcal{O}_v \subset F$ denote its valuation ring, $m_v \subset \mathcal{O}_v$ its maximal ideal, and $\kappa(v)$ its residue field.

For any scheme X and any integer i we let $X^{(i)}$ denote the set of points in X of codimension i . Given a point $x \in X$, $\kappa(x)$ will denote its residue field, and \mathcal{M}_x will denote the maximal ideal of the local ring $\mathcal{O}_{X,x}$.

The category Sm_k is always endowed with the Nisnevich topology [59, 62], unless otherwise explicitly stated. Thus for us “sheaf” always means sheaf in the Nisnevich topology.

We will let Set denote the category of sets, Ab that of abelian groups. A *space* is a simplicial object in the category of sheaves of sets on Sm_k [59]. We will also assume the reader is familiar with the notions and results of *loc. cit.*

We denote by Sm'_k the category of *essentially smooth k -schemes*. For us, an *essentially smooth k -scheme* is a noetherian k -scheme X which is the inverse limit of a left filtering system $(X_\alpha)_\alpha$ with each transition morphism $X_\beta \rightarrow X_\alpha$ being an étale affine morphism between smooth k -schemes (see [29]).

For any $F \in \mathcal{F}_k$ the k -scheme $Spec(F)$ is essentially k -smooth. For each point $x \in X \in Sm_k$, the local scheme $X_x := Spec(\mathcal{O}_{X,x})$ of X at x as well as its henselization $X_x^h := Spec(\mathcal{O}_{X,x}^h)$ are essentially smooth k -schemes.

In the same way the complement of the closed point in $\text{Spec}(\mathcal{O}_{X,x})$ or X_x^h is essentially smooth over k . We will sometime make the abuse of writing “smooth k -scheme” instead of “essentially smooth k -scheme”, if no confusion can arise.

Given a presheaf of sets on Sm_k , that is to say a functor $F: (Sm_k)^{op} \rightarrow \text{Sets}$, and an essentially smooth k -scheme $X = \lim_{\alpha} X_{\alpha}$ we set $F(X) := \text{colimit}_{\alpha} F(X_{\alpha})$. From the results of [29] this is well defined. When $X = \text{Spec}(A)$ is affine we will also simply denote this set by $F(A)$.

Acknowledgements I want to take this opportunity to warmly thank Mike Hopkins and Marc Levine for their interest in this work during the past years when I conceived it; their remarks, comments, and some discussions on and around this subject helped me to improve it very much.

Contents

1	Introduction	1
2	Unramified Sheaves and Strongly \mathbb{A}^1-Invariant Sheaves	15
2.1	Unramified Sheaves of Sets	15
2.2	Strongly \mathbb{A}^1 -Invariant Sheaves of Groups	23
2.3	\mathbb{Z} -Graded Strongly \mathbb{A}^1 -Invariant Sheaves of Abelian Groups	36
3	Unramified Milnor–Witt K-Theories	49
3.1	Milnor–Witt K-Theory of Fields	49
3.2	Unramified Milnor–Witt K -Theories	55
3.3	Milnor–Witt K-Theory and Strongly \mathbb{A}^1 -Invariant Sheaves	70
4	Geometric Versus Canonical Transfers	81
4.1	The Gersten Complex in Codimension 2	81
4.2	Geometric Versus Cohomological Transfers on M_{-n}	96
4.3	Proof of the Main Theorem	102
5	The Rost–Schmid Complex of a Strongly \mathbb{A}^1-Invariant Sheaf	113
5.1	Absolute Transfers and the Rost–Schmid Complex	113
5.2	The Rost–Schmid Complex Is a Complex	124
5.3	Gersten Complex <i>Versus</i> Rost–Schmid Complex	138
6	\mathbb{A}^1-Homotopy Sheaves and \mathbb{A}^1-Homology Sheaves	149
6.1	Strong \mathbb{A}^1 -Invariance of the Sheaves $\pi_n^{\mathbb{A}^1}$, $n \geq 1$	149
6.2	\mathbb{A}^1 -Derived Category and Eilenberg–MacLane Spaces	157
6.3	The Hurewicz Theorem and some of Its Consequences	163
7	\mathbb{A}^1-Coverings, $\pi_1^{\mathbb{A}^1}(\mathbb{P}^n)$ and $\pi_1^{\mathbb{A}^1}(SL_n)$	177
7.1	\mathbb{A}^1 -Coverings, Universal \mathbb{A}^1 -Covering and $\pi_1^{\mathbb{A}^1}$	177
7.2	Basic Computation: $\pi_1^{\mathbb{A}^1}(\mathbb{P}^n)$ and $\pi_1^{\mathbb{A}^1}(SL_n)$ for $n \geq 2$	184
7.3	The Computation of $\pi_1^{\mathbb{A}^1}(\mathbb{P}^1)$	188

8	\mathbb{A}^1-Homotopy and Algebraic Vector Bundles	199
8.1	\mathbb{A}^1 -Homotopy Classification of Vector Bundles	199
8.2	The Theory of the Euler Class for Affine Smooth Schemes ...	204
8.3	A Result Concerning Stably Free Vector Bundles	206
9	The Affine B.G. Property for the Linear Groups and the Grassmanian	209
9.1	Preliminaries and Recollections on Regularity	209
9.2	Preliminaries and Recollections on Patching Projective Modules	216
9.3	The Affine B-G Properties for the Grassmanian and the General Linear Groups	221
A	The (Affine) B.G. Property for Simplicial Sheaves	227
A.1	Some Recollections on the B.G. Property	227
A.2	The Affine Replacement of a Simplicial Presheaf	230
A.3	The Affine B.G. Property in the Nisnevich Topology	231
A.4	A Technical Result	237
B	Recollection on Obstruction Theory	241
B.1	The Postnikov Tower of a Morphism	241
B.2	Twisted Eilenberg–MacLane Objects	243
B.3	The Obstruction Theory We Need	248
	References	255
	Index	259